

RICHARD CLAY AND SONS, LIMITED,
LONDON AND BUNGAY.

First Printed March, 1889.

Reprinted January and June, 1890; April, 1891; 1892, 1893, 1894

A TEXT-BOOK
OF
EUCLID'S ELEMENTS

BOOKS I.—IV.



BY

H. S. HALL, M.A.

FORMERLY SCHOLAR OF CHRIST'S COLLEGE, CAMBRIDGE

AND

F. H. STEVENS, M.A.

FORMERLY SCHOLAR OF QUEEN'S COLLEGE, OXFORD
MASTERS OF THE MILITARY SIDE, CLIFTON COLLEGE

LONDON
MACMILLAN AND CO.
AND NEW YORK
1894

All Rights Reserved

BOOK III.

	PAGE
DEFINITIONS, &c.,	149
PROPOSITIONS 1—37,	153
NOTE ON THE METHOD OF LIMITS AS APPLIED TO TANGENCY . . .	213

Theorems and Examples of Book III.

I. ON THE CENTRE AND CHORDS OF A CIRCLE	215
II. ON THE TANGENT AND THE CONTACT OF CIRCLES. The Common Tangent to Two Circles, Problems on Tangency, Orthogonal Circles	217
III. ON ANGLES IN SEGMENTS, AND ANGLES AT THE CENTRES AND CIRCUMFERENCES OF CIRCLES. The Orthocentre of a Triangle, and properties of the Pedal Triangle, Loci, Simson's Line	222
IV. ON THE CIRCLE IN CONNECTION WITH RECTANGLES. Further Problems on Tangency	233
V. ON MAXIMA AND MINIMA	239
VI. HARDER MISCELLANEOUS EXAMPLES	246

BOOK IV.

DEFINITIONS, &c.	250
PROPOSITIONS 1—16	251
NOTE ON REGULAR POLYGONS	274

Theorems and Examples on Book IV.

I. ON THE TRIANGLE AND ITS CIRCLES. Circumscribed, Inscribed, and Escribed Circles, The Nine-points Circle	277
II. MISCELLANEOUS EXAMPLES	283

APPENDIX TO BOOK III.

I. ON POLAR AND POLAR	i
II. ON THE RADICAL AXIS	

CONTENTS.

BOOK I.

	PAGE
DEFINITIONS, POSTULATES, AXIOMS	1
SECTION I. PROPOSITIONS 1—26	11
SECTION II. PARALLELS AND PARALLELOGRAMS.	
PROPOSITIONS 27—31	59
SECTION III. THE AREAS OF PARALLELOGRAMS AND TRIANGLES.	
PROPOSITIONS 35—48	66
<i>Theorems and Examples on Book I.</i>	
ANALYSIS, SYNTHESIS	87
I. ON THE IDENTICAL EQUALITY OF TRIANGLES	90
II. ON INEQUALITIES	93
III. ON PARALLELS	95
IV. ON PARALLELOGRAMS	96
V. MISCELLANEOUS THEOREMS AND EXAMPLES	109
VI. ON THE CONCURRENCE OF STRAIGHT LINES IN A TRI- ANGLE	102
VII. ON THE CONSTRUCTION OF TRIANGLES WITH GIVEN PARTS	107
VIII. ON AREAS	109
IX. ON LOCI	114
X. ON THE INTERSECTION OF LOCI	117

BOOK II.

DEFINITIONS, &c.	129
PROPOSITIONS 1—11	122
THEOREMS AND EXAMPLES ON BOOK II.	141

press, and especially to the Rev. H. C. Watson, of Clifton College, who added to these services much kind assistance in the revision of proof-sheets.

H. S. HALL,

F. H. STEVENS.

February, 1889.

PREFACE.

THIS edition of the first Four Books of Euclid's 'Elements' contains in addition to the text a large and carefully chosen collection of exercises together with Appendices giving the most important elementary developments of Euclidean Geometry.

The text has been carefully revised, and special attention given to those points which experience has shown to present difficulties to beginners.

In the course of this revision the enunciations have been altered as little as possible; and very few departures have been made from Euclid's proofs: in each case changes have been adopted only where the old text has been generally found a cause of difficulty; and such changes are for the most part in favour of well-recognised alternatives.

For example, the ambiguity has been removed from the Enunciations of Propositions 18 and 19 of Book I.: the fact that Propositions 8 and 26 establish the complete

identical equality of the two triangles considered has been strongly urged; and thus the redundant step has been removed from Proposition 34. In Book II. Simson's arrangement of Proposition 13 has been abandoned for a well-known alternative proof. In Book III. Proposition 35 is not given at length, and its place is taken by a simple equivalent. Propositions 35 and 36 have been treated generally, and it has not been thought necessary to do more than call attention in a note to the special cases.

These are the chief deviations from the ordinary text as regards method and arrangement of proof: they are points familiar as difficulties to most teachers, and to name them indicates sufficiently, without further enumeration, the general principles which have guided our revision.

A few alternative proofs of difficult propositions are given for the convenience of those teachers who care to use them.

One purpose of the book is to gradually familiarise the student with the use of legitimate symbols and abbreviations; for a geometrical argument may thus be thrown into a form which is not only more readily seized by an advanced reader, but is useful as a guide to the way in which Euclid's propositions may be handled in written work. On the other hand, we think it very desirable to defer the introduction of symbols until the beginner has learnt that they can only be properly used in Pure Geometry as abbreviations for verbal argument: and we hope thus to prevent the slovenly and inaccurate habits which are very apt to arise from their employment before this principle is fully recognised.

Accordingly in Book I. we have used no contractions or symbols of any kind, though we have introduced verbal

alterations into the text wherever it appeared that conciseness or clearness would be gained.

In Book II. abbreviated forms of constantly recurring words are used, and the phrases *therefore* and *is equal to* are replaced by the usual symbols.

In the Third and Fourth Books, and in additional matter throughout the whole, we have employed all such signs and abbreviations as we believe to add to the clearness of the reasoning, care being taken that the symbols chosen are compatible with a rigorous geometrical method, and are recognised by the majority of teachers.

It must be understood that our use of symbols, and the removal of unnecessary verbiage and repetition, by no means implies a desire to secure brevity at all hazards. On the contrary, nothing appears to us more mischievous than an abridgement which is attained by omitting steps, or condensing two or more steps into one. Such uses spring from the pressure of examinations; but an examination is not, or ought not to be, a mere race; and while we wish to indicate generally in the latter books how a geometrical argument may be abbreviated for the purposes of written work, we have not thought well to reduce the propositions to the bare skeleton so often presented to an Examiner. Indeed it does not follow that the form most suitable for the page of a text-book is also best adapted to examination purposes; for the object to be attained in each case is entirely different. The text-book should present the argument in the clearest possible manner to the mind of a reader to whom it is new: the written proposition need only convey to the Examiner the assurance that the proposition has been thoroughly grasped and remembered by the pupil.

From first to last we have kept in mind the undoubted

fact that a very small proportion of those who study Elementary Geometry, and study it with profit, are destined to become mathematicians in any real sense; and that to a large majority of students, Euclid is intended to serve not so much as a first lesson in *mathematical* reasoning, as the first, and sometimes the only, model of formal and rigid argument presented in an elementary education.

This consideration has determined not only the full treatment of the earlier Books, but the retention of the formal, if somewhat cumbrous, methods of Euclid in many places where proofs of greater brevity and mathematical elegance are available.

We hope that the additional matter introduced into the book will provide sufficient exercise for pupils whose study of Euclid is preliminary to a mathematical education.

The questions distributed through the text follow very easily from the propositions to which they are attached, and we think that teachers are likely to find in them all that is needed for an average pupil reading the subject for the first time.

The Theorems and Examples at the end of each Book contain questions of a slightly more difficult type: they have been very carefully classified and arranged, and brought into close connection with typical examples worked out either partially or in full; and it is hoped that this section of the book, on which much thought has been expended, will do something towards removing that extreme want of freedom in solving deductions that is so commonly found even among students who have a good knowledge of the text of Euclid.

In the course of our work we have made ourselves acquainted with most modern English books on Euclidean

Geometry: among these we have to express our special indebtedness to the text-book recently published by the Association for the Improvement of Geometrical Teaching; and we must also mention the Edition of Euclid's Elements prepared by Mr J. S. Mackay, whose historical notes and frequent references to original authorities have been of the utmost service to us.

Our Treatment of Maxima and Minima on page 239 is based upon suggestions derived from a Discussion of the subject which took place at the annual meeting of the Geometrical Association in January 1887.

An appendix containing an elementary account of the properties of Pole and Polar and Radical Axis appears at the end of the volume.

Of the Riders and deductions some are original; but the greater part have been drawn from that large store of floating material which has furnished Examination Papers for the last 30 years, and must necessarily form the basis of any elementary collection. Proofs which have been found in two or more books without acknowledgement have been regarded as common property.


As regards figures, in accordance with a usage not uncommon in recent editions of Euclid, we have made a distinction between given lines and lines of construction.

Throughout the book we have italicised those deductions on which we desired to lay special stress as being in themselves important geometrical results: this arrangement we think will be useful to teachers who have little time to devote to riders, or who wish to sketch out a suitable course for revision.

We have in conclusion to tender our thanks to many of our friends for the valuable criticism and advice which we received from them as the book was passing through the

PROPOSITION 30. THEOREM.

Straight lines which are parallel to the same straight line are parallel to one another.



Let the straight lines AB , CD be each parallel to the straight line PQ :

then shall AB and CD be parallel to one another.

Construction. Draw any straight line EF cutting AB , CD , and PQ in the points G , H , and K .

Proof. Then because AB and PQ are parallel, and EF meets them,

therefore the angle AGK is equal to the alternate angle GKQ .
I. 29.

And because CD and PQ are parallel, and EF meets them, therefore the exterior angle GHD is equal to the interior opposite angle HKG .
I. 29.

Therefore the angle AGH is equal to the angle GHD ;

and these are alternate angles;

therefore AB and CD are parallel. I. 27.

Q.E.D.

NOTE. (If PQ lies between AB and CD , the Proposition may be established in a similar manner, though in this case it scarcely needs proof; for it is inconceivable that two straight lines, which do not meet an intermediate straight line, should meet one another.

The truth of this Proposition may be readily deduced from Playfair's Axiom, of which it is the converse.

For if AB and CD were not parallel, they would meet when produced. Then there would be two intersecting straight lines both parallel to a third straight line: which is impossible.

Therefore AB and CD never meet; that is, they are parallel.)

EUCLID'S ELEMENTS.

BOOK I.

DEFINITIONS.

1. A **point** is that which has position, but no magnitude.

2. A **line** is that which has length without breadth.

The extremities of a line are points, and the intersection of two lines is a point.

3. A **straight line** is that which lies evenly between its extreme points.

Any portion cut off from a straight line is called a **segment** of it.

4. A **surface** is that which has length and breadth, but no thickness.

The boundaries of a surface are lines.

5. A **plane surface** is one in which any two points being taken, the straight line between them lies wholly in that surface.

A plane surface is frequently referred to simply as a plane.

NOTE. Euclid regards a point merely as a *mark of position*, and he therefore attaches to it no idea of size and shape.


Similarly he considers that the properties of a line arise only from its *length* and *position*, without reference to that minute breadth which every line must really have *if actually drawn*, even though the most perfect instruments are used.

The definition of a surface is to be understood in a similar way.


6. " A **plane angle** is the inclination of two straight lines to one another, which meet together, but are not in the same straight line.

The point at which the straight lines meet is called the **vertex** of the angle, and the straight lines themselves the **arms** of the angle.

When several angles are at one point O , any one of them is expressed by three letters, of which the letter that refers to the vertex is put between the other two. Thus if the straight lines OA , OB , OC meet at the point O , the angle contained by the straight lines OA , OB is named the angle AOB or BOA ; and the angle contained by OA , OC is named the angle AOC or COA . Similarly the angle contained by OB , OC is referred to as the angle BOC or COB . But if there be only one angle at a point, it may be expressed by a single letter, as *the angle at O* .



Of the two straight lines OB , OC shewn in the adjoining figure, we recognize that OC is *more inclined* than OB to the straight line OA : this we express by saying that the angle AOC is greater than the angle AOB . Thus an angle must be regarded as having *magnitude*.




It should be observed that the angle AOC is the sum of the angles AOB and BOC ; and that AOB is the difference of the angles AOC and BOC .

The beginner is cautioned against supposing that the size of an angle is altered either by increasing or diminishing the length of its arms.


[Another view of an angle is recognized in many branches of mathematics; and though not employed by Euclid, it is here given because it furnishes more clearly than any other a conception of what is meant by the *magnitude* of an angle.

Suppose that the straight line OP in the figure is capable of revolution about the point O , like the hand of a watch, but in the opposite direction; and suppose that in this way it has passed successively from the position OA to the positions occupied by OB and OC .




Such a line must have undergone *more turning* in passing from OA to OC , than in passing from OA to OB ; and consequently the angle AOC is said to be greater than the angle AOB .]


7. When a straight line standing on another straight line makes the adjacent angles equal to one another, each of the angles is called a **right angle**; and the straight line which stands on the other is called a **perpendicular** to it.




8. An **obtuse angle** is an angle which is greater than one right angle, but less than two right angles.




9. An **acute angle** is an angle which is less than a right angle.



[In the adjoining figure the straight line **OB** may be supposed to have arrived at its present position, from the position occupied by **OA**, by revolution about the point **O** in either of the two directions indicated by the arrows: thus two straight lines drawn from a point may be considered as forming two angles, (marked (i) and (ii) in the figure) of which the greater (ii) is said to be **reflex**.]



If the arms **OA**, **OB** are in the same straight line, the angle formed by them on either side is called a **straight angle**.]




10. Any portion of a plane surface bounded by one or more lines, straight or curved, is called a **plane figure**.

The sum of the bounding lines is called the **perimeter** of the figure.

Two figures are said to be equal in area, when they enclose equal portions of a plane surface.

✓ 11. A **circle** is a plane figure contained by one line, which is called the **circumference**, and is such that all straight lines drawn from a certain point within the figure to the circumference are equal to one another: this point is called the **centre** of the circle.



A **radius** of a circle is a straight line drawn from the centre to the circumference.

12. A **diameter** of a circle is a straight line drawn through the centre, and terminated both ways by the circumference.

13. A **semicircle** is the figure bounded by a diameter of a circle and the part of the circumference cut off by the diameter.

14. A **segment of a circle** is the figure bounded by a straight line and the part of the circumference which it

15. **Rectilineal figures** are those which are bounded by straight lines.

16. A **triangle** is a plane figure bounded by *three* straight lines.

Any one of the angular points of a triangle may be regarded as its **vertex**; and the opposite side is then called the **base**.

17. A **quadrilateral** is a plane figure bounded by *four* straight lines.

The straight line which joins opposite angular points in a quadrilateral is called a **diagonal**.

18. A **polygon** is a plane figure bounded by more than four straight lines.


19. An **equilateral triangle** is a triangle whose three sides are equal.



20. An **isosceles triangle** is a triangle two of whose sides are equal.



21. A **scalene triangle** is a triangle which has three unequal sides.



22. A **right-angled triangle** is a triangle which has a right angle.



The side opposite to the right angle in a right-angled triangle is called the **hypotenuse**.

23. An **obtuse-angled triangle** is a triangle which has an obtuse angle.




24. An **acute-angled triangle** is a triangle which has *three* acute angles.




[It will be seen hereafter (Book I. Proposition 17) *that every triangle must have at least two acute angles.*]

25. ✓ **Parallel straight lines** are such as, being in the same plane, do not meet, however far they are produced in either direction.


26. A **Parallelogram** is a four-sided figure which has its opposite sides parallel.



27. A **rectangle** is a parallelogram which has one of its angles a right angle.



28. A **square** is a four-sided figure which has all its sides equal and all its angles right angles.



[It may easily be shewn that if a quadrilateral has all its sides equal and *one* angle a right angle, then *all* its angles will be right angles.]

29. ✓ A **rhombus** is a four-sided figure which has all its sides equal, but its angles are not right angles.



30. ✓ A **trapezium** is a four-sided figure which has *two* of its sides parallel.



ON THE POSTULATES.

In order to effect the *constructions* necessary to the study of geometry, it must be supposed that certain instruments are available; but it has always been held that such instruments should be as few in number, and as simple in character as possible.

For the purposes of the first Six Books a *straight ruler* and a pair of compasses are all that are needed; and in the following **Postulates**, or requests, Euclid demands the use of such instruments, and assumes that they suffice, theoretically as well as practically, to carry out the processes mentioned below.

POSTULATES. *man's requests*

Let it be granted,

1. That a straight line may be drawn from any one point to any other point.

When we draw a straight line from the point A to the point B, we are said to *join AB*.

2. That a *finite*, that is to say, a terminated straight line may be produced to any length in that straight line.

3. That a circle may be described from any centre, at any distance from that centre, that is, with a radius equal to any finite straight line drawn from the centre.

It is important to notice that the Postulates include no means of *direct measurement*: hence the straight ruler is not supposed to be *graduated*; and the compasses, in accordance with Euclid's use, are not to be employed for *transferring distances* from one part of a figure to another.

ON THE AXIOMS.

The science of Geometry is based upon certain simple statements, the truth of which is assumed at the outset to be self-evident.

These self-evident truths, called by Euclid *Common Notions*, are now known as the **Axioms**.

✓ The necessary characteristics of an Axiom are

(i) That it should be *self-evident*; that is, that its truth should be immediately accepted without proof.

(ii) That it should be *fundamental*; that is, that its truth should not be derivable from any other truth more simple than itself.

(iii) That it should supply a basis for the establishment of further truths.

These characteristics may be summed up in the following definition.

✓ (DEFINITION. An **Axiom** is a self-evident truth, which neither requires nor is capable of proof, but which serves as a foundation for future reasoning.)

Axioms are of two kinds, *general* and *geometrical*.

General Axioms apply to *magnitudes of all kinds*. Geometrical Axioms refer exclusively to *geometrical magnitudes*, such as have been already indicated in the definitions.

GENERAL AXIOMS.

1. Things which are equal to the same thing are equal to one another.

2. If equals be added to equals, the wholes are equal.*

3. If equals be taken from equals, the remainders are equal.

4. If equals be added to unequals, the wholes are unequal, the greater sum being that which includes the greater of the unequals.

5. If equals be taken from unequals, the remainders are unequal, the greater remainder being that which is left from the greater of the unequals.

6. Things which are double of the same thing, or of equal things, are equal to one another.

7. Things which are halves of the same thing, or of equal things, are equal to one another.

9.* The whole is greater than its part.

* To preserve the classification of general and geometrical axioms, we have placed Euclid's *ninth* axiom before the *eighth*.

GEOMETRICAL AXIOMS.

8. Magnitudes which can be made to coincide with one another, are equal.

This axiom affords the ultimate test of the equality of two geometrical magnitudes. It implies that any line, angle, or figure, may be supposed to be taken up from its position, and without change in size or form, laid down upon a second line, angle, or figure, for the purpose of comparison.

This process is called **superposition**, and the first magnitude is said to be **applied** to the other.


10. Two straight lines cannot enclose a space.

11. All right angles are equal.

[The statement that all right angles are equal, admits of proof, and is therefore perhaps out of place as an Axiom.]

12. If a straight line meet two straight lines so as to make the interior angles on one side of it together less than two right angles, these straight lines will meet if continually produced on the side on which are the angles which are together less than two right angles.

That is to say, if the two straight lines AB and CD are met by the straight line EH at F and G , in such a way that the angles BFG , DGF are together less than two right angles, it is asserted that AB and CD will meet if continually produced in the direction of B and D .



[Axiom 12 has been objected to on the double ground that it cannot be considered self-evident, and that its truth may be deduced from simpler principles. It is employed for the first time in the 29th Proposition of Book I., where a short discussion of the difficulty will be found.

The converse of this Axiom is proved in Book I. Proposition 17.]

INTRODUCTORY.

Plane Geometry deals with the properties of all lines and figures that may be drawn upon a plane surface.

Euclid in his first Six Books confines himself to the properties of straight lines, rectilineal figures, and circles.

The *Definitions* indicate the subject-matter of these books: the *Postulates and Axioms* lay down the fundamental principles which regulate all investigation and argument relating to this subject-matter.

Euclid's method of exposition divides the subject into a number of separate discussions, called **propositions**; each proposition, though in one sense complete in itself, is derived from results previously obtained, and itself leads up to subsequent propositions.

Propositions are of two kinds, **Problems** and **Theorems**.

(A **Problem** proposes to effect some geometrical construction, such as to draw some particular line, or to construct some required figure.)

(A **Theorem** proposes to demonstrate some geometrical truth.)

A Proposition consists of the following parts:

The General Enunciation, the Particular Enunciation, the Construction, and the Demonstration or Proof.

(1) The **General Enunciation** is a preliminary statement, describing in general terms the purpose of the proposition.

In a *problem* the Enunciation states the construction which it is proposed to effect: it therefore names first the **Data**, or things given, secondly the **Quæsitæ**, or things required.

In a *theorem* the Enunciation states the property which it is proposed to demonstrate: it names first, the **Hypothesis**, or the conditions assumed; secondly, the **Conclusion**, or the assertion to be proved.

(ii) The **Particular Enunciation** repeats in special terms the statement already made, and refers it to a diagram, which enables the reader to follow the reasoning more easily. }

(iii) The **Construction** then directs the drawing of such straight lines and circles as may be required to effect the purpose of a problem, or to prove the truth of a theorem. }

(iv) Lastly, the **Demonstration** proves that the object proposed in a problem has been accomplished, or that the property stated in a theorem is true. }

Euclid's reasoning is said to be **Deductive**, because by a connected chain of argument it **deduces** new truths from truths already proved or admitted.

The initial letters Q. E. F., placed at the end of a problem, stand for **Quod erat Faciendum**, *which was to be done*.

The letters Q. E. D. are appended to a theorem, and stand for **Quod erat Demonstrandum**, *which was to be proved*.

A **Corollary** is a statement the truth of which follows readily from an established proposition; it is therefore appended to the proposition as an inference or deduction, which usually requires no further proof.

The following symbols and abbreviations may be employed in writing out the propositions of Book I., though their use is not recommended to beginners.


\therefore	for	therefore,	par ^l (or)	for	parallel,
$=$	„	is, or are, equal to,	par ^m	„	parallelogram,
\angle	„	angle,	sq.	„	square,
rt. \angle	„	right angle,	rectil.	„	rectilineal,
Δ	„	triangle,	st. line	„	straight line,
perp.	„	perpendicular,	pt.	„	a point;

and all obvious contractions of words, such as opp., adj., diag., &c., for opposite, adjacent, diagonal, &c.

SECTION I.

PROPOSITION 1. PROBLEM.

To describe an equilateral triangle on a given finite straight line.



Let AB be the given straight line.

It is required to describe an equilateral triangle on AB .

Construction. From centre A , with radius AB , describe the circle BCD . *Post. 3.*

From centre B , with radius BA , describe the circle ACE . *Post. 3.*

From the point C at which the circles cut one another, draw the straight lines CA and CB to the points A and B . *Post. 1.*

Then shall ABC be an equilateral triangle.

Proof. Because A is the centre of the circle BCD ,
therefore AC is equal to AB . *Def. 11.*

And because B is the centre of the circle ACE ,
therefore BC is equal to BA . *Def. 11.*

But it has been shewn that AC is equal to AB ;

therefore AC and BC are each equal to AB .

But things which are equal to the same thing are equal to one another. *Ax. 1.*


Therefore AC is equal to BC .

Therefore CA , AB , BC are equal to one another.

Therefore the triangle ABC is equilateral;
and it is described on the given straight line AB . *Q. E. F.*

PROPOSITION 2. PROBLEM.

From a given point to draw a straight line equal to a given straight line.



Let *A* be the given point, and *BC* the given straight line.
It is required to draw from the point *A* a straight line equal to *BC*.

Construction. Join *AB*; *Post.* 1.

and on *AB* describe an equilateral triangle *DAB*. 1. 1.

From centre *B*, with radius *BC*, describe the circle *CGH*.

Post. 3.

Produce *DB* to meet the circle *CGH* at *G*. *Post.* 2.

From centre *D*, with radius *DG*, describe the circle *GKF*.

Produce *DA* to meet the circle *GKF* at *F*. *Post.* 2.

Then *AF* shall be equal to *BC*.

Proof. Because *B* is the centre of the circle *CGH*,
therefore *BC* is equal to *BG*. *Def.* 11.

And because *D* is the centre of the circle *GKF*,
therefore *DF* is equal to *DG*; *Def.* 11.

and *DA*, *DB*, parts of them are equal; *Def.* 19.
therefore the remainder *AF* is equal to the remainder *BG*.
Ax. 3.

And it has been shewn that *BC* is equal to *BG*;

therefore *AF* and *BC* are each equal to *BG*.

But things which are equal to the same thing are equal to one another. *Ax.* 1.


Therefore *AF* is equal to *BC*;

and it has been drawn from the given point *A*. Q. E. F.

[This Proposition is rendered necessary by the restriction, tacitly imposed by Euclid, that compasses shall not be used to *transfer distances*.]

PROPOSITION 3. PROBLEM.

From the greater of two given straight lines to cut off a part equal to the less.



Let AB and C be the two given straight lines, of which AB is the greater.

It is required to cut off from AB a part equal to C.

Construction. From the point A draw the straight line AD equal to C;
and from centre A, with radius AD, describe the circle DEF, meeting AB at E. I. 2.
Post. 3.

Then AE shall be equal to C.

Proof. Because A is the centre of the circle DEF,
therefore AE is equal to AD. Def. 11.

But C is equal to AD. Constr.

Therefore AE and C are each equal to AD.

Therefore AE is equal to C;

and it has been cut off from the given straight line AB.

Q. E. D.

EXERCISES.

1. On a given straight line describe an isosceles triangle having each of the equal sides equal to a given straight line.

2. On a given base describe an isosceles triangle having each of the equal sides double of the base.


3. In the figure of 1. 2, if AB is equal to BC, shew that D, the vertex of the equilateral triangle, will fall on the circumference of the circle CGH.

Obs. Every triangle has six **parts**, namely its three sides and three angles.)

Two triangles are said to be **equal in all respects**, when they can be made to coincide with one another by *superposition* (see note on Axiom 8), and in this case each part of the one is equal to a corresponding part of the other.

PROPOSITION 4. THEOREM.

If two triangles have two sides of the one equal to two sides of the other, each to each, and have also the angles contained by those sides equal: then shall their bases or third sides be equal, and the triangles shall be equal in area, and their remaining angles shall be equal, each to each, namely those to which the equal sides are opposite: that is to say, the triangles shall be equal in all respects.



Let ABC , DEF be two triangles, which have the side AB equal to the side DE , the side AC equal to the side DF , and the contained angle BAC equal to the contained angle EDF .

Then shall the base BC be equal to the base EF , and the triangle ABC shall be equal to the triangle DEF in area; and the remaining angles shall be equal, each to each, to which the equal sides are opposite,

namely the angle ABC to the angle DEF ,
and the angle ACB to the angle DFE .

For if the triangle ABC be applied to the triangle DEF ,
so that the point A may be on the point D ,

and the straight line AB along the straight line DE ,

then because AB is equal to DE ,

Hyp.

therefore the point B must coincide with the point E .

And because AB falls along DE,
and the angle BAC is equal to the angle EDF, *Hyp.*
therefore AC must fall along DF.

And because AC is equal to DF, *Hyp.*
therefore the point C must coincide with the point F.

Then B coinciding with E, and C with F,
the base BC must coincide with the base EF;
for if not, two straight lines would enclose a space; which
is impossible. *Ax. 10.*

Thus the base BC coincides with the base EF, and is
therefore equal to it. *Ax. 3.*

And the triangle ABC coincides with the triangle DEF,
and is therefore equal to it in area. *Ax. 8.*

And the remaining angles of the one coincide with the re-
maining angles of the other, and are therefore equal to them,
namely, the angle ABC to the angle DEF,
and the angle ACB to the angle DFE.

That is, the triangles are equal in all respects. *Q. E. D.*

NOTE. It follows that two triangles which are equal in their
several parts are equal also in *area*; but it should be observed that
equality of area in two triangles does not necessarily imply equality in
their several parts: that is to say, triangles may be equal in *area*,
without being of the same *shape*.

(Two triangles which are equal in all respects have *identity of form
and magnitude*, and are therefore said to be *identically equal*, or
congruent.)

The following application of Proposition 4 anticipates
the chief difficulty of Proposition 5.

In the equal sides AB, AC of an isosceles triangle
ABC, the points X and Y are taken, so that AX
is equal to AY; and XY and CX are joined.

Shew that BY is equal to CX.


In the two triangles XAC, YAB,

XA is equal to YA, and AC is equal to AB; *Hyp.*
that is, the two sides XA, AC are equal to the two
sides YA, AB, each to each;

and the angle at A, which is contained by these
sides, is common to both triangles:

therefore the triangles are equal in all respects;

so that XC is equal to YB.




I. 4.

Q. E. D.

PROPOSITION 5. THEOREM.

The angles at the base of an isosceles triangle are equal to one another; and if the equal sides be produced, the angles on the other side of the base shall also be equal to one another.



Let $\triangle ABC$ be an isosceles triangle, having the side AB equal to the side AC , and let the straight lines AB , AC be produced to D and E :

then shall the angle ABC be equal to the angle ACB , and the angle CBD to the angle BCE .

Construction. In BD take any point F ; and from AE the greater cut off AG equal to AF the less. I. 3.
Join FC , GB .

Proof. Then in the triangles FAC , GAB ,

Because $\begin{cases} FA \text{ is equal to } GA, & \text{Constr.} \\ \text{and } AC \text{ is equal to } AB, & \text{Hyp.} \\ \text{also the contained angle at } A \text{ is common to the} \\ \text{two triangles;} \end{cases}$

therefore the triangle FAC is equal to the triangle GAB in all respects; I. 4.

that is, the base FC is equal to the base GB ,

and the angle ACF is equal to the angle ABG ,

also the angle AFC is equal to the angle AGB .

Again, because the whole AF is equal to the whole AG ,
of which the parts AB , AC are equal, Hyp.

therefore the remainder BF is equal to the remainder CG .

Then in the two triangles BFC , CGB ,

Because $\left\{ \begin{array}{l} BF \text{ is equal to } CG, \text{ } \textit{Proved.} \\ \text{and } FC \text{ is equal to } GB, \text{ } \textit{Proved.} \\ \text{also the contained angle } BFC \text{ is equal to the} \\ \text{contained angle } CGB, \text{ } \textit{Proved.} \end{array} \right.$
therefore the triangles BFC , CGB are equal in all respects;
so that the angle FBC is equal to the angle GCB ,
and the angle BCF to the angle CBG . I. 4.

Now it has been shewn that the whole angle ABG is equal
to the whole angle ACF ,
and that parts of these, namely the angles CBG , BCF , are
also equal;

therefore the remaining angle ABC is equal to the remain-
ing angle ACB ;

and these are the angles at the base of the triangle ABC .

Also it has been shewn that the angle FBC is equal to the
angle GCB ;

and these are the angles on the other side of the base. Q.E.D.

(COROLLARY. *Hence if a triangle is equilateral it is
also equiangular.*)

EXERCISES.

1. AB is a given straight line and C a given point outside it: shew
how to find any points in AB such that their distance from C shall be
equal to a given length L . Can such points always be found?

2. If the vertex C and one extremity A of the base of an isosceles
triangle are given, find the other extremity B , supposing it to lie on a
given straight line PQ .


3. Describe a rhombus having given two opposite angular points
 A and C , and the length of each side.

4. $AMNB$ is a straight line; on AB describe a triangle ABC such
that the side AC shall be equal to AN and the side BC to MB .

5. In Prop. 2 the point A may be joined to *either* extremity of BC .
Draw the figure and prove the proposition in the case when A is joined
to C .

The following proof is sometimes given as a substitute for the first part of Proposition 5.

PROPOSITION 5. ALTERNATIVE PROOF.



Let ABC be an isosceles triangle, having AB equal to AC :
then shall the angle ABC be equal to the angle ACB .

Suppose the triangle ABC to be taken up, turned over and laid down again in the position $A'B'C'$, where $A'B'$, $A'C'$, $B'C'$ represent the new positions of AB , AC , BC .

Then $A'B'$ is equal to $A'C'$; and $A'B'$ is AB in its new position,
therefore AB is equal to $A'C'$;

{ in the same way AC is equal to $A'B'$;

and the included angle BAC is equal to the included angle $C'A'B'$, for
they are the same angle in different positions ;

therefore the triangle ABC is equal to the triangle $A'C'B'$ in all respects :

so that the angle ABC is equal to the angle $A'C'B'$. I. 4.

But the angle $A'C'B'$ is the angle ACB in its new position ;

therefore the angle ABC is equal to the angle ACB .

Q. E. D.

EXERCISES.

CHIEFLY ON PROPOSITIONS 4 AND 5.

1. Two circles have the same centre O ; OAD and OBE are straight lines drawn to cut the smaller circle in A and B and the larger circle in D and E : prove that

(i) $AD = BE$.

(ii) $DB = EA$.

(iii) The angle DAB is equal to the angle EBA .

(iv) The angle ODB is equal to the angle OEA .

2. $ABCD$ is a square, and L , M , and N are the middle points of AB , BC , and CD : prove that

(i) $LM = LN$.

(ii) $AM = DM$.

(iii) $AN \perp AL$.

(iv) $BN = DM$.

[Draw a separate figure in each case].

3. O is the centre of a circle and OA , OB are radii ; OM divides the angle AOB into two equal parts and cuts the line AB in M : prove that $AM = BM$.

4. ABC , DBC are two isosceles triangles described on the same base BC but on opposite sides of it : prove that the angle ABD is equal to the angle ACD .

5. ABC , DBC are two isosceles triangles described on the same base BC , but on opposite sides of it : prove that if AD be joined, each of the angles BAC , BDC will be divided into two equal parts.

6. PQR , SQR are two isosceles triangles described on the same base QR , and on the same side of it : shew that the angle PQS is equal to the angle PRS , and that the line PS divides the angle QPR into two equal parts.

7. If in the figure of Exercise 5 the line AD meets BC in E , prove that $BE = EC$.

8. $ABCD$ is a rhombus and AC is joined : prove that the angle DAB is equal to the angle DCB .

9. $ABCD$ is a quadrilateral having the opposite sides BC , AD equal, and also the angle BCD equal to the angle ADC : prove that BD is equal to AC .

10. AB , AC are the equal sides of an isosceles triangle ; L , M , N are the middle points of AB , BC , and CA respectively : prove that $LM = MN$.

Prove also that the angle ALM is equal to the angle ANM .

• DEFINITION. (Each of two Theorems is said to be the **Converse** of the other, when the hypothesis of each is the conclusion of the other.)


(It will be seen, on comparing the hypotheses and conclusions of Props. 5 and 6, that each proposition is the converse of the other.)

NOTE. (Proposition 6 furnishes the first instance of an *indirect method of proof*, frequently used by Euclid. It consists in shewing that an absurdity must result from supposing the theorem to be otherwise than true. This form of demonstration is known as the **Reductio ad Absurdum**, and is most commonly employed in establishing the converse of some foregoing theorem.)

It must not be supposed that the converse of a true theorem is itself necessarily true : for instance, it will be seen in Prop. 8, C, that if two triangles have their sides equal, each to each, then their angles will also be equal, each to each ; but it may easily be shewn by means of a figure that the converse of this theorem is not necessarily true.

PROPOSITION 6. THEOREM.

If two angles of a triangle be equal to one another, then the sides also which subtend, or are opposite to, the equal angles, shall be equal to one another.



Let ABC be a triangle, having the angle ABC equal to the angle ACB :

then shall the side AC be equal to the side AB .

Construction. For if AC be not equal to AB , one of them must be greater than the other.

If possible, let AB be the greater ;
and from it cut off BD equal to AC . 1. 3.
Join DC .


Proof. Then in the triangles DBC , ACB ,
Because $\left\{ \begin{array}{l} DB \text{ is equal to } AC, \\ \text{and } BC \text{ is common to both,} \\ \text{also the contained angle } DBC \text{ is equal to the} \\ \text{contained angle } ACB ; \end{array} \right. \begin{array}{l} \text{Constr.} \\ \text{Hyp.} \end{array}$
therefore the triangle DBC is equal in area to the triangle ACB , 1. 4.
the part equal to the whole ; which is absurd. Ax. 9.

Therefore AB is not unequal to AC ;
that is, AB is equal to AC . Q.E.D.

COROLLARY. (*Hence if a triangle is equiangular it is also equilateral.*)

PROPOSITION 7. THEOREM.

On the same base, and on the same side of it, there cannot be two triangles having their sides which are terminated at one extremity of the base equal to one another, and likewise those which are terminated at the other extremity equal to one another.



If it be possible, on the same base AB , and on the same side of it, let there be two triangles ACB , ADB , having their sides AC , AD , which are terminated at A , equal to one another, and likewise their sides BC , BD , which are terminated at B , equal to one another.

CASE I. " When the vertex of each triangle is without the other triangle.

Construction. Join CD . *Post.* 1.


Proof. Then in the triangle ACD ,
because AC is equal to AD , *Hyp.*
therefore the angle ACD is equal to the angle ADC . 1. 5.

But the whole angle ACD is greater than its part, the angle BCD ,

therefore also the angle ADC is greater than the angle BCD ;
still more then is the angle BDC greater than the angle BCD .

Again, in the triangle BCD ,
because BC is equal to BD , *Hyp.*
therefore the angle BDC is equal to the angle BCD : 1. 5.
but it was shewn to be greater; which is impossible.

CASE II. When one of the vertices, as D, is within the other triangle, ACE.



Construction. As before, join CD; Post. 1.
and produce AC, AD to E and F. Post. 2.

Then in the triangle ACD, because AC is equal to AD, *Hyp.*
therefore the angles ECD, FDC, on the other side of the
base, are equal to one another. I. 5.

But the angle ECD is greater than its part, the angle BCD;
therefore the angle FDC is also greater than the angle
BCD:

still more than is the angle BDC greater than the angle
BCD.

Again, in the triangle BCD,
because BC is equal to BD, *Hyp.*
therefore the angle BDC is equal to the angle BCD: I. 5.
but it has been shewn to be greater; which is impossible.


The case in which the vertex of one triangle is on a
side of the other needs no demonstration.

Therefore AC cannot be equal to AD, and at the same
time, BC equal to BD. Q.E.D.

NOTE. The sides AC, AD are called **conterminous** sides; similarly
the sides BC, BD are conterminous.

PROPOSITION 8. THEOREM.

*If two triangles have two sides of the one equal to two
sides of the other, each to each, and have likewise their bases
equal, then the angle which is contained by the two sides of
the one shall be equal to the angle which is contained by
the two sides of the other.*



Let ABC , DEF be two triangles, having the two sides BA , AC equal to the two sides ED , DF , each to each, namely BA to ED , and AC to DF , and also the base BC equal to the base EF :

then shall the angle BAC be equal to the angle EDF .

Proof. For if the triangle ABC be applied to the triangle DEF , so that the point B may be on E , and the straight line BC along EF ;

then because BC is equal to EF , *Hyp.*
therefore the point C must coincide with the point F .

Then, BC coinciding with EF ,
it follows that BA and AC must coincide with ED and DF :
for if not, they would have a different situation, as EG , GF :
then, on the same base and on the same side of it there
would be two triangles having their *conterminous* sides
equal.

But this is impossible. I. 7.
Therefore the sides BA , AC coincide with the sides ED , DF .
That is, the angle BAC coincides with the angle EDF , and is
therefore equal to it. *Ax. 8.*

Q. E. D.


NOTE. In this Proposition the three sides of one triangle are given equal respectively to the three sides of the other; and from this it is shewn that the two triangles may be made to coincide with one another.

Hence we are led to the following important Corollary.

COROLLARY. *If in two triangles the three sides of the one are equal to the three sides of the other, each to each, then the triangles are equal in all respects.*

The following proof of Prop. 8 is worthy of attention as it is independent of Prop. 7, which frequently presents difficulty to a beginner.

✓ PROPOSITION 8. ALTERNATIVE PROOF.



Let ABC and DEF be two triangles, which have the sides BA, AC equal respectively to the sides ED, DF , and the base BC equal to the base EF :

then shall the angle BAC be equal to the angle EDF .

For apply the triangle ABC to the triangle DEF , so that B may fall on E , and BC along EF , and so that the point A may be on the side of EF remote from D .

then C must fall on F , since BC is equal to EF .

Let $A'EF$ be the new position of the triangle ABC .

If neither DF, FA' nor DE, EA' are in one straight line, join DA' .

CASE I. When DA' intersects EF .

Then because ED is equal to EA' ,

therefore the angle EDA' is equal to the angle $EA'D$. 1. 5.

Again because FD is equal to FA' ,

therefore the angle FDA' is equal to the angle $FA'D$. 1. 5.

Hence the whole angle EDF is equal to the whole angle $EA'F$;

that is, the angle EDF is equal to the angle BAC .


Two cases remain which may be dealt with in a similar manner: namely,

CASE II. When DA' meets EF produced.

CASE III. When one pair of sides, as DF, FA' , are in one straight line.

PROPOSITION 9. PROBLEM.

To bisect a given angle, that is, to divide it into two equal parts.



Let BAC be the given angle:
it is required to bisect it.

Construction. In AB take any point D ;
and from AC cut off AE equal to AD . I. 3.

Join DE ;

and on DE , on the side remote from A , describe an equi-
lateral triangle DEF . I. 1.

Join AF .

Then shall the straight line AF bisect the angle BAC .

Proof. For in the two triangles DAF , EAF ,

Because $\left\{ \begin{array}{l} DA \text{ is equal to } EA, \\ \text{and } AF \text{ is common to both;} \\ \text{and the third side } DF \text{ is equal to the third side } EF; \end{array} \right. \begin{array}{l} \text{Constr.} \\ \\ \text{Def. 19.} \end{array}$

therefore the angle DAF is equal to the angle EAF . I. 8.


Therefore the given angle BAC is bisected by the straight
line AF . Q.E.F.

EXERCISES.

1. (If in the above figure the equilateral triangle DFE were described on the same side of DE as A , what different cases would arise? And under what circumstances would the construction fail?)
2. (In the same figure, shew that AF also bisects the angle DFE .)
3. (Divide an angle into four equal parts.)

PROPOSITION 10. PROBLEM.

To bisect a given finite straight line, that is, to divide it into two equal parts.



Let AB be the given straight line :
it is required to divide it into two equal parts.

Constr. On AB describe an equilateral triangle ABC , I. 1.
and bisect the angle ACB by the straight line CD , meeting
 AB at D . I. 9.

Then shall AB be bisected at the point D .

Proof. For in the triangles ACD , BCD ,
 Because $\left\{ \begin{array}{l} AC \text{ is equal to } BC, \quad \text{Def. 19.} \\ \text{and } CD \text{ is common to both;} \\ \text{also the contained angle } ACD \text{ is equal to the con-} \\ \text{tained angle } BCD; \quad \text{' Constr.} \end{array} \right.$

Therefore the triangles are equal in all respects:

so that the base AD is equal to the base BD . I. 4.

Therefore the straight line AB is bisected at the point D .

Q. E. F.

EXERCISES.

1. Shew that the straight line which bisects the vertical angle of an isosceles triangle, also bisects the base.

2. On a given base describe an isosceles triangle such that the sum of its equal sides may be equal to a given straight line.

PROPOSITION 11. PROBLEM.

To draw a straight line at right angles to a given straight line, from a given point in the same.

A D C E B

Let AB be the given straight line, and C the given point in it.

It is required to draw from the point C a straight line at right angles to AB .

Construction. In AC take any point D ,
and from CB cut off CE equal to CD . I. 3.

On DE describe the equilateral triangle DFE . I. 1.

Join CF .

Then shall the straight line CF be at right angles to AB .

Proof. For in the triangles DCF , ECF ,
Because { DC is equal to EC , *Constr.*
and CF is common to both;
and the third side DF is equal to the third side EF : *Def. 19.*

Therefore the angle DCF is equal to the angle ECF : I. 8.
and these are adjacent angles.

But when a straight line, standing on another straight line, makes the adjacent angles equal to one another, each of these angles is called a right angle; *Def. 7.*

therefore each of the angles DCF , ECF is a right angle.

Therefore CF is at right angles to AB ,
and has been drawn from a point C in it. Q.E.F.

EXERCISE.

In the figure of the above proposition, shew that any point in FC , or FC produced, is equidistant from D and E .

EXERCISES ON PROPOSITIONS 1 TO 12.

1. Shew that the straight line which joins the vertex of an isosceles triangle to the middle point of the base is perpendicular to the base.

2. Shew that the straight lines which join the extremities of the base of an isosceles triangle to the middle points of the opposite sides, are equal to one another.

3. Two given points in the base of an isosceles triangle are equidistant from the extremities of the base: shew that they are also equidistant from the vertex.

4. If the opposite sides of a quadrilateral are equal, shew that the opposite angles are also equal.

5. Any two isosceles triangles XAB , YAB stand on the same base AB : shew that the angle XAY is equal to the angle XBY ; and that the angle AXY is equal to the angle BXY .

6. Shew that the opposite angles of a rhombus are bisected by the diagonal which joins them.

7. Shew that the straight lines which bisect the base angles of an isosceles triangle form with the base a triangle which is also isosceles.

8. ABC is an isosceles triangle having AB equal to AC ; and the angles at B and C are bisected by straight lines which meet at O : shew that OA bisects the angle BAC .

9. Shew that the triangle formed by joining the middle points of the sides of an equilateral triangle is also equilateral.

10. The equal sides BA , CA of an isosceles triangle BAC are produced beyond the vertex A to the points E and F , so that AE is equal to AF ; and FB , EC are joined: shew that FB is equal to EC .

11. Shew that the diagonals of a rhombus bisect one another at right angles.

12. In the equal sides AB , AC of an isosceles triangle ABC two points X and Y are taken, so that AX is equal to AY ; and CX and BY are drawn intersecting in O : shew that

- (i) the triangle BOC is isosceles;
- (ii) AO bisects the vertical angle BAC ;
- (iii) AO , if produced, bisects BC at right angles.



13. Describe an isosceles triangle, having given the base and the length of the perpendicular drawn from the vertex to the base.

14. In a given straight line find a point that is equidistant from two given points.

In what case is this impossible?

PROPOSITION 13. THEOREM.

If one straight line stand upon another straight line, then the adjacent angles shall be either two right angles, or together equal to two right angles.



Let the straight line AB stand upon the straight line DC : then the adjacent angles DBA , ABC shall be either two right angles, or together equal to two right angles.

CASE I. For if the angle DBA is equal to the angle ABC , each of them is a right angle. *Def. 7.*

CASE II. But if the angle DBA is not equal to the angle ABC ;

from B draw BE at right angles to CD . *I. 11.*

Proof. Now the angle DBA is made up of the two angles DBE , EBA ;

to each of these equals add the angle ABC ;
then the two angles DBA , ABC are together equal to the three angles DBE , EBA , ABC . *Ax. 2.*

Again, the angle EBC is made up of the two angles EBA , ABC ;

to each of these equals add the angle DBE .

Then the two angles DBE , EBC are together equal to the three angles DBE , EBA , ABC . *Ax. 2.*

But the two angles DBA , ABC have been shown to be equal to the same three angles;

therefore the angles DBA , ABC are together equal to the angles DBE , EBC . *Ax. 1.*

But the angles DBE , EBC are two right angles; *Constr.*
therefore the angles DBA , ABC are together equal to two right angles. *Q. E. D.*

DEFINITIONS.

(i) The **complement** of an acute angle is its *defect from a right angle*, that is, the angle by which it falls short of a right angle.

Thus two angles are **complementary**, when their sum is a right angle.

(ii) The **supplement** of an angle is its *defect from two right angles*, that is, the angle by which it falls short of two right angles.

Thus two angles are **supplementary**, when their sum is two right angles.

COROLLARY. *Angles which are complementary or supplementary to the same angle are equal to one another.*


EXERCISES.

1. If the two exterior angles formed by producing a side of a triangle both ways are equal, shew that the triangle is isosceles.

2. The bisectors of the adjacent angles which one straight line makes with another contain a right angle.

NOTE. In the adjoining figure AOB is a given angle; and one of its arms AO is produced to C: the adjacent angles AOB, BOC are bisected by OX, OY.

Then OX and OY are called respectively the **internal** and **external bisectors** of the angle AOB.



Hence Exercise 2 may be thus enunciated:


The internal and external bisectors of an angle are at right angles to one another.

3. Shew that the angles AOX and COY are complementary.

4. Shew that the angles BOX and COX are supplementary; and also that the angles AOY and BOY are supplementary.

PROPOSITION 14. THEOREM.

If, at a point in a straight line, two other straight lines, on opposite sides of it, make the adjacent angles together equal to two right angles, then these two straight lines shall be in one and the same straight line.



At the point B in the straight line AB, let the two straight lines BC, BD, on the opposite sides of AB, make the adjacent angles ABC, ABD together equal to two right angles :

then BD shall be in the same straight line with BC.

Proof. For if BD be not in the same straight line with BC, if possible, let BE be in the same straight line with BC.

Then because AB meets the straight line CBE, therefore the adjacent angles CBA, ABE are together equal to two right angles. I. 13.

But the angles CBA, ABD are also together equal to two right angles. Hyp.

Therefore the angles CBA, ABE are together equal to the angles CBA, ABD. Ax. 11.

From each of these equals take the common angle CBA; then the remaining angle ABE is equal to the remaining angle ABD; the part equal to the whole; which is impossible.

Therefore BE is not in the same straight line with BC.

And in the same way it may be shewn that no other line but BD can be in the same straight line with BC.

Therefore BD is in the same straight line with BC. Q. E. D.


EXERCISE.

ABCD is a rhombus; and the diagonal AC is bisected at O. If O is joined to the angular points B and D; shew that OB and OD are in one straight line.

Obs. (When two straight lines intersect at a point, four angles are formed; and any two of these angles which are not adjacent, are said to be **vertically opposite** to one another.)

PROPOSITION 15. THEOREM.

If two straight lines intersect one another, then the vertically opposite angles shall be equal.



Let the two straight lines AB, CD cut one another at the point E:

then shall the angle AEC be equal to the angle DEB,
and the angle CEB to the angle AED.

Proof. Because AE makes with CD the adjacent angles CEA, AED,

therefore these angles are together equal to two right angles. I. 13.

Again, because DE makes with AB the adjacent angles AED, DEB,

therefore these also are together equal to two right angles.
Therefore the angles CEA, AED are together equal to the angles AED, DEB.

From each of these equals, take the common angle AED;
then the remaining angle CEA is equal to the remaining angle DEB. Ax. 3.


In a similar way it may be shewn that the angle CEB is equal to the angle AED. Q. E. D.

COROLLARY 1. *From this it is manifest that, if two straight lines cut one another, the angles which they make at the point where they cut, are together equal to four right angles.*

COROLLARY 2. *Consequently, when any number of straight lines meet at a point, the sum of the angles made by consecutive lines is equal to four right angles.*

PROPOSITION 16. THEOREM.

If one side of a triangle be produced, then the exterior angle shall be greater than either of the interior opposite angles.



Let ABC be a triangle, and let one side BC be produced to D : then shall the exterior angle ACD be greater than either of the interior opposite angles CBA , BAC .

Construction. Bisect AC at E : I. 10.
Join BE ; and produce it to F , making EF equal to BE . I. 3.
Join FC .

Proof. Then in the triangles AEB , CEF ,
Because $\left\{ \begin{array}{l} AE \text{ is equal to } CE, \\ \text{and } EB \text{ to } EF; \\ \text{also the angle } AEB \text{ is equal to the vertically} \\ \text{opposite angle } CEF; \end{array} \right. \begin{array}{l} \text{Constr.} \\ \text{Constr.} \\ \text{I. 15.} \end{array}$
therefore the triangle AEB is equal to the triangle CEF in all respects: I. 4.

so that the angle BAE is equal to the angle ECF .

But the angle ECD is greater than its part, the angle ECF ;
therefore the angle ECD is greater than the angle BAE ;
that is, the angle ACD is greater than the angle BAC .

In a similar way, if BC be bisected, and the side AC produced to G , it may be shewn that the angle BCG is greater than the angle ABC .

But the angle BCG is equal to the angle ACD : I. 15.
therefore also the angle ACD is greater than the angle ABC .

Q. E. D.

PROPOSITION 17. THEOREM.

Any two angles of a triangle are together less than two right angles.



Let ABC be a triangle: then shall any two of its angles, as ABC , ACB , be together less than two right angles.

Construction. Produce the side BC to D .

Proof. Then because ACD is an exterior angle of the triangle ABC ,
therefore it is greater than the interior opposite angle ABC . I. 16.

To each of these add the angle ACB :
then the angles ACD , ACB are together greater than the angles ABC , ACB . Ax. 4.

But the adjacent angles ACD , ACB are together equal to two right angles. I. 13.

Therefore the angles ABC , ACB are together less than two right angles.

Similarly it may be shewn that the angles BAC , ACB , as also the angles CAB , ABC , are together less than two right angles. Q. E. D.

NOTE. It follows from this Proposition that every triangle must have at least two acute angles: for if one angle is obtuse, or a right angle, each of the other angles must be less than a right angle.

EXERCISES.


1. Enunciate this Proposition so as to shew that it is the converse of Axiom 12.

2. If any side of a triangle is produced both ways, the exterior angles so formed are together greater than two right angles.

3. Shew how a proof of Proposition 17 may be obtained by joining each vertex in turn to any point in the opposite side.

PROPOSITION 18. THEOREM.

If one side of a triangle be greater than another, then the angle opposite to the greater side shall be greater than the angle opposite to the less.



Let ABC be a triangle, in which the side AC is greater than the side AB :

then shall the angle ABC be greater than the angle ACB .

Construction. From AC , the greater, cut off a part AD equal to AB . I. 3.

Join BD .

Proof. Then in the triangle ABD ,
because AB is equal to AD ,

therefore the angle ABD is equal to the angle ADB . I. 5.

But the exterior angle ADB of the triangle BDC is greater than the interior opposite angle DCB , that is, greater than the angle ACB . I. 16.

Therefore also the angle ABD is greater than the angle ACB ; still more then is the angle ABC greater than the angle ACB . Q. E. D.

Euclid enunciated Proposition 18 as follows :


The greater side of every triangle has the greater angle opposite to it.

[This form of enunciation is found to be a common source of difficulty with beginners, who fail to distinguish what is *assumed* in it and what is to be *proved*.]

* [For Exercises see page 38.]

PROPOSITION 19. THEOREM.

If one angle of a triangle be greater than another, then the side opposite to the greater angle shall be greater than the side opposite to the less.



Let ABC be a triangle in which the angle ABC is greater than the angle ACB :

then shall the side AC be greater than the side AB .

Proof. For if AC be not greater than AB ,
it must be either equal to, or less than AB .

But AC is not equal to AB ,
for then the angle ABC would be equal to the angle ACB ; 1. 5.
but it is not. *Hyp.*

Neither is AC less than AB ;
for then the angle ABC would be less than the angle ACB ; 1. 18.
but it is not: *Hyp.*

Therefore AC is neither equal to, nor less than AB .

That is, AC is greater than AB . Q. E. D.

NOTE. The mode of demonstration used in this Proposition is known as the **Proof by Exhaustion**. It is applicable to cases in which one of certain mutually exclusive suppositions must necessarily be true; and it consists in shewing the falsity of each of those suppositions in turn *with one exception*: hence the truth of the remaining supposition is inferred.


Euclid enunciated Proposition 19 as follows :

The greater angle of every triangle is subtended by the greater side, or, has the greater side opposite to it.

[For Exercises see page 38.]

PROPOSITION 20. THEOREM.

Any two sides of a triangle are together greater than the third side.



Let ABC be a triangle:
then shall any two of its sides be together greater than the third side :

namely, BA , AC , shall be greater than CB ;

AC , CB greater than BA ;

and CB , BA greater than AC .

Construction. Produce BA to the point D , making AD equal to AC . I. 3.

Join DC .

Proof. Then in the triangle ADC ,
because AD is equal to AC , *Constr.*
therefore the angle ACD is equal to the angle ADC . I. 5.
But the angle BCD is greater than the angle ACD ; *Ax. 9.*
therefore also the angle BCD is greater than the angle ADC ,
that is, than the angle BDC .

And in the triangle BCD ,
because the angle BCD is greater than the angle BDC , *Pr.*
therefore the side BD is greater than the side CB . I. 19.


But BA and AC are together equal to BD ;
therefore BA and AC are together greater than CB .

Similarly it may be shewn
that AC , CB are together greater than BA ;
and CB , BA are together greater than AC . Q. E. D.

[For Exercises see page 38.]

PROPOSITION 21. THEOREM.

If from the ends of a side of a triangle, there be drawn two straight lines to a point within the triangle, then these straight lines shall be less than the other two sides of the triangle, but shall contain a greater angle.



Let ABC be a triangle, and from B, C , the ends of the side BC , let the two straight lines BD, CD be drawn to a point D within the triangle :

then (i) BD and DC shall be together less than BA and AC ;
 (ii) the angle BDC shall be greater than the angle BAC .

Construction. Produce BD to meet AC in E .

Proof. (i) In the triangle BAE , the two sides BA, AE are together greater than the third side BE : I. 20.

to each of these add EC ;

then BA, AC are together greater than BE, EC . *Ax. 2.*

Again, in the triangle DEC , the two sides DE, EC are together greater than DC : I. 20.

to each of these add BD ;

then BE, EC are together greater than BD, DC .

But it has been shewn that BA, AC are together greater than BE, EC :

still more then are BA, AC greater than BD, DC .

(ii) Again, the exterior angle BDC of the triangle DEC is, greater than the interior opposite angle DEC ; I. 16.

and the exterior angle DEC of the triangle BAE is greater than the interior opposite angle BAE , that is, than the angle BAC ; I. 16.

still more then is the angle BDC greater than the angle BAC .

Q.E.D.

EXERCISES

ON PROPOSITIONS 18 AND 19.

1. The hypotenuse is the greatest side of a right-angled triangle.
2. If two angles of a triangle are equal to one another, the sides also, which subtend the equal angles, are equal to one another. Prop. 6. Prove this indirectly by using the result of Prop. 18.
3. BC , the base of an isosceles triangle ABC , is produced to any point D ; shew that AD is greater than either of the equal sides.
4. If in a quadrilateral the greatest and least sides are opposite to one another, then each of the angles adjacent to the least side is greater than its opposite angle.
5. In a triangle ABC , if AC is not greater than AB , shew that any straight line drawn through the vertex A and terminated by the base BC , is less than AB .
6. ABC is a triangle, in which OB , OC bisect the angles ABC , ACB respectively: shew that, if AB is greater than AC , then OB is greater than OC .

ON PROPOSITION 20.


7. The difference of any two sides of a triangle is less than the third side.
8. In a quadrilateral, if two opposite sides which are not parallel are produced to meet one another; shew that the perimeter of the greater of the two triangles so formed is greater than the perimeter of the quadrilateral.
9. The sum of the distances of any point from the three angular points of a triangle is greater than half its perimeter.
10. The perimeter of a quadrilateral is greater than the sum of its diagonals.
11. Obtain a proof of Proposition 20 by bisecting an angle by a straight line which meets the opposite side.

ON PROPOSITION 21.

12. In Proposition 21 shew that the angle BDC is greater than the angle BAC by joining AD , and producing it towards the base.
13. The sum of the distances of any point within a triangle from its angular points is less than the perimeter of the triangle.

PROPOSITION 22. PROBLEM.

To describe a triangle having its sides equal to three given straight lines, any two of which are together greater than the third.



Let A, B, C be the three given straight lines, of which any two are together greater than the third.

It is required to describe a triangle of which the sides shall be equal to A, B, C .

Construction. Take a straight line DE terminated at the point D , but unlimited towards E .

Make DF equal to A , FG equal to B , and GH equal to C . 3.

From centre F , with radius FD , describe the circle DLK .

From centre G with radius GH , describe the circle MHK , cutting the former circle at K .

Join FK, GK .

Then shall the triangle KFG have its sides equal to the three straight lines A, B, C .

Proof. Because F is the centre of the circle DLK ,
 therefore FK is equal to FD : Def. 11.
 but FD is equal to A ; Constr.
 therefore also FK is equal to A . Ax. 1.

Again, because G is the centre of the circle MHK ,
 therefore GK is equal to GH : Def. 11.
 but GH is equal to C ; Constr.
 therefore also GK is equal to C . Ax. 1.

And FG is equal to B . Constr.


Therefore the triangle KFG has its sides KF, FG, GK equal respectively to the three given lines A, B, C . Q.E.F.

EXERCISE. (

On a given base describe a triangle, whose remaining sides shall be equal to two given straight lines. Point out how the construction fails, if any one of the three given lines is greater than the sum of the other two.

PROPOSITION 23. PROBLEM.

At a given point in a given straight line, to make an angle equal to a given angle.



Let AB be the given straight line, and A the given point in it; and let DCE be the given angle.

It is required to draw from A a straight line making with AB an angle equal to the given angle DCE .

Construction. In CD , CE take any points D and E ; and join DE .

From AB cut off AF equal to CD . I. 3.

On AF describe the triangle FAG , having the remaining sides AG , FG equal respectively to CE , ED . I. 22.

Then shall the angle FAG be equal to the angle DCE .

Proof. For in the triangles FAG , DCE ,


Because $\left\{ \begin{array}{l} \text{FA is equal to DC,} \\ \text{and AG is equal to CE;} \\ \text{and the base FG is equal to the base DE;} \end{array} \right. \begin{array}{l} \text{Constr.} \\ \text{Constr.} \\ \text{Constr.} \end{array}$

therefore the angle FAG is equal to the angle DCE . I. 8.

That is, AG makes with AB , at the given point A , an angle equal to the given angle DCE . Q.E.F.

PROPOSITION 24.

If two triangles have two sides of the one equal to two sides of the other, each to each, but the angle contained by the two sides of one greater than the angle contained by the corresponding sides of the other; then the base of that which has the greater angle shall be greater than the base of the other.



Let ABC , DEF be two triangles, in which the two sides BA , AC are equal to the two sides ED , DF , each to each, but the angle BAC greater than the angle EDF :

then shall the base BC be greater than the base EF .

* Of the two sides DE , DF , let DE be that which is not greater than DF .


Construction. At the point D , in the straight line ED , and on the same side of it as DF , make the angle EDG equal to the angle BAC . I. 23.

Make DG equal to DF or AC ; I. 3.
and join EG , GF .

Proof. Then in the triangles BAC , EDG ,
 Because $\left\{ \begin{array}{l} BA \text{ is equal to } ED, \\ \text{and } AC \text{ is equal to } DG, \\ \text{also the contained angle } BAC \text{ is equal to the} \\ \text{contained angle } EDG; \end{array} \right. \begin{array}{l} \text{Hyp.} \\ \text{Constr.} \\ \text{Constr.} \end{array}$
 Therefore the triangle BAC is equal to the triangle EDG in
 all respects: I. 4.

so that the base BC is equal to the base EG .

* See note on the next page.



Again, in the triangle FDG ,
because DG is equal to DF ,
therefore the angle DFG is equal to the angle DGF , I. 5.
but the angle DGF is greater than the angle EGF ;
therefore also the angle DFG is greater than the angle EGF ;
still more then is the angle EFG greater than the angle EGF .

And in the triangle EFG ,
because the angle EFG is greater than the angle EGF ,
therefore the side EG is greater than the side EF ; I. 19.
but EG was shewn to be equal to BC ;
therefore BC is greater than EF . Q.E.D.

* This condition was inserted by Simson to ensure that, in the complete construction, the point F should fall *below* EG . Without this condition it would be necessary to consider three cases: for F might fall *above*, or *upon*, or *below* EG ; and each figure would require separate proof.

We are however scarcely at liberty to employ Simson's condition without *proving* that it fulfils the object for which it was introduced.

This may be done as follows:

Let EG , DF , produced if necessary, intersect at K .

Then, since DE is not greater than DF ,
that is, since DE is not greater than DG ,
therefore the angle DGE is not greater than the angle DEG . I. 18.

But the exterior angle DKG is greater than the angle DEK : I. 16.
therefore the angle DKG is greater than the angle DGK .

Hence DG is greater than DK . I. 19.

But DG is equal to DF ;
therefore DF is greater than DK .
So that the point F must fall *below* EG .

Or the following method may be adopted.

PROPOSITION 24. [ALTERNATIVE PROOF.]


In the triangles ABC , DEF ,
let BA be equal to ED ,
and AC equal to DF ,
but let the angle BAC be greater than
the angle EDF :
then shall the base BC be greater than
the base EF .

For apply the triangle DEF to the
triangle ABC , so that D may fall on A ,
and DE along AB :

then because DE is equal to AB ,
therefore E must fall on B .

And because the angle EDF is less than the angle BAC ,
therefore DF must fall between AB and AC .

Let DF occupy the position AG .




Hyp.

CASE I. If G falls on BC :

Then G must be between B and C :
therefore BC is greater than BG .

But BG is equal to EF :
therefore BC is greater than EF .




CASE II. If G does not fall on BC .
Bisect the angle CAG by the straight line AK
which meets BC in K . I. 9.

Join GK .


Then in the triangles GAK , CAK ,
Because $\left\{ \begin{array}{l} GA \text{ is equal to } CA, \\ \text{and } AK \text{ is common to both;} \\ \text{and the angle } GAK \text{ is equal to the} \\ \text{angle } CAK; \end{array} \right. \quad \begin{array}{l} \text{Hyp.} \\ \text{Constr.} \end{array}$
therefore GK is equal to CK . I. 4.

But in the triangle BKG ,
the two sides BK , KG are together greater than the third side BG , I. 20.
that is, BK , CK are together greater than BG ;
therefore BC is greater than BG , or EF . Q.E.D.



PROPOSITION 25. THEOREM.

If two triangles have two sides of the one equal to two sides of the other, each to each, but the base of one greater than the base of the other; then the angle contained by the sides of that which has the greater base, shall be greater than the angle contained by the corresponding sides of the other.



Let ABC , DEF be two triangles which have the two sides BA , AC equal to the two sides ED , DF , each to each, but the base BC greater than the base EF :

then shall the angle BAC be greater than the angle EDF .

Proof. For if the angle BAC be not greater than the angle EDF , it must be either equal to, or less than the angle EDF .

But the angle BAC is not equal to the angle EDF ,
for then the base BC would be equal to the base EF ; i. 4.
but it is not. *Hyp.*

Neither is the angle BAC less than the angle EDF ,
for then the base BC would be less than the base EF ; i. 24.
but it is not. *Hyp.*

Therefore the angle BAC is neither equal to, nor less than the angle EDF ;
that is, the angle BAC is greater than the angle EDF . Q.E.D.


EXERCISE.

In a triangle ABC , the vertex A is joined to X , the middle point of the base BC ; shew that the angle AXB is obtuse or acute, according as AB is greater or less than AC .

PROPOSITION 26. THEOREM.

If two triangles have two angles of the one equal to two angles of the other, each to each, and a side of one equal to a side of the other, these sides being either adjacent to the equal angles, or opposite to equal angles in each; then shall the triangles be equal in all respects.

CASE I. When the equal sides are adjacent to the equal angles in the two triangles.



Let ABC , DEF be two triangles, which have the angles ABC , ACB , equal to the two angles DEF , DFE , each to each; and the side BC equal to the side EF : then shall the triangle ABC be equal to the triangle DEF in all respects;

that is, AB shall be equal to DE , and AC to DF , and the angle BAC shall be equal to the angle EDF .


For if AB be not equal to DE , one must be greater than the other. If possible, let AB be greater than DE .

Construction. From BA cut off BG equal to ED , I. 3. and join GC .

Proof. Then in the two triangles GBC , DEF ,
 Because $\begin{cases} GB \text{ is equal to } DE, & \text{Constr.} \\ \text{and } BC \text{ to } EF, & \text{Hyp.} \\ \text{also the contained angle } GBC \text{ is equal to the} & \\ \text{contained angle } DEF; & \text{Hyp.} \end{cases}$

therefore the triangles are equal in all respects; I. 4., so that the angle GCB is equal to the angle DFE .

But the angle ACB is equal to the angle DFE ; Hyp. therefore also the angle GCB is equal to the angle ACB ; Ax. 1. the part equal to the whole, which is impossible.



Therefore AB is not unequal to DE ,
that is, AB is equal to DE .

Hence in the triangles ABC , DEF ,

AB is equal to DE ,

Proved.

and BC is equal to EF ;

Hyp.

Because

also the contained angle ABC is equal to the
contained angle DEF ;


Hyp.

therefore the triangles are equal in all respects: 1. 4.

so that the side AC is equal to the side DF ;

and the angle EAC to the angle EDF . Q.E.D.

CASE II. When the equal sides are *opposite* to equal
angles in the two triangles.



Let ABC , DEF be two triangles which have the angles
 ABC , ACB equal to the angles DEF , DFE , each to each,
and the side AB equal to the side DE :

then shall the triangles ABC , DEF be equal in all respects;

that is, BC shall be equal to EF , and AC to DF ,

and the angle BAC shall be equal to the angle EDF .

For if BC be not equal to EF , one must be greater than the other. If possible, let BC be greater than EF .

Construction. From BC cut off BH equal to EF ,
and join AH . I. 3.

Proof. Then in the triangles ABH , DEF ,
Because $\left\{ \begin{array}{l} AB \text{ is equal to } DE, \\ \text{and } BH \text{ to } EF, \\ \text{also the contained angle } ABH \text{ is equal to the} \\ \text{contained angle } DEF; \end{array} \right. \begin{array}{l} \text{Hyp.} \\ \text{Constr.} \\ \text{Hyp.} \end{array}$
therefore the triangles are equal in all respects, I. 4.
so that the angle AHB is equal to the angle DFE .

But the angle DFE is equal to the angle ACB ; *Hyp.*
therefore the angle AHB is equal to the angle ACB ; *Ax. 1.*
that is, an exterior angle of the triangle ACH is equal to
an interior opposite angle; which is impossible. I. 16.

Therefore BC is not unequal to EF ,
that is, BC is equal to EF .

Hence in the triangles ABC , DEF ,
Because $\left\{ \begin{array}{l} AB \text{ is equal to } DE, \\ \text{and } BC \text{ is equal to } EF; \\ \text{also the contained angle } ABC \text{ is equal to the} \\ \text{contained angle } DEF; \end{array} \right. \begin{array}{l} \text{Hyp.} \\ \text{Proved.} \\ \text{Hyp.} \end{array}$
therefore the triangles are equal in all respects; I. 4.
so that the side AC is equal to the side DF ,
and the angle BAC to the angle EDF .

Q.E.D.

COROLLARY. In both cases of this Proposition it is seen
that the triangles may be made to coincide with one another;
and they are therefore equal in area.

ON THE IDENTICAL EQUALITY OF TRIANGLES

At the close of the first section of Book I., it is worth while to call special attention to those Propositions (viz. Props. 4, 8, 26) which deal with the *identical equality* of two triangles.

The results of these Propositions may be summarized thus:

Two triangles are equal to one another in all respects, when the following parts in each are equal, each to each.

1. Two sides, and the included angle. Prop. 4.
2. The three sides. Prop. 8, Cor.
3. (a) Two angles, and the adjacent side.
 (b) Two angles, and the side opposite one of them. } Prop. 26.

From this the beginner will perhaps surmise that two triangles may be shewn to be equal in all respects, when they have *three parts* equal, each to each; but to this statement two obvious exceptions must be made.

(i) When in two triangles the *three angles* of one are equal to the *three angles* of the other, each to each, it does *not* necessarily follow that the triangles are equal in all respects.

(ii) When in two triangles two sides of the one are equal to two sides of the other, each to each, and one angle equal to one angle, these not being the angles included by the equal sides; the triangles are *not* necessarily equal in all respects.

In these cases a further condition must be added to the hypothesis, before we can assert the identical equality of the two triangles.

[See Theorems and Exercises on Book I., Ex. 13, Page 92.]

We observe that in each of the three cases already proved of identical equality in two triangles, namely in Propositions 4, 8, 26, it is shewn that the triangles may be made to *coincide with one another*; so that they are equal in *area*, as in all other respects. Euclid however restricted himself to the use of Prop. 4, when he required to deduce the equality in *area* of two triangles from the equality of certain of their parts.

This restriction has been abandoned in the present text-book.
 [See note to Prop. 34.]

EXERCISES ON PROPOSITIONS 12—26.

1. If BX and CY , the bisectors of the angles at the base BC of an isosceles triangle ABC , meet the opposite sides in X and Y ; shew that the triangles YBC , XCB are equal in all respects.

2. Shew that the perpendiculars drawn from the extremities of the base of an isosceles triangle to the opposite sides are equal.

3. Any point on the bisector of an angle is equidistant from the arms of the angle.

4. Through C , the middle point of a straight line AB , any straight line is drawn, and perpendiculars AX and BY are dropped upon it from A and B : shew that AX is equal to BY .

5. If the bisector of the vertical angle of a triangle is at right angles to the base, the triangle is isosceles.

6. The perpendicular is the shortest straight line that can be drawn from a given point to a given straight line; and of others, that which is nearer to the perpendicular is less than the more remote; and two, and only two equal straight lines can be drawn from the given point to the given straight line, one on each side of the perpendicular.

7. From two given points on the same side of a given straight line, draw two straight lines, which shall meet in the given straight line and make equal angles with it.

Let AB be the given straight line, and P, Q the given points.

It is required to draw from P and Q to a point in AB , two straight lines that shall be equally inclined to AB .

Construction. From P draw PH perpendicular to AB : produce PH to P' , making HP' equal to PH . Draw QP' , meeting AB in K . Join PK .



Then PK, QK shall be the required lines. [Supply the proof.]

8. In a given straight line find a point which is equidistant from two given intersecting straight lines. In what case is this impossible?

9. Through a given point draw a straight line such that the perpendiculars drawn to it from two given points may be equal.

In what case is this impossible?


SECTION II.

PARALLEL STRAIGHT LINES AND PARALLELOGRAMS.

DEFINITION. Parallel straight lines are such as, being in the same plane, do not meet however far they are produced in both directions.

When two straight lines **AB**, **CD** are met by a third straight line **EF**, *eight* angles are formed, to which for the sake of distinction particular names are given.

Thus in the adjoining figure,
 1, 2, 7, 8 are called **exterior** angles,
 3, 4, 5, 6 are called **interior** angles,
 4 and 6 are said to be **alternate** angles ;
 'so also the angles 3 and 5 are alternate to one another.



Of the angles 2 and 6, 2 is referred to as the **exterior** angle, and 6 as the **interior opposite** angle on the same side of **EF**.

2 and 6 are sometimes called **corresponding** angles.

So also, 1 and 5, 7 and 3, 8 and 4 are corresponding angles.

Euclid's treatment of parallel straight lines is based upon his twelfth Axiom, which we here repeat.


AXIOM 12. If a straight line cut two straight lines so as to make the two interior angles on the same side of it together less than two right angles, these straight lines, being continually produced, will at length meet on that side on which are the angles which are together less than two right angles.

Thus in the figure given above, if the two angles 3 and 6 are together less than two right angles, it is asserted that **AB** and **CD** will meet towards **B** and **D**.

This Axiom is used to establish I. 29 : some remarks upon it will be found in a note on that Proposition.

PROPOSITION 27. THEOREM.

If a straight line, falling on two other straight lines, make the alternate angles equal to one another, then the straight lines shall be parallel.



Let the straight line EF cut the two straight lines AB , CD at G and H , so as to make the alternate angles AGH , GHD equal to one another:

then shall AB and CD be parallel.

Proof. For if AB and CD be not parallel, they will meet, if produced, either towards B and D , or towards A and C .

If possible, let AB and CD , when produced, meet towards B and D , at the point K .

Then KGH is a triangle, of which one side KG is produced to A :

therefore the exterior angle AGH is greater than the interior opposite angle GKH . I. 16.

But the angle AGH is equal to the angle GKH : *Hyp.*
hence the angles AGH and GKH are both equal and unequal;
which is impossible.

Therefore AB and CD cannot meet when produced towards B and D .

Similarly it may be shewn that they cannot meet towards A and C :


therefore they are parallel.

Q. E. D.

PROPOSITION 28. THEOREM.

If a straight line, falling on two other straight lines, make an exterior angle equal to the interior opposite angle on the same side of the line; or if it make the interior angles on the same side together equal to two right angles, then the two straight lines shall be parallel.

E.



Let the straight line EF cut the two straight lines AB, CD in G and H: and

First, let the exterior angle EGB be equal to the interior opposite angle GHD:

then shall AB and CD be parallel.

Proof. Because the angle EGB is equal to the angle GHD; and, because the angle EGB is also equal to the vertically opposite angle AGH; I. 15.

therefore the angle AGH is equal to the angle GHD;

but these are alternate angles;

therefore AB and CD are parallel. I.

Q. E. D.

Secondly, let the two interior angles BGH, GHD be together equal to two right angles:

then shall AB and CD be parallel.

Proof. Because the angles BGH, GHD are together equal to two right angles; *Hyp.*

and because the adjacent angles BGH, AGH are also together equal to two right angles; I. 13.

therefore the angles BGH, AGH are together equal to the two angles BGH, GHD.


From these equals take the common angle BGH: then the remaining angle AGH is equal to the remaining angle GHD: and these are alternate angles;

therefore AB and CD are parallel. I. 27.

Q. E. D.

PROPOSITION 29. THEOREM.

If a straight line fall on two parallel straight lines, then it shall make the alternate angles equal to one another, and the exterior angle equal to the interior opposite angle on the same side; and also the two interior angles on the same side equal to two right angles.



Let the straight line EF fall on the parallel straight lines AB, CD:

- then (i) the alternate angles AGH, GHD shall be equal to one another;
- (ii) the exterior angle EGB shall be equal to the interior opposite angle GHD;
- (iii) the two interior angles BGH, GHD shall be together equal to two right angles.

Proof. (i) For if the angle AGH be not equal to the angle GHD, one of them must be greater than the other.

If possible, let the angle AGH be greater than the angle GHD;

add to each the angle BGH:

then the angles AGH, BGH are together greater than the angles BGH, GHD.

But the adjacent angles AGH, BGH are together equal to two right angles;


therefore the angles BGH, GHD are together less than two right angles;

therefore AB and CD meet towards B and D. *Ax. 12.*

But they never meet, since they are parallel. *Hyp.*

Therefore the angle AGH is not unequal to the angle GHD: that is, the alternate angles AGH, GHD are equal.

(Over)



(ii) Again, because the angle AGH is equal to the vertically opposite angle EGB; I. 15.

and because the angle AGH is equal to the angle GHD;

Proved.

therefore the exterior angle EGB is equal to the interior opposite angle GHD.

(iii) Lastly, the angle EGB is equal to the angle GHD;

Proved.

add to each the angle BGH;

then the angles EGB, BGH are together equal to the angles BGH, GHD.

But the adjacent angles EGB, BGH are together equal to two right angles: I. 13.

therefore also the two interior angles BGH, GHD are together equal to two right angles. Q.E.D.

EXERCISES ON PROPOSITIONS 27, 28, 29.

1. Two straight lines AB, CD bisect one another at O: shew that the straight lines joining AC and BD are parallel. [I. 27.]

2. Straight lines which are perpendicular to the same straight line are parallel to one another. [I. 27 or I. 28.]

3. If a straight line meet two or more parallel straight lines, and is perpendicular to one of them, it is also perpendicular to all the others. [I. 29.]

4. If two straight lines are parallel to two other straight lines, each to each, then the angles contained by the first pair are equal respectively to the angles contained by the second pair. [I. 29.]

NOTE ON THE TWELFTH AXIOM.

It must be admitted that Euclid's twelfth Axiom is unsatisfactory as the basis of a theory of parallel straight lines. It cannot be regarded as either simple or self-evident, and it therefore falls short of the essential characteristics of an axiom: nor is the difficulty entirely removed by considering it as a corollary to Proposition 17, of which it is the converse.

Many substitutes have been proposed; but we need only notice here the system which has met with most general approval.

This system rests on the following hypothesis, which is put forward as a fundamental Axiom.

AXIOM. *Two intersecting straight lines cannot be both parallel to a third straight line.*

This statement is known as **Playfair's Axiom**; and though it is not altogether free from objection, it is recommended as both simpler and more fundamental than that employed by Euclid, and more readily admitted without proof.

Propositions 27 and 28 having been proved in the usual way, the first part of Proposition 29 is then given thus.

PROPOSITION 29. [ALTERNATIVE PROOF.]

If a straight line fall on two parallel straight lines, then it shall make the alternate angles equal.

Let the straight line EF meet the two parallel straight lines AB, CD, at G and H:

then shall the alternate angles AGH, GHD be equal.

For if the angle AGH is not equal to the angle GHD:

at G in the straight line HG make the angle HGP equal to the angle GHD, and alternate to it. I. 23.

Then PG and CD are parallel. I. 27.


But AB and CD are parallel: *Hyp.*
therefore the two intersecting straight lines AG, PG are both parallel to CD:

which is impossible.

Playfair's Axiom.

Therefore the angle AGH is not unequal to the angle GHD, that is, the alternate angles AGH, GHD are equal. Q.E.D.

The second and third parts of the Proposition may then be deduced as in the text; and Euclid's Axiom 12 follows as a Corollary.



It may happen that the data of the problem are so related to one another that the resulting loci do not intersect; in this case the problem is impossible.

For example, if in Ex. 1, page 118, the length of the given median is *less than* the given altitude, the straight line CD will not be intersected by the circle, and no triangle can fulfil the conditions of the problem. If the length of the median is *equal* to the given altitude, *one* point is common to the two loci; and consequently only one solution of the problem exists; and we have seen that there are two solutions, if the median is greater than the altitude.


In examples of this kind the student should make a point of investigating the relations which must exist among the data, in order that the problem may be possible; and he must observe that if under certain relations *two* solutions are possible, and under other relations no solution exists, there will always be some *intermediate* relation under which *one and only one* solution is possible.

EXAMPLES.

1. Find a point in a given straight line which is equidistant from two given points.
2. Find a point which is at given distances from each of two given straight lines. How many solutions are possible?
3. On a given base construct a triangle, having given one angle at the base and the length of the opposite side. Examine the relations which must exist among the data in order that there may be two solutions, one solution, or that the problem may be impossible.
4. On the base of a given triangle construct a second triangle equal in area to the first, and having its vertex in a given straight line.
5. Construct an isosceles triangle equal in area to a given triangle, and standing on the same base.
6. Find a point which is at a given distance from a given point, and is equidistant from two given parallel straight lines.

PROPOSITION 31. PROBLEM.

To draw a straight line through a given point parallel to a given straight line.



Let A be the given point, and BC the given straight line. It is required to draw through A a straight line parallel to BC.

Construction. In BC take any point D; and join AD. At the point A in DA, make the angle DAE equal to the angle ADC, and alternate to it I. 23.

and produce EA to F.

Then shall EF be parallel to BC.

Proof. Because the straight line AD, meeting the two straight lines EF, BC, makes the alternate angles EAD, ADC equal; Constr.

therefore EF is parallel to BC; I. 27.

and it has been drawn through the given point A.

Q. E. F.

EXERCISES.

1. Any straight line drawn parallel to the base of an isosceles triangle makes equal angles with the sides.

2. If from any point in the bisector of an angle a straight line is drawn parallel to either arm of the angle, the triangle thus formed is isosceles.


3. From a given point draw a straight line that shall make with a given straight line an angle equal to a given angle.

4. From X, a point in the base BC of an isosceles triangle ABC, a straight line is drawn at right angles to the base, cutting AB in Y, and CA produced in Z: shew the triangle AYZ is isosceles.

5. If the straight line which bisects an exterior angle of a triangle is parallel to the opposite side, shew that the triangle is isosceles.

PROPOSITION 32. THEOREM.

If a side of a triangle be produced, then the exterior angle shall be equal to the sum of the two interior opposite angles: also the three interior angles of a triangle are together equal to two right angles.



Let ABC be a triangle, and let one of its sides BC be produced to D ;

then (i) the exterior angle ACD shall be equal to the sum of the two interior opposite angles CAB , ABC ;

(ii) the three interior angles ABC , BCA , CAB shall be together equal to two right angles.

Construction. Through C draw CE parallel to BA . I. 31.

Proof. (i) Then because BA and CE are parallel, and AC meets them,

therefore the angle ACE is equal to the alternate angle CAB . I. 29.

Again, because BA and CE are parallel, and BD meets them, therefore the exterior angle ECD is equal to the interior opposite angle ABC . I. 29.

Therefore the whole exterior angle ACD is equal to the sum of the two interior opposite angles CAB , ABC .

(ii) Again, since the angle ACD is equal to the sum of the angles CAB , ABC ; *Proved.*

to each of these equals add the angle BCA :

then the angles BCA , ACD are together equal to the three angles BCA , CAB , ABC .

But the adjacent angles BCA , ACD are together equal to two right angles; I. 13.

therefore also the angles BCA , CAB , ABC are together equal to two right angles. Q. E. D.

From this Proposition we draw the following important inferences.

1. *If two triangles have two angles of the one equal to two angles of the other, each to each, then the third angle of the one is equal to the third angle of the other.*

2. *In any right-angled triangle the two acute angles are complementary.*

3. *In a right-angled isosceles triangle each of the equal angles is half a right angle.*

4. *If one angle of a triangle is equal to the sum of the other two, the triangle is right-angled.*

5. *The sum of the angles of any quadrilateral figure is equal to four right angles.*

6. *Each angle of an equilateral triangle is two-thirds of a right angle.*

EXERCISES ON PROPOSITION 32

1. Prove that the three angles of a triangle are together equal to two right angles,

(i) by drawing through the vertex a straight line parallel to the base;

(ii) by joining the vertex to any point in the base.

2. If the base of any triangle is produced both ways, shew that the sum of the two exterior angles diminished by the vertical angle is equal to two right angles.

3. If two straight lines are perpendicular to two other straight lines, each to each, the acute angle between the first pair is equal to the acute angle between the second pair.

4. Every right-angled triangle is divided into two isosceles triangles by a straight line drawn from the right angle to the middle point of the hypotenuse.

Hence the joining line is equal to half the hypotenuse.

5. Draw a straight line at right angles to a given finite straight line from one of its extremities, without producing the given straight line.

[Let AB be the given straight line. On AB describe any isosceles triangle ACB. Produce BC to D, making CD equal to BC. Join AD. Then shall AD be perpendicular to AB.]


6. *Trisect a right angle.*

7. The angle contained by the bisectors of the angles at the base of an isosceles triangle is equal to an exterior angle formed by producing the base.

8. The angle contained by the bisectors of two adjacent angles of a quadrilateral is equal to half the sum of the remaining angles.

The following theorems were added as corollaries to Proposition 32 by Robert Simson.

COROLLARY 1. *All the interior angles of any rectilineal figure, with four right angles, are together equal to twice as many right angles as the figure has sides.*



Let ABCDE be any rectilineal figure.

Take F, any point within it,

and join F to each of the angular points of the figure.

Then the figure is divided into as many triangles as it has sides.

And the three angles of each triangle are together equal to two right angles. I. 32.


Hence all the angles of all the triangles are together equal to twice as many right angles as the figure has sides.

But all the angles of all the triangles make up the interior angles of the figure, together with the angles at F;

and the angles at F are together equal to four right angles: I. 15, Cor.

Therefore all the interior angles of the figure, with four right angles, are together equal to twice as many right angles as the figure has sides. Q. E. D.

COROLLARY 2. *If the sides of a rectilineal figure, which has no re-entrant angle, are produced in order, then all the exterior angles so formed are together equal to four right angles*



For at each angular point of the figure, the interior angle and the exterior angle are together equal to two right angles. • •

I. 13.

Therefore all the interior angles, with all the exterior angles, are together equal to twice as many right angles as the figure has sides.

But all the interior angles, with four right angles, are together equal to twice as many right angles as the figure has sides.

I. 32, Cor. 1.

Therefore all the interior angles, with all the exterior angles, are together equal to all the interior angles, with four right angles.

Therefore the exterior angles are together equal to four right angles.

∴ Q. E. D.

EXERCISES ON SIMSON'S COROLLARIES.

[A polygon is said to be **regular** when it has all its sides and all its angles equal.]

1. Express in terms of a right angle the magnitude of each angle of (i) a regular hexagon, (ii) a regular octagon.

2. If one side of a regular hexagon is produced, shew that the exterior angle is equal to the angle of an equilateral triangle.

3. Prove Simson's first Corollary by joining one vertex of the rectilineal figure to each of the other vertices.

4. Find the magnitude of each angle of a regular polygon of n sides.

5. If the alternate sides of any polygon be produced to meet, the sum of the included angles, together with eight right angles, will be equal to twice as many right angles as the figure has sides.

PROPOSITION 33. THEOREM.

The straight lines which join the extremities of two equal and parallel straight lines towards the same parts are themselves equal and parallel.



Let AB and CD be equal and parallel straight lines; and let them be joined towards the same parts by the straight lines AC and BD:

then shall AC and BD be equal and parallel.

Construction. Join BC.

Proof. Then because AB and CD are parallel, and BC meets them,

therefore the alternate angles ABC, BCD are equal. I. 29.

Now in the triangles ABC, DCB,

Because $\left\{ \begin{array}{l} \text{AB is equal to DC,} \\ \text{and BC is common to both;} \\ \text{also the angle ABC is equal to the angle} \\ \text{DCB;} \end{array} \right. \begin{array}{l} \text{Hyp.} \\ \\ \text{Proved} \end{array}$

therefore the triangles are equal in all respects; I. 4.

so that the base AC is equal to the base DB,

and the angle ACB equal to the angle DBC;

but these are alternate angles;

therefore AC and BD are parallel: I. 27


and it has been shewn that they are also equal.

Q. E. D.

DEFINITION. A **Parallelogram** is a four-sided figure whose opposite sides are parallel.

PROPOSITION 34. THEOREM.

The opposite sides and angles of a parallelogram are equal to one another, and each diagonal bisects the parallelogram.



Let $ACDB$ be a parallelogram, of which BC is a diagonal: then shall the opposite sides and angles of the figure be equal to one another; and the diagonal BC shall bisect it.

Proof. Because AB and CD are parallel, and BC meets them,

therefore the alternate angles ABC , DCB are equal. 1. 29.

Again, because AC and BD are parallel, and BC meets them,

therefore the alternate angles ACB , DBC are equal. 1. 29.

Hence in the triangles ABC , DCB ,

Because $\begin{cases} \text{the angle } ABC \text{ is equal to the angle } DCB, \\ \text{and the angle } ACB \text{ is equal to the angle } DBC; \\ \text{also the side } BC, \text{ which is adjacent to the equal} \\ \text{angles, is common to both,} \end{cases}$

therefore the two triangles ABC , DCB are equal in all respects; 1. 26.

so that AB is equal to DC , and AC to DB ;

and the angle BAC is equal to the angle CDB .

Also, because the angle ABC is equal to the angle DCB ,

and the angle CBD equal to the angle BCA ,

therefore the whole angle ABD is equal to the whole angle DCA .

And since it has been shewn that the triangles ABC , DCB are equal in all respects,

therefore the diagonal BC bisects the parallelogram $ACDB$.

Q. E. D.

[See note on next page.]

NOTE. To the proof which is here given Euclid added an application of Proposition 4, with a view to shewing that the triangles **ABC**, **DCB** are equal in area, and that therefore the diagonal **BC** bisects the parallelogram. This equality of area is however sufficiently established by the step which depends upon I. 26. [See page 48.]

EXERCISES.

1. *If one angle of a parallelogram is a right angle, all its angles are right angles.*
2. *If the opposite sides of a quadrilateral are equal, the figure is a parallelogram.*
3. *If the opposite angles of a quadrilateral are equal, the figure is a parallelogram.*
4. *If a quadrilateral has all its sides equal and one angle a right angle, all its angles are right angles.*
5. *The diagonals of a parallelogram bisect each other.*
6. *If the diagonals of a quadrilateral bisect each other, the figure is a parallelogram.*
7. *If two opposite angles of a parallelogram are bisected by the diagonal which joins them, the figure is equilateral.*
8. *If the diagonals of a parallelogram are equal, all its angles are right angles.*
9. *In a parallelogram which is not rectangular the diagonals are unequal.*
10. *Any straight line drawn through the middle point of a diagonal of a parallelogram and terminated by a pair of opposite sides, is bisected at that point.*
11. *If two parallelograms have two adjacent sides of one equal to two adjacent sides of the other, each to each, and one angle of one equal to one angle of the other, the parallelograms are equal in all respects.*
12. *Two rectangles are equal if two adjacent sides of one are equal to two adjacent sides of the other, each to each.*
13. *In a parallelogram the perpendiculars drawn from one pair of opposite angles to the diagonal which joins the other pair are equal.*
14. *If **ABCD** is a parallelogram, and **X**, **Y** respectively the middle points of the sides **AD**, **BC**; shew that the figure **AYCX** is a parallelogram.*

MISCELLANEOUS EXERCISES ON SECTIONS I. AND II.

1. Shew that the construction in Proposition 2 may generally be performed in eight different ways. Point out the exceptional case.
2. The bisectors of two vertically opposite angles are in the same straight line.
3. In the figure of Proposition 16, if AF is joined, shew
 - (i) that AF is equal to BC;
 - (ii) that the triangle ABC is equal to the triangle CFA in all respects.
4. ABC is a triangle right-angled at B, and BC is produced to D: shew that the angle ACD is obtuse.
- X 5. Shew that in any regular polygon of n sides each angle contains $\frac{2(n-2)}{n}$ right angles.
6. The angle contained by the bisectors of the angles at the base of any triangle is equal to the vertical angle together with half the sum of the base angles.
7. The angle contained by the bisectors of two exterior angles of any triangle is equal to half the sum of the two corresponding interior angles.
8. If perpendiculars are drawn to two intersecting straight lines from any point between them, shew that the bisector of the angle between the perpendiculars is parallel to (or coincident with) the bisector of the angle between the given straight lines.
9. If two points P, Q be taken in the equal sides of an isosceles triangle ABC, so that BP is equal to CQ, shew that PQ is parallel to BC.
10. ABC and DEF are two triangles, such that AB, BC are equal and parallel to DE, EF, each to each; shew that AC is equal and parallel to DF.
11. Prove the second Corollary to Prop. 32 by drawing through any angular point lines parallel to all the sides.
12. If two sides of a quadrilateral are parallel, and the remaining two sides equal but not parallel, shew that the opposite angles are supplementary; also that the diagonals are equal.

SECTION III.

THE AREAS OF PARALLELOGRAMS AND TRIANGLES.

Hitherto when two figures have been said to be *equal*, it has been implied that they are *identically* equal, that is, equal in all respects.

In Section III. of Euclid's first Book, we have to consider the equality in *area* of parallelograms and triangles which are not necessarily equal in all respects.

[The ultimate test of equality, as we have already seen, is afforded by Axiom 8, which asserts that magnitudes which *may be made to coincide with one another* are equal. Now figures which are not identically equal, cannot be made to coincide without first undergoing some change of form: hence the method of direct *superposition* is unsuited to the purposes of the present section.


We shall see however from Euclid's proof of Proposition 35, that two figures which are not identically equal, may nevertheless be so related to a third figure, that it is possible to infer the equality of their areas.]

DEFINITIONS.

1. (The **Altitude** of a parallelogram with reference to a given side as base, is the perpendicular distance between the base and the opposite side.)
2. (The **Altitude** of a triangle with reference to a given side as base, is the perpendicular distance of the opposite vertex from the base.)

PROPOSITION 35. THEOREM.

Parallelograms on the same base, and between the same parallels, are equal in area.



Let the parallelograms $ABCD$, $EBCF$ be on the same base BC , and between the same parallels BC , AF :

then shall the parallelogram $ABCD$ be equal in area to the parallelogram $EBCF$.

CASE I. If the sides of the given parallelograms, opposite to the common base BC , are terminated at the same point D :

then because each of the parallelograms is double of the triangle BDC ;

I. 34.

therefore they are equal to one another. Ax. 6.

CASE II. But if the sides AD , EF , opposite to the base BC , are not terminated at the same point :

then because $ABCD$ is a parallelogram,

therefore AD is equal to the opposite side BC ; I. 34.

and for a similar reason, EF is equal to BC ;

therefore AD is equal to EF . Ax. 1.

Hence the whole, or remainder, EA is equal to the whole, or remainder, FD .

Then in the triangles FDC , EAB ,

Because $\left\{ \begin{array}{l} FD \text{ is equal to } EA, \\ \text{and } DC \text{ is equal to the opposite side } AB, \\ \text{also the exterior angle } FDC \text{ is equal to the interior} \\ \text{opposite angle } EAB, \end{array} \right. \quad \begin{array}{l} \text{Proved.} \\ \text{I. 34.} \\ \text{I. 29.} \end{array}$

therefore the triangle FDC is equal to the triangle EAB . I. 4.


From the whole figure $ABCF$ take the triangle FDC ;
and from the same figure take the equal triangle EAB ;

then the remainders are equal ; Ax. 3.

that is, the parallelogram $ABCD$ is equal to the parallelogram $EBCF$. Q. E. D.

PROPOSITION 36. THEOREM.

Parallelograms on equal bases, and between the same parallels, are equal in area.



Let $ABCD$, $EFGH$ be parallelograms on equal bases BC , FG , and between the same parallels AH , BG : then shall the parallelogram $ABCD$ be equal to the parallelogram $EFGH$.

Construction. Join BE , CH .

Proof. Then because BC is equal to FG ; *Hyp.*
 and FG is equal to the opposite side EH ; 1. 34.
 therefore BC is equal to EH : *Ax.* 1.
 and they are also parallel; *Hyp.*
 therefore BE and CH , which join them towards the same parts, are also equal and parallel. 1. 33.

Therefore $EBCH$ is a parallelogram. *Def.* 26.

Now the parallelogram $ABCD$ is equal to $EBCH$; 1. 35.
 for they are on the same base BC , and between the same parallels BC , AH .

Also the parallelogram $EFGH$ is equal to $EBCH$; 1. 35.
 for they are on the same base EH , and between the same parallels AH , BG .

Therefore the parallelogram $ABCD$ is equal to the parallelogram $EFGH$. *Ax.* 1.

Q. E. D.


From the last two Propositions we infer that :

- (i) *A parallelogram is equal in area to a rectangle of equal base and equal altitude.*
- (ii) *Parallelograms on equal bases and of equal altitudes are equal in area.*

- (iii) *Of two parallelograms of equal altitudes, that is the greater which has the greater base; and of two parallelograms on equal bases, that is the greater which has the greater altitude.*

PROPOSITION 37. THEOREM.

Triangles on the same base, and between the same parallels, are equal in area.



Let the triangles ABC , DBC be upon the same base BC , and between the same parallels BC , AD .

Then shall the triangle ABC be equal to the triangle DBC .

Construction. Through B draw BE parallel to CA , to meet DA produced in E ; I. 31.

through C draw CF parallel to BD , to meet AD produced in F .

Proof. Then, by construction, each of the figures $EBCA$, $DBC F$ is a parallelogram. Def. 26.

And $EBCA$ is equal to $DBC F$;
 for they are on the same base BC , and between the same parallels BC , EF . I. 35.

And the triangle ABC is half of the parallelogram $EBCA$,
 for the diagonal AB bisects it. I. 34.

Also the triangle DBC is half of the parallelogram $DBC F$,
 for the diagonal DC bisects it. I. 34.


But the halves of equal things are equal; *Ax. 7*;
 therefore the triangle ABC is equal to the triangle DBC .

Q.E.D.

[For Exercises see page 73.]

PROPOSITION 38. THEOREM.

Triangles on equal bases, and between the same parallels, are equal in area.



Let the triangles ABC , DEF be on equal bases BC , EF , and between the same parallels BF , AD :

then shall the triangle ABC be equal to the triangle DEF .

Construction. Through B draw BG parallel to CA , to meet DA produced in G ; I. 31.
through F draw FH parallel to ED , to meet AD produced in H .

Proof. Then, by construction, each of the figures $GBCA$, $DEFH$ is a parallelogram. Def. 26.

And $GBCA$ is equal to $DEFH$;
for they are on equal bases BC , EF , and between the same parallels BF , GH . I. 36.

And the triangle ABC is half of the parallelogram $GBCA$,
for the diagonal AB bisects it. I. 34.

Also the triangle DEF is half the parallelogram $DEFH$,
for the diagonal DF bisects it. I. 34.

But the halves of equal things are equal: *Ax. 7.*
therefore the triangle ABC is equal to the triangle DEF .

Q.E.D.

From this Proposition we infer that :


(i) *Triangles on equal bases and of equal altitude are equal in area.*

(ii) *Of two triangles of the same altitude, that is the greater which has the greater base : and of two triangles on the same base, or on equal bases, that is the greater which has the greater altitude.*

[For Exercises see page 73.]

PROPOSITION 39. THEOREM.

Equal triangles on the same base, and on the same side of it, are between the same parallels.



Let the triangles ABC , DBC which stand on the same base BC , and on the same side of it, be equal in area :
 then shall they be between the same parallels ;
 that is, if AD be joined, AD shall be parallel to BC .

Construction. For if AD be not parallel to BC ,
 if possible, through A draw AE parallel to BC , I. 31.
 meeting BD , or BD produced, in E .
 Join EC .

Proof. Now the triangle ABC is equal to the triangle EBC ,
 for they are on the same base BC , and between the same
 parallels BC , AE . I. 37.

But the triangle ABC is equal to the triangle DBC ; *Hyp.*
 therefore also the triangle DBC is equal to the triangle EBC ;
 the whole equal to the part ; which is impossible.

Therefore AE is not parallel to BC .

Similarly it can be shewn that no other straight line
 through A , except AD , is parallel to BC .

Therefore AD is parallel to BC .

Q.E.D.


From this Proposition it follows that :

Equal triangles on the same base have equal altitudes.

[For Exercises see page 73.]

PROPOSITION 40. THEOREM.

Equal triangles, on equal bases in the same straight line, and on the same side of it, are between the same parallels.



Let the triangles ABC , DEF which stand on equal bases BC , EF , in the same straight line BF , and on the same side of it, be equal in area :

then shall they be between the same parallels :

that is, if AD be joined, AD shall be parallel to BF .

Construction. For if AD be not parallel to BF ,
if possible, through A draw AG parallel to BF , i. 31.
meeting ED , or ED produced, in G .

Join GF .

Proof. Now the triangle ABC is equal to the triangle GEF ,
for they are on equal bases BC , EF , and between the
same parallels BF , AG . i. 38.

But the triangle ABC is equal to the triangle DEF : *Hyp.*
therefore also the triangle DEF is equal to the triangle GEF :
the whole equal to the part; which is impossible.

Therefore AG is not parallel to BF .

Similarly it can be shewn that no other straight line
through A , except AD , is parallel to BF .

Therefore AD is parallel to BF .

Q.E.D.

From this Proposition it follows that :

- (i) *Equal triangles on equal bases have equal altitudes.*
- (ii) *Equal triangles of equal altitudes have equal bases.*

EXERCISES ON PROPOSITIONS 37—40. •

DEFINITION. Each of the three straight lines which join the angular points of a triangle to the middle points of the opposite sides is called a **Median** of the triangle.

ON PROP. 37.

1. If, in the figure of Prop. 37, AC and BD intersect in K, shew that
 - (i) the triangles AKB, DKC are equal in area.
 - (ii) the quadrilaterals EBKA, FCKD are equal.
2. In the figure of 1. 16, shew that the triangles ABC, FBC are equal in area.
3. On the base of a given triangle construct a second triangle, equal in area to the first, and having its vertex in a given straight line.
4. Describe an isosceles triangle equal in area to a given triangle and standing on the same base.

ON PROP. 38.

5. A triangle is divided by each of its medians into two parts of equal area.
6. A parallelogram is divided by its diagonals into four triangles of equal area.
7. ABC is a triangle, and its base BC is bisected at X; if Y be any point in the median AX, shew that the triangles ABY, ACY are equal in area.
8. In AC, a diagonal of the parallelogram ABCD, any point X is taken, and XB, XD are drawn; shew that the triangle BAX is equal to the triangle DAX.
9. If two triangles have two sides of one respectively equal to two sides of the other, and the angles contained by those sides supplementary, the triangles are equal in area.

ON PROP. 39.


10. The straight line which joins the middle points of two sides of a triangle is parallel to the third side.
11. If two straight lines AB, CD intersect in O, so that the triangle AOC is equal to the triangle DOB, shew that AD and CB are parallel.

ON PROP. 40.

12. Deduce Prop. 40 from Prop. 39 by joining AE, AF in the figure of page 72.

PROPOSITION 41. THEOREM.

If a parallelogram and a triangle be on the same base and between the same parallels, the parallelogram shall be double of the triangle.



Let the parallelogram $ABCD$, and the triangle EBC be upon the same base BC , and between the same parallels BC , AE :

then shall the parallelogram $ABCD$ be double of the triangle EBC .

Construction. Join AC .

Proof. Then the triangle ABC is equal to the triangle EBC , for they are on the same base BC , and between the same parallels BC , AE . I. 37.

But the parallelogram $ABCD$ is double of the triangle ABC , for the diagonal AC bisects the parallelogram. I. 34.

Therefore the parallelogram $ABCD$ is also double of the triangle EBC . Q.E.D.

EXERCISES.

1. $ABCD$ is a parallelogram, and X , Y are the middle points of the sides AD , BC ; if Z is any point in XY , or XY produced, shew that the triangle AZB is one quarter of the parallelogram $ABCD$.


2. Describe a right-angled isosceles triangle equal to a given square.

3. If $ABCD$ is a parallelogram, and X , Y any points in DC and AD respectively: shew that the triangles AXB , BYC are equal in area.

4. $ABCD$ is a parallelogram, and P is any point within it; shew that the sum of the triangles PAB , PCD is equal to half the parallelogram.

PROPOSITION 42. PROBLEM.

To describe a parallelogram that shall be equal to a given triangle, and have one of its angles equal to a given angle.



Let ABC be the given triangle, and D the given angle.
It is required to describe a parallelogram equal to ABC , and having one of its angles equal to D .

Construction. Bisect BC at E . I. 10.

At E in CE , make the angle CEF equal to D ; I. 23.

through A draw AFG parallel to EC ; I. 31.

and through C draw CG parallel to EF .

Then $FECG$ shall be the parallelogram required.

Join AE .

Proof. Now the triangles ABE , AEC are equal,
for they are on equal bases BE , EC , and between the same
parallels; I. 38.

therefore the triangle ABC is double of the triangle AEC .

But $FECG$ is a parallelogram by construction: *Def.* 26.

and it is double of the triangle AEC ,

for they are on the same base EC , and between the same
parallels EC and AG . I. 41.

Therefore the parallelogram $FECG$ is equal to the triangle
 ABC ;

and it has one of its angles CEF equal to the given angle D .

Q. E. F.

EXERCISES.

1. Describe a parallelogram equal to a given square standing on the same base, and having an angle equal to half a right angle.

2. Describe a rhombus equal to a given parallelogram and standing on the same base. When does the construction fail?


DEFINITION. (If in the diagonal of a parallelogram any point is taken, and straight lines are drawn through it parallel to the sides of the parallelogram; then of the four parallelograms into which the whole figure is divided, the two through which the diagonal passes are called **Parallelograms about that diagonal**, and the other two, which with these make up the whole figure, are called the **complements** of the parallelograms about the diagonal.)

Thus in the figure given below, $AEKH$, $KGCF$ are parallelograms about the diagonal AC ; and $HKFD$, $EBGK$ are the complements of those parallelograms.

NOTE. A parallelogram is often named by *two* letters only, these being placed at opposite angular points.

PROPOSITION 43. THEOREM.

The complements of the parallelograms about the diagonal of any parallelogram, are equal to one another.



Let $ABCD$ be a parallelogram, and KD , KB the complements of the parallelograms EH , GF about the diagonal AC : then shall the complement BK be equal to the complement KD .

Proof. Because EH is a parallelogram, and AK its diagonal, therefore the triangle AEK is equal to the triangle AHK . I. 34.

• For a similar reason the triangle KGC is equal to the triangle KFC .

Hence the triangles AEK , KGC are together equal to the triangles AHK , KFC .

But the whole triangle ABC is equal to the whole triangle ADC , for AC bisects the parallelogram $ABCD$; I. 34. therefore the remainder, the complement BK , is equal to the remainder, the complement KD . Q.E.D.


EXERCISES.

In the figure of Prop. 43, prove that

- (i) The parallelogram ED is equal to the parallelogram BH .
- (ii) If KB , KD are joined, the triangle AKB is equal to the triangle AKD .

PROPOSITION 44. PROBLEM.

To a given straight line to apply a parallelogram which shall be equal to a given triangle, and have one of its angles equal to a given angle.




Let AB be the given straight line, C the given triangle, and D the given angle.

It is required to apply to the straight line AB a parallelogram equal to the triangle C , and having an angle equal to the angle D .

Construction. On AB produced describe a parallelogram $BEFG$ equal to the triangle C , and having the angle EBG equal to the angle D ; I. 22 and I. 42*. through A draw AH parallel to BG or EF , to meet FG produced in H . I. 31.

Join HB .

* This step of the construction is effected by first describing on AB produced a triangle whose sides are respectively equal to those of the triangle C (I. 22); and by then making a parallelogram equal to the triangle so drawn, and having an angle equal to D (I. 42).



Then because AH and EF are parallel, and HF meets them, therefore the angles AHF, HFE are together equal to two right angles: I. 29.

hence the angles BHF, HFE are together less than two right angles;

therefore HB and FE will meet if produced towards B and E. Ac. 12.

Produce them to meet at K.

Through K draw KL parallel to EA or FH; I. 31.
and produce HA, GB to meet KL in the points L and M.

Then shall BL be the parallelogram required.

Proof. Now FHLK is a parallelogram, *Constr.*
and LB, BF are the complements of the parallelograms about the diagonal HK:

therefore LB is equal to BF. I. 43.

But the triangle C is equal to BF; *Constr.*
therefore LB is equal to the triangle C.


And because the angle GBE is equal to the vertically opposite angle ABM, I. 15.

and is likewise equal to the angle D; *Constr.*
therefore the angle ABM is equal to the angle D.

Therefore the parallelogram LB, which is applied to the straight line AB, is equal to the triangle C, and has the angle ABM equal to the angle D. Q.E.F.

PROPOSITION 45. PROBLEM.

To describe a parallelogram equal to a given rectilineal figure, and having an angle equal to a given angle.



Let $ABCD$ be the given rectilineal figure, and E the given angle.

It is required to describe a parallelogram equal to $ABCD$, and having an angle equal to E .

Suppose the given rectilineal figure to be a quadrilateral.

Construction. Join BD .

Describe the parallelogram FH equal to the triangle ABD , and having the angle FKH equal to the angle E . I. 42.

To GH apply the parallelogram GM , equal to the triangle DBC , and having the angle GHM equal to E . I. 44.

Then shall $FKML$ be the parallelogram required.


Proof. Because each of the angles GHM , FKH is equal to E , therefore the angle FKH is equal to the angle GHM .

To each of these equals add the angle GKH ; then the angles FKH , GKH are together equal to the angles GHM , GKH .

But since FK , GH are parallel, and KH meets them, therefore the angles FKH , GKH are together equal to two right angles: I. 29.

therefore also the angles GHM , GKH are together equal to two right angles:

therefore KH , HM are in the same straight line. I. 14.



Again, because KM , FG are parallel, and HG meets them, therefore the alternate angles MHG , HGF are equal : I. 29
to each of these equals add the angle HGL ;
then the angles MHG , HGL are together equal to the angles HGF , HGL .

But because HM , GL are parallel, and HG meets them, therefore the angles MHG , HGL are together equal to two right angles : I. 29.
therefore also the angles HGF , HGL are together equal to two right angles :

therefore FG , GL are in the same straight line. I. 14.

And because KF and ML are each parallel to HG , *Constr.*

therefore KF is parallel to ML ; I. 30.

and KM , FL are parallel ; *Constr.*

therefore $FKML$ is a parallelogram. *Def. 26.*

And because the parallelogram PH is equal to the triangle ABD , *Constr.*

and the parallelogram GM to the triangle DBC ; *Constr.*
therefore the whole parallelogram $FKML$ is equal to the whole figure $ABCD$;


and it has the angle FKM equal to the angle E .

By a series of similar steps, a parallelogram may be constructed equal to a rectilineal figure of more than four sides.

Q.E.F.

PROPOSITION 46. PROBLEM.

To describe a square on a given straight line.



Let AB be the given straight line :
it is required to describe a square on AB .

Constr. From A draw AC at right angles to AB ; 1. 11.
and make AD equal to AB . 1. 3.

Through D draw DE parallel to AB ; 1. 31.
and through B draw BE parallel to AD , meeting DE in E .
Then shall $ADEB$ be a square.

Proof. For, by construction, $ADEB$ is a parallelogram :
therefore AB is equal to DE , and AD to BE . 1. 31.

But AD is equal to AB ; *Constr.*
therefore the four straight lines AB , AD , DE , EB are equal
to one another ;

that is, the figure $ADEB$ is equilateral.

Again, since AB , DE are parallel, and AD meets them,
therefore the angles BAD , ADE are together equal to two
right angles ; 1. 29.

but the angle BAD is a right angle ; *Constr.*

therefore also the angle ADE is a right angle.

And the opposite angles of a parallelogram are equal ; 1. 34.

therefore each of the angles DEB , EBA is a right angle : •

that is the figure $ADEB$ is rectangular. •


Hence it is a square, and it is described on AB .

• Q.E.F.

COROLLARY. *If one angle of a parallelogram is a right angle, all its angles are right angles.*

PROPOSITION 47. THEOREM.

In a right-angled triangle the square described on the hypotenuse is equal to the sum of the squares described on the other two sides.



Let ABC be a right-angled triangle, having the angle BAC a right angle :
then shall the square described on the hypotenuse BC be equal to the sum of the squares described on BA , AC .

Construction. On BC describe the square $BDEC$; i. 46.
and on BA , AC describe the squares $BAGF$, $ACKH$.
Through A draw AL parallel to BD or CE ; i. 31.
and join AD , FC .

Proof. Then because each of the angles BAC , BAG is a right angle,
therefore CA and AG are in the same straight line. i. 14.

Now the angle CBD is equal to the angle FBA ,
for each of them is a right angle.

Add to each the angle ABC :
then the whole angle ABD is equal to the whole angle FBC .

Then in the triangles ABD, FBC,
 Because $\left\{ \begin{array}{l} \text{AB is equal to FB,} \\ \text{and BD is equal to BC,} \\ \text{also the angle ABD is equal to the angle FBC;} \end{array} \right.$
 therefore the triangle ABD is equal to the triangle FBC. I. 4.

Now the parallelogram BL is double of the triangle ABD, for they are on the same base BD, and between the same parallels BD, AL. I. 41.

And the square GB is double of the triangle FBC, for they are on the same base FB, and between the same parallels FB, GC. I. 41.

But doubles of equals are equal: Ax. 6.
 therefore the parallelogram BL is equal to the square GB.

In a similar way, by joining AE, BK, it can be shewn that the parallelogram CL is equal to the square CH.

Therefore the whole square BE is equal to the sum of the squares GB, HC:

that is, the square described on the hypotenuse BC is equal to the sum of the squares described on the two sides BA, AC. Q.E.D.

NOTE. It is not necessary to the proof of this Proposition that the three squares should be described *external* to the triangle ABC; and since *each* square may be drawn either *towards* or *away from* the triangle, it may be shewn that there are $2 \times 2 \times 2$, or *eight*, possible constructions.

EXERCISES.

In the figure of this Proposition, shew that

- (i) If BG, CH are joined, these straight lines are parallel;
- (ii) The points F, A, K are in one straight line;
- (iii) FC and AD are at right angles to one another;
- (iv) If GH, KE, FD are joined, the triangle GAH is equal to the given triangle in all respects; and the triangles FBD, KCE are each equal in area to the triangle ABC.

[See Ex. 9, p. 73.]

2. On the sides AB , AC of *any* triangle ABC , squares $ABFG$, $ACKH$ are described both toward the triangle, or both on the side remote from it: shew that the straight lines BH and CG are equal.

3. On the sides of any triangle ABC , equilateral triangles BCX , CAY , ABZ are described, all externally, or all towards the triangle: shew that AX , BY , CZ are all equal.

4. The square described on the diagonal of a given square, is double of the given square.


5. ABC is an equilateral triangle, and AX is the perpendicular drawn from A to BC : shew that the square on AX is three times the square on BX .

6. Describe a square equal to the sum of two given squares.

7. From the vertex A of a triangle ABC , AX is drawn perpendicular to the base: shew that the difference of the squares on the sides AB and AC , is equal to the difference of the squares on BX and CX , the segments of the base.

8. If from any point O within a triangle ABC , perpendiculars OX , OY , OZ are drawn to the sides BC , CA , AB respectively; shew that the sum of the squares on the segments AZ , BX , CY is equal to the sum of the squares on the segments AY , CX , BZ .

PROPOSITION 47. ALTERNATIVE PROOF.



Let CAB be a right-angled triangle, having the angle at A a right angle:

then shall the square on the hypotenuse BC be equal to the sum of the squares on BA , AC .

On AB describe the square $ABFG$. I. 46.
 From FG and GA cut off respectively FD and GK , each equal to AC . I. 3.

On GK describe the square $GKEH$: I. 46.
 then HG and GF are in the same straight line. I. 11.

Join CE , ED , DB .

It will first be shewn that the figure $CEDB$ is the square on CB .

Now CA is equal to KG ; add to each AK :

therefore CK is equal to AG .

Similarly DH is equal to GF :

hence the four lines BA , CK , DH , BF are all equal.

Then in the triangles BAC , CKE ,

Because $\left\{ \begin{array}{l} BA \text{ is equal to } CK, \\ \text{and } AC \text{ is equal to } KE; \\ \text{also the contained angle } BAC \text{ is equal to the contained} \\ \text{angle } CKE, \text{ being right angles;} \end{array} \right.$ Proved.
Constr.

therefore the triangles BAC , CKE are equal in all respects. I. 4.

Similarly the four triangles BAC , CKE , DHE , BFD may be shewn to be equal in all respects.

Therefore the four straight lines BC , CE , ED , DB are all equal;

that is, the figure $CEDB$ is equilateral.

Again the angle CBA is equal to the angle DBF ; Proved.

add to each the angle ABD :

then the angle CBD is equal to the angle ABF :

therefore the angle CBD is a right angle.

Hence the figure $CEDB$ is the square on BC . Def. 28.

And $EHGK$ is equal to the square on AC . Constr.

Now the square $CEDB$ is made up of the two triangles BAC , CKE , and the rectilinear figure $AKEDB$;

therefore the square $CEDB$ is equal to the triangles EHD , DFB together with the same rectilinear figure;

but these make up the squares $EHGK$, $AGFB$:

hence the square $CEDB$ is equal to the sum of the squares $EHGK$, $AGFB$:


that is, the square on the hypotenuse BC is equal to the sum of the squares on the two sides CA , AB . Q. E. D.

Obs. The following properties of a square, though not formally enunciated by Euclid, are employed in subsequent proofs. [See I. 48.]

- (i) *The squares on equal straight lines are equal.*
- (ii) *Equal squares stand upon equal straight lines.*

PROPOSITION 48. THEOREM.

If the square described on one side of a triangle be equal to the sum of the squares described on the other two sides, then the angle contained by these two sides shall be a right angle.



Let ABC be a triangle; and let the square described on BC be equal to the sum of the squares described on BA , AC :
then shall the angle BAC be a right angle.

Construction. From A draw AD at right angles to AC ; 1. 11.
and make AD equal to AB . 1. 3.
Join DC .

Proof. Then, because AD is equal to AB , *Constr.*
therefore the square on AD is equal to the square on AB .

To each of these add the square on CA ;
then the sum of the squares on CA , AD is equal to the sum
of the squares on CA , AB .

But, because the angle DAC is a right angle, *Constr.*
therefore the square on DC is equal to the sum of the
squares on CA , AD . 1. 47.

And, by hypothesis, the square on BC is equal to the sum
of the squares on CA , AB ;

therefore the square on DC is equal to the square on BC :
therefore also the side DC is equal to the side BC .

Then in the triangles DAC , BAC ,

Because $\left\{ \begin{array}{l} DA \text{ is equal to } BA, \\ \text{and } AC \text{ is common to both;} \\ \text{also the third side } DC \text{ is equal to the third side} \\ BC; \end{array} \right. \begin{array}{l} \text{Constr.} \\ \\ \text{Proved.} \end{array}$

therefore the angle DAC is equal to the angle BAC . 1. 8.

But DAC is a right angle; *Constr.*
therefore also BAC is a right angle. Q. E. D.

THEOREMS AND EXAMPLES ON BOOK I.

INTRODUCTORY.

HINTS TOWARDS THE SOLUTION OF GEOMETRICAL EXERCISES. ANALYSIS. SYNTHESIS.

It is commonly found that exercises in Pure Geometry present to a beginner far more difficulty than examples in any other branch of Elementary Mathematics. This seems to be due to the following causes.

(i) The main Propositions in the text of Euclid must be not merely understood, but thoroughly digested, before the exercises depending upon them can be successfully attempted.

(ii) The variety of such exercises is practically unlimited; and it is impossible to lay down for their treatment any definite methods, such as the student has been accustomed to find in the rules of Elementary Arithmetic and Algebra.

(iii) The arrangement of Euclid's Propositions, though perhaps the most *convincing* of all forms of argument, affords in most cases little clue as to the way in which the proof or construction *was discovered*.

Euclid's propositions are arranged **synthetically**: that is to say, they start from the hypothesis or data; they next proceed to a construction in accordance with postulates, and problems already solved; then by successive steps based on known theorems, they finally establish the result indicated by the enunciation.

Thus Geometrical Synthesis is a *building up* of known results, in order to obtain a *new* result.

But as this is not the way in which constructions or proofs are usually discovered, we draw the attention of the student to the following hints.


Begin by *assuming* the result it is desired to establish; then by working backwards, trace the consequences of the assumption, and try to ascertain its dependence on some simpler theorem which is already known to be true, or on some condition which suggests the necessary construction. If this attempt is successful, the steps of the argument may in general be re-arranged in reverse order, and the construction and proof presented in a synthetic form.

This unravelling of the conditions of a proposition in order to trace it back to some earlier principle on which it depends, is called **geometrical analysis**: it is the natural way of attacking most exercises of a more difficult type, and it is especially adapted to the solution of *problems*.

These directions are so general that they cannot be said to amount to a *method*: all that can be claimed for Geometrical Analysis is that it furnishes a mode of *searching for a suggestion*, and its success will necessarily depend on the skill and ingenuity with which it is employed: these may be expected to come with experience, but a thorough grasp of the chief Propositions of Euclid is essential to attaining them.

The practical application of these hints is illustrated by the following examples.

1. *Construct an isosceles triangle having given the base, and the sum of one of the equal sides and the perpendicular drawn from the vertex to the base.*



Let AB be the given base, and K the sum of one side and the perpendicular drawn from the vertex to the base.

ANALYSIS. Suppose ABC to be the required triangle.

From C draw CX perpendicular to AB :

then AB is bisected at X .

I. 26.

Now if we produce CX to H , making XH equal to K ,
it follows that $CH = CA$;

and if AH is joined,

we notice that the angle $CAH =$ the angle CHA .

I. 5.

Now the straight lines XH and AH can be drawn *before the position of C is known*;

Hence we have the following construction, which we arrange synthetically.

SYNTHESIS. Bisect AB at X :
from X draw XH perpendicular to AB , making XH equal to K .
Join AH .

At the point A in HA , make the angle HAC equal to the angle AHX ; and join CB .


Then ACB shall be the triangle required.

First, the triangle is isosceles, for $AC = BC$. I. 4.
Again, since the angle $HAC =$ the angle AHX , Constr.
 $\therefore HC = AC$. I. 6.

To each add CX ;
then the sum of AC , $CX =$ the sum of HC , CX .
 $= HX$.

That is, the sum of AC , $CX = K$. Q. E. F.

2. To divide a given straight line so that the square on one part may be double of the square on the other.



Let AB be the given straight line.

ANALYSIS. Suppose AB to be divided as required at X : that is, suppose the square on AX to be double of the square on XB .

Now we remember that in an isosceles right-angled triangle, the square on the hypotenuse is double of the square on either of the equal sides.

This suggests to us to draw BC perpendicular to AB , and to make BC equal to BX .

Join XC .

Then the square on XC is double of the square on XB , I. 47.
 $\therefore XC = AX$.

And when we join AC , we notice that
the angle $XAC =$ the angle XCA . I. 5.

Hence the exterior angle CXB is double of the angle XAC . I. 32.

But the angle CXB is half of a right angle : I. 32.
 \therefore the angle XAC is one-fourth of a right angle.

This supplies the clue to the following construction :—

SYNTHESIS. From **B** draw **BD** perpendicular to **AB** ;
 and from **A** draw **AC**, making **BAC** one-fourth of a right angle.
 From **C**, the intersection of **AC** and **BD**, draw **CX**, making the angle \angle
ACX equal to the angle **BAC**. I. 23.
 Then **AB** shall be divided as required at **X**.
 For since the angle **XCA** = the angle **XAC**,
 \therefore **XA** = **XC**. I. 6.
 And because the angle **BXC** = the sum of the angles **BAC**, **ACX**, I. 32.
 \therefore the angle **BXC** is half a right angle ;
 and the angle at **B** is a right angle ;
 therefore the angle **BCX** is half a right angle ; I. 32.
 \therefore the angle **BXC** = the angle **BCX** ;
 \therefore **BX** = **BC**.
 Hence the square on **XC** is double of the square on **XB** : I. 47.
 that is, the square on **AX** is double of the square on **XB**. Q. E. F.

I. ON THE IDENTICAL EQUALITY OF TRIANGLES.

See Propositions 4, 8, 26.

1. If in a triangle the perpendicular from the vertex on the base bisects the base, then the triangle is isosceles.

2. If the bisector of the vertical angle of a triangle is also perpendicular to the base, the triangle is isosceles.

3. If the bisector of the vertical angle of a triangle also bisects the base, the triangle is isosceles.

[Produce the bisector, and complete the construction after the manner of I. 16.]

4. If in a triangle a pair of straight lines drawn from the extremities of the base, making equal angles with the sides, are equal, the triangle is isosceles.

5. If in a triangle the perpendiculars drawn from the extremities of the base to the opposite sides are equal, the triangle is isosceles.

6. Two triangles **ABC**, **ABD** on the same base **AB**, and on opposite sides of it, are such that **AC** is equal to **AD**, and **BC** is equal to **BD** : shew that the line joining the points **C** and **D** is perpendicular to **AB**.

7. If from the extremities of the base of an isosceles triangle perpendiculars are drawn to the opposite sides, shew that the straight line joining the vertex to the intersection of these perpendiculars bisects the vertical angle.


8. ABC is a triangle in which the vertical angle BAC is bisected by the straight line AX : from B draw BD perpendicular to AX , and produce it to meet AC , or AC produced, in E ; then shew that BD is equal to DE .

9. In a quadrilateral $ABCD$, AB is equal to AD , and BC is equal to DC : shew that the diagonal AC bisects each of the angles which it joins.

10. In a quadrilateral $ABCD$ the opposite sides AD , BC are equal, and also the diagonals AC , BD are equal: if AC and BD intersect at K , shew that each of the triangles AKB , DKC is isosceles.

11. If one angle of a triangle be equal to the sum of the other two, the greatest side is double of the distance of its middle point from the opposite angle.

12. Two right-angled triangles which have their hypotenuses equal, and one side of one equal to one side of the other, are identically equal.



Let ABC , DEF be two Δ^s right-angled at B and E , having AC equal to DF , and AB equal to DE :

then shall the Δ^s be identically equal.

For apply the Δ ABC to the Δ DEF , so that A may fall on D , and AB along DE ; and so that C may fall on the side of DE remote from F .

Let C' be the point on which C falls.

Then since $AB = DE$,

$\therefore B$ must fall on E ;

so that DEC' represents the Δ ABC in its new position.

Now each of the \angle^s DEF , DEC' is a rt. \angle ;

$\therefore EF$ and EC' are in one st. line.

Hyp.
I. 11.

Then in the Δ $C'DF$,

because $DF = DC'$,

\therefore the \angle $DFC' =$ the \angle $DC'F$.

Hence in the two Δ^s DEF , DEC' ,

the \angle $DEF =$ the \angle DEC' , being rt. \angle^s ;

Because { and the \angle $DFE =$ the \angle $DC'E$;
also the side DE is common to both;

Proved.


\therefore the Δ^s DEF , DEC' are equal in all respects;

I. 26.

that is, the Δ^s DEF , ABC are equal in all respects.

Q.E.D.

13. If two triangles have two sides of the one equal to two sides of the other, each to each, and have likewise the angles opposite to one pair of equal sides equal, then the angles opposite to the other pair of equal sides are either equal or supplementary, and in the former case the triangles are equal in all respects.



Let ABC , DEF be two triangles, having the side AB equal to the side DE , the side AC equal to the side DF , and the $\angle ABC$ equal to the $\angle DEF$; then shall the $\angle^s ACB$, DFE be either equal or supplementary, and in the former case the triangles shall be equal in all respects.

If the $\angle BAC$ — the $\angle EDF$,
then the triangles are equal in all respects. I. 4.

But if the $\angle BAC$ be not equal to the $\angle EDF$, one of them must be the greater.

Let the $\angle EDF$ be greater than the $\angle BAC$.

At D in ED make the $\angle EDF'$ equal to the $\angle BAC$.

Then the $\triangle^s BAC$, EDF' are equal in all respects. I. 26.

$\therefore AC = DF'$;

but $AC = DF$;

Hyp.

$\therefore DF = DF'$;

\therefore the $\angle DFF' =$ the $\angle DF'E$. I. 5.

But the $\angle^s DF'E$, $DF'E$ are supplementary, I. 13.

\therefore the $\angle^s DFF'$, $DF'E$ are supplementary:

that is, the $\angle^s DFE$, ACB are supplementary.

Q. E. D.

Three cases of this theorem deserve special attention.

It has been proved that if the angles ACB , DFE are not *equal*, they are *supplementary* :

And we know that of angles which are supplementary and unequal, one must be acute and the other obtuse. *

COROLLARIES. Hence, in addition to the hypothesis of this theorem,

- (i) If the angles ACB , DFE , opposite to the two equal sides AB , DE are both acute, both obtuse, or if one of them is a right angle,
it follows that these angles are equal,
and therefore that the triangles are equal in all respects.
- (ii) If the two given angles are right angles or obtuse angles,
it follows that the angles ACB , DFE must be both acute, and therefore equal, by (i) :
so that the triangles are equal in all respects.
- (iii) If in each triangle the side opposite the given angle is not less than the other given side ; that is, if AC and DF are not less than AB and DE respectively, then
the angles ACB , DFE cannot be greater than the angles ABC , DEF respectively ;
therefore the angles ACB , DFE , are both acute ;
hence, as above, they are equal ;
and the triangles ABC , DEF are equal in all respects.

• II. ON INEQUALITIES.

See Propositions 16, 17, 18, 19, 20, 21, 24, 25.

1. In a triangle ABC , if AC is not greater than AB , shew that any straight line drawn through the vertex A , and terminated by the base BC , is less than AB .

2. ABC is a triangle, and the vertical angle BAC is bisected by a straight line which meets the base BC in X ; shew that BA is greater than BX , and CA greater than CX . Hence obtain a proof of I. 20.

3. The perpendicular is the shortest straight line that can be drawn from a given point to a given straight line ; and of others, that which is nearer to the perpendicular is less than the more remote ; and two, and only two equal straight lines can be drawn from the given point to the given straight line, one on each side of the perpendicular.

4. The sum of the distances of any point from the three angular points of a triangle is greater than half its perimeter.

5. The sum of the distances of any point within a triangle from its angular points is less than the perimeter of the triangle.

6. The perimeter of a quadrilateral is greater than the sum of its diagonals.

7. The sum of the diagonals of a quadrilateral is less than the sum of the four straight lines drawn from the angular points to any given point. Prove this, and point out the exceptional case.

8. In a triangle any two sides are together greater than twice the median which bisects the remaining side. [See Def. p. 73.]

[Produce the median, and complete the construction after the manner of I. 16.]

9. In any triangle the sum of the medians is less than the perimeter.

10. In a triangle an angle is acute, obtuse, or a right angle, according as the median drawn from it is greater than, less than, or equal to half the opposite side. [See Ex. 4, p. 59.]

11. The diagonals of a rhombus are unequal.

12. If the vertical angle of a triangle is contained by unequal sides, and if from the vertex the median and the bisector of the angle are drawn, then the median lies within the angle contained by the bisector and the longer side.

Let ABC be a Δ , in which AB is greater than AC ; let AX be the median drawn from A , and AP the bisector of the vertical $\angle BAC$:

then shall AX lie between AP and AB .

Produce AX to K , making XK equal to AX . Join KC .

Then the $\Delta^s BXA, CXK$ may be shewn to be equal in all respects: I. 4.
hence $BA = CK$, and the $\angle BAX = \angle CKX$.

But since BA is greater than AC , Hyp.


$\therefore CK$ is greater than AC ;

\therefore the $\angle CAK$ is greater than the $\angle CKA$: I. 18.

that is, the $\angle CAX$ is greater than the $\angle BAX$:

\therefore the $\angle CAX$ must be more than half the vert. $\angle BAC$;

hence AX lies within the angle BAP . Q.E.D.



13. If two sides of a triangle are unequal, and if from their point of intersection three straight lines are drawn, namely the bisector of the vertical angle, the median, and the perpendicular to the base, the first is intermediate in position and magnitude to the other two.

III. ON PARALLELS.

See Propositions 27—31.

1. If a straight line meets two parallel straight lines, and the two interior angles on the same side are bisected; shew that the bisectors meet at right angles. [r. 29, r. 32.]

2. The straight lines drawn from any point in the bisector of an angle parallel to the arms of the angle, and terminated by them, are equal; and the resulting figure is a rhombus.

3. AB and CD are two straight lines intersecting at D , and the adjacent angles so formed are bisected: if through any point X in DC a straight line YXZ be drawn parallel to AB and meeting the bisectors in Y and Z , shew that XY is equal to XZ .

4. If two straight lines are parallel to two other straight lines, each to each; and if the angles contained by each pair are bisected; shew that the bisecting lines are parallel.

5. The middle point of any straight line which meets two parallel straight lines, and is terminated by them, is equidistant from the parallels.

6. A straight line drawn between two parallels and terminated by them, is bisected; shew that any other straight line passing through the middle point and terminated by the parallels, is also bisected at that point.

7. If through a point equidistant from two parallel straight lines, two straight lines are drawn cutting the parallels, the portions of the latter thus intercepted are equal.

PROBLEMS.

8. AB and CD are two given straight lines, and X is a given point in AB : find a point Y in AB such that YX may be equal to the perpendicular distance of Y from CD .

9. ABC is an isosceles triangle; required to draw a straight line DE parallel to the base BC , and meeting the equal sides in D and E , so that BD , DE , EC may be all equal.

10. ABC is any triangle; required to draw a straight line DE parallel to the base BC , and meeting the other sides in D and E , so that DE may be equal to the sum of BD and CE .

11. ABC is any triangle; required to draw a straight line parallel to the base BC , and meeting the other sides in D and E , so that DE may be equal to the difference of BD and CE .

IV. ON PARALLELOGRAMS.

See Propositions 33, 34, and the deductions from these Props. given on page 64.

1. *The straight line drawn through the middle point of a side of a triangle parallel to the base, bisects the remaining side.*

Let ABC be a \triangle , and Z the middle point of the side AB . Through Z , ZY is drawn par^l to BC ; then shall Y be the middle point of AC .

Through Z draw ZX par^l to AC . I. 31.

Then in the \triangle s AZY , ZBX ,
because ZY and BC are par^l,
 \therefore the $\angle AZY =$ the $\angle ZBX$; I. 29
and because ZX and AC are par^l,
 \therefore the $\angle ZAY =$ the $\angle BZX$; I. 29.
also $AZ = ZB$; Hyp.

$\therefore AY = ZX$.

But $ZXCY$ is a parm by construction;

$\therefore ZX = YC$.


Hence $AY = YC$;

that is, AC is bisected at Y .

I. 26.

I. 34.

Q.E.D.



2. *The straight line which joins the middle points of two sides of a triangle, is parallel to the third side.*

Let ABC be a \triangle , and Z , Y the middle points of the sides AB , AC :

then shall ZY be par^l to BC .

Produce ZY to V , making YV equal to ZY .

Join CV .

Then in the \triangle s AYZ , CYV ,

Because { $AY = CY$, Hyp.
and $YZ = YV$, Constr.
and the $\angle AYZ =$ the vert. opp. $\angle CYV$;

I. 15.

I. 4.

$\therefore AZ = CV$,
and the $\angle ZAY =$ the $\angle VCY$;

hence CV is par^l to AZ .

I. 27.

But CV is equal to AZ , that is, to BZ ;

Hyp.


$\therefore CV$ is equal and par^l to BZ ;

$\therefore ZV$ is equal and par^l to BC ;

I. 33.

that is, ZY is par^l to BC .

Q.E.D.



[A second proof of this proposition may be derived from I. 38, 39.]

3. *The straight line which joins the middle points of two sides of a triangle is equal to half the third side.*

4. *Shew that the three straight lines which join the middle points of the sides of a triangle, divide it into four triangles which are identically equal.*

5. *Any straight line drawn from the vertex of a triangle to the base is bisected by the straight line which joins the middle points of the other sides of the triangle.*

6. *Given the three middle points of the sides of a triangle, construct the triangle.*

7. *AB, AC are two given straight lines, and P is a given point between them; required to draw through P a straight line, terminated by AB, AC, and bisected by P.*

8. *ABCD is a parallelogram, and X, Y are the middle points of the opposite sides AD, BC: shew that BX and DY trisect the diagonal AC.*

9. *If the middle points of adjacent sides of any quadrilateral be joined, the figure thus formed is a parallelogram.*

10. *Shew that the straight lines which join the middle points of opposite sides of a quadrilateral, bisect one another.*

11. *The straight line which joins the middle points of the oblique sides of a trapezium, is parallel to the two parallel sides, and passes through the middle points of the diagonals.*


12. *The straight line which joins the middle points of the oblique sides of a trapezium is equal to half the sum of the parallel sides; and the portion intercepted between the diagonals is equal to half the difference of the parallel sides.*

Definition. If from the extremities of one straight line perpendiculars are drawn to another, the portion of the latter intercepted between the perpendiculars is said to be the **Orthogonal Projection** of the first line upon the second.




Thus in the adjoining figures, if from the extremities of the straight line AB the perpendiculars AX, BY are drawn to PQ, then XY is the *orthogonal projection* of AB on PQ.

13. A given straight line AB is bisected at C ; shew that the projections of AC , CB on any other straight line are equal.



Q



Let XZ , ZY be the projections of AC , CB on any straight line PQ : then XZ and ZY shall be equal.

Through A draw a straight line parallel to PQ , meeting CZ , BY or these lines produced, in H , K . 1. 31.

Now AX , CZ , BY are parallel, for they are perp. to PQ ; 1. 28.

\therefore the figures XH , HY are paral^l;

$\therefore AH = XZ$, and $HK = ZY$. 1. 31.

But through C , the middle point of AB , a side of the $\triangle ABK$, CH has been drawn parallel to the side BK ;

$\therefore CH$ bisects AK :

Ex. 1, p. 96.

that is, $AH = HK$;

$\therefore XZ = ZY$.

Q. E. D.

11. If three parallel straight lines make equal intercepts on a fourth straight line which meets them, they will also make equal intercepts on any other straight line which meets them.

15. Equal and parallel straight lines have equal projections on any other straight line.

16. AB is a given straight line bisected at O ; and AX , BY are perpendiculars drawn from A and B on any other straight line: shew that OX is equal to OY .

17. AB is a given straight line bisected at O : and AX , BY and OZ are perpendiculars drawn to any straight line PQ , which does not pass between A and B : shew that OZ is equal to half the sum of AX , BY .

[OZ is said to be the **Arithmetic Mean** between AX and BY .]

18. AB is a given straight line bisected at O ; and through A , B and O parallel straight lines are drawn to meet a given straight line PQ in X , Y , Z : shew that OZ is equal to half the sum, or half the difference of AX and BY , according as A and B lie on the same side or on opposite sides of PQ .

19. *To divide a given finite straight line into any number of equal parts.*


[For example, required to divide the straight line AB into *five* equal parts.

From A draw AC , a straight line of unlimited length, making any angle with AB .

In AC take *any* point P , and mark off successive parts PQ , QR , RS , ST each equal to AP .

Join BT ; and through P , Q , R , S draw parallels to BT .

It may be shewn by Ex. 14, p. 98, that these parallels divide AB into five equal parts.]



20. *If through an angle of a parallelogram any straight line is drawn, the perpendicular drawn to it from the opposite angle is equal to the sum or difference of the perpendiculars drawn to it from the two remaining angles, according as the given straight line falls without the parallelogram, or intersects it.*

[Through the opposite angle draw a straight line parallel to the given straight line, so as to meet the perpendicular from one of the remaining angles, produced if necessary: then apply i. 34, i. 26. Or proceed as in the following example.]

21. From the angular points of a parallelogram perpendiculars are drawn to any straight line which is without the parallelogram: shew that the sum of the perpendiculars drawn from one pair of opposite angles is equal to the sum of those drawn from the other pair.

[Draw the diagonals, and from their point of intersection let fall a perpendicular upon the given straight line. See Ex. 17, p. 98.]

22. The sum of the perpendiculars drawn from any point in the base of an isosceles triangle to the equal sides is equal to the perpendicular drawn from either extremity of the base to the opposite side.

[It follows that the sum of the distances of *any* point in the base of an isosceles triangle from the equal sides is **constant**, that is, the same whatever point in the base is taken.]

23. In the base produced of an isosceles triangle any point is taken: shew that the difference of its distances from the equal sides is *constant*.

24. The sum of the perpendiculars drawn from any point within an equilateral triangle to the three sides is equal to the perpendicular drawn from any one of the angular points to the opposite side, and is therefore constant.

PROBLEMS.

[Problems marked (*) admit of more than one solution.]

*25. Draw a straight line through a given point, so that the part of it intercepted between two given parallel straight lines may be of given length.

26. Draw a straight line parallel to a given straight line, so that the part intercepted between two other given straight lines may be of given length.

27. Draw a straight line equally inclined to two given straight lines that meet, so that the part intercepted between them may be of given length.

28. AB, AC are two given straight lines, and P is a given point *without* the angle contained by them. It is required to draw through P a straight line to meet the given lines, so that the part intercepted between them may be equal to the part between P and the nearer line.

V. MISCELLANEOUS THEOREMS AND EXAMPLES.

Chiefly on I. 32.

1. A is the vertex of an isosceles triangle ABC , and BA is produced to D , so that AD is equal to BA ; if DC is drawn, shew that BCD is a right angle.

2. The straight line joining the middle point of the hypotenuse of a right-angled triangle to the right angle is equal to half the hypotenuse.

3. From the extremities of the base of a triangle perpendiculars are drawn to the opposite sides (produced if necessary); shew that the straight lines which join the middle point of the base to the feet of the perpendiculars are equal.

4. In a triangle ABC , AD is drawn perpendicular to BC ; and X, Y, Z are the middle points of the sides BC, CA, AB respectively: shew that each of the angles ZXY, ZDY is equal to the angle BAC .

5. In a right-angled triangle, if a perpendicular be drawn from the right angle to the hypotenuse, the two triangles thus formed are equiangular to one another.


6. In a right-angled triangle two straight lines are drawn from the right angle, one bisecting the hypotenuse, the other perpendicular to it: shew that they contain an angle equal to the difference of the two acute angles of the triangle. [See above, Ex. 2 and Ex. 5.]

7. In a triangle if a perpendicular be drawn from one extremity of the base to the bisector of the vertical angle, (i) it will make with either of the sides containing the vertical angle an angle equal to half the sum of the angles at the base; (ii) it will make with the base an angle equal to half the difference of the angles at the base.

Let ABC be the given Δ , and AH the bisector of the vertical $\angle BAC$.

Let CLK meet AH at right angles.

(i) Then shall each of the $\angle^s AKC, ACK$ be equal to half the sum of the $\angle^s ABC, ACB$.



In the $\Delta^s AKL, ACL$,
 Because $\left\{ \begin{array}{l} \text{the } \angle KAL = \text{the } \angle CAL, \\ \text{also the } \angle ALK = \text{the } \angle ALC, \text{ being rt. } \angle^s; \\ \text{and } AL \text{ is common to both } \Delta^s; \end{array} \right.$
 $\therefore \text{the } \angle AKL = \text{the } \angle ACL.$ I. 26.

Again, the $\angle AKC = \text{the sum of the } \angle^s KBC, KCB$; I. 32.
 that is, the $\angle ACK = \text{the sum of the } \angle^s KBC, KCB$.

To each add the $\angle ACK$,
 then twice the $\angle ACK = \text{the sum of the } \angle^s ABC, ACB$,
 $\therefore \text{the } \angle ACK = \text{half the sum of the } \angle^s ABC, ACB$.

(ii) The $\angle KCB$ shall be equal to half the difference of the $\angle^s ACB, ABC$.

As before, the $\angle ACK = \text{the sum of the } \angle^s KBC, KCB$.

To each of these add the $\angle KCB$:

then the $\angle ACB = \text{the } \angle KBC \text{ together with twice the } \angle KCB$.

$\therefore \text{twice the } \angle KCB = \text{the difference of the } \angle^s ACB, KBC$,
 that is, the $\angle KCB = \text{half the difference of the } \angle^s ACB, ABC$.

COROLLARY. If X be the middle point of the base, and XL be joined, it may be shewn by Ex. 3, p. 97, that XL is half BK ; that is, that XL is half the difference of the sides AB, AC .

8. In any triangle the angle contained by the bisector of the vertical angle and the perpendicular from the vertex to the base is equal to half the difference of the angles at the base. [See Ex. 3, p. 59.]

9. In a triangle ABC the side AC is produced to D , and the angles BAC, BCD are bisected by straight lines which meet at F ; shew that they contain an angle equal to half the angle at B .

10. If in a right-angled triangle one of the acute angles is double of the other, shew that the hypotenuse is double of the shorter side.

11. If in a diagonal of a parallelogram any two points equidistant from its extremities be joined to the opposite angles, the figure thus formed will be also a parallelogram.

12. ABC is a given equilateral triangle, and in the sides BC , CA , AB the points X , Y , Z are taken respectively, so that BX , CY and AZ are all equal. AX , BY , CZ are now drawn, intersecting in P , Q , R : shew that the triangle PQR is equilateral.

13. If in the sides AB , BC , CD , DA of a parallelogram $ABCD$ four points P , Q , R , S be taken in order, one in each side, so that AP , BQ , CR , DS are all equal; shew that the figure $PQRS$ is a parallelogram.

14. In the figure of r. 1, if the circles intersect at F , and if CA and CB are produced to meet the circles in P and Q respectively; shew that the points P , F , Q are in the same straight line; and shew also that the triangle CPQ is equilateral.

[Problems marked (*) admit of more than one solution.]

15. Through two given points draw two straight lines forming with a straight line given in position, an equilateral triangle.

*16. From a given point it is required to draw to two parallel straight lines two equal straight lines at right angles to one another.

*17. Three given straight lines meet at a point; draw another straight line so that the two portions of it intercepted between the given lines may be equal to one another.

18. From a given point draw three straight lines of given lengths, so that their extremities may be in the same straight line, and intercept equal distances on that line. [See Fig. to r. 16.]

19. Use the properties of the equilateral triangle to trisect a given finite straight line.

20. In a given triangle inscribe a rhombus, having one of its angles coinciding with an angle of the triangle.

VI. ON THE CONCURRENCE OF STRAIGHT LINES IN A TRIANGLE.

DEFINITIONS. (i) Three or more straight lines are said to be **concurrent** when they meet in one point.

(ii) Three or more points are said to be **collinear** when they lie upon one straight line.

We here give some propositions relating to the concurrence of certain groups of straight lines drawn in a triangle: the importance of these theorems will be more fully appreciated when the student is familiar with Books III. and IV.


1. *The perpendiculars drawn to the sides of a triangle from their middle points are concurrent.*

Let ABC be a Δ , and X, Y, Z the middle points of its sides:

then shall the perp^s drawn to the sides from X, Y, Z be concurrent.

From Z and Y draw perp^s to AB, AC ; these perp^s, since they cannot be parallel, will meet at point O . *Ax. 12.*

Join OX .



It is required to prove that OX is perp. to BC .

Join OA, OB, OC .

Because $\left\{ \begin{array}{l} \text{In the } \Delta^s OYA, OYC, \\ YA = YC, \\ \text{and } OY \text{ is common to both;} \\ \text{also the } \angle OYA = \text{the } \angle OYC, \text{ being rt. } \angle^s. \end{array} \right. \quad \begin{array}{l} \text{Hyp.} \\ \text{I. 4.} \end{array}$

$\therefore OA = OC$.

Similarly, from the $\Delta^s OZA, OZB$, it may be proved that $OA = OB$.

Hence OA, OB, OC are all equal.

Again, in the $\Delta^s OXB, OXC$

Because $\left\{ \begin{array}{l} BX = CX, \\ \text{and } XO \text{ is common to both;} \\ \text{also } OB = OC: \end{array} \right. \quad \begin{array}{l} \text{Hyp.} \\ \text{Proved.} \\ \text{I. 8.} \end{array}$

$\therefore \text{the } \angle OXB = \text{the } \angle OXC$;

but these are adjacent \angle^s ;

\therefore they are rt. \angle^s ;

that is, OX is perp. to BC . *Def. 7.*

Hence the three perp^s OX, OY, OZ meet in the point O .

Q. E. D.

2. *The bisectors of the angles of a triangle are concurrent.*

Let ABC be a Δ . Bisect the $\angle^s ABC, BCA$, by straight lines which must meet at some point O . *Ax. 12.*

Join AO .


It is required to prove that AO bisects the $\angle BAC$.

From O draw OP, OQ, OR perp. to the sides of the Δ .

Then in the $\Delta^s OBP, OBR$,


Because $\left\{ \begin{array}{l} \text{the } \angle OBP = \text{the } \angle OBR, \\ \text{and the } \angle OPB = \text{the } \angle ORB, \text{ being rt. } \angle^s, \\ \text{and } OB \text{ is common;} \end{array} \right. \quad \begin{array}{l} \text{Constr.} \\ \text{I. 26.} \end{array}$

$\therefore OP = OR$.



Similarly from the Δ^s OCP, OCQ,
it may be shewn that $OP = OQ$,
 $\therefore OP, OQ, OR$ are all equal.

Again in the Δ^s ORA, OQA,
the \angle^s ORA, OQA are rt. \angle^s ,
Because { and the hypotenuse OA is
common,
also $OR = OQ$; *Proved.*
 \therefore the \angle RAO = the \angle QAO.



Ex. 12, p. 91.

That is, AO is the bisector of the \angle BAC.

Hence the bisectors of the three \angle^s meet at the point O.

Q. E. D.

3. The bisectors of two exterior angles of a triangle and the bisector of the third angle are concurrent.


Let ABC be a Δ , of which the sides AB, AC are produced to any points D and E.

Bisect the \angle^s DBC, ECB by straight lines which must meet at some point O. Ax. 12.
Join AO.

It is required to prove that AO bisects the angle BAC.

From O draw OP, OQ, OR perp. to the sides of the Δ .

Then in the Δ^s OBP, OBR,
(the \angle OBP = the \angle OBR, *Constr.*
Because { also the \angle OPB = the \angle ORB,
being rt. \angle^s ,
and OB is common;
 $\therefore OP = OR$.



I. 26.

Similarly in the Δ^s OCP, OCQ,
it may be shewn that $OP = OQ$:
 $\therefore OP, OQ, OR$ are all equal.

Again in the Δ^s ORA, OQA,
the \angle^s ORA, OQA are rt. \angle^s ,
Because { and the hypotenuse OA is common,
also $OR = OQ$;
 \therefore the \angle RAO = the \angle QAO.

Proved.
Ex. 12, p. 91.

That is, AO is the bisector of the \angle BAC.
the bisectors of the two exterior \angle^s DBC, ECB,
and of the interior \angle BAC meet at the point O.

Q. E. D.

4. *The medians of a triangle are concurrent.*

Let ABC be a Δ . Let BY and CZ be two of its medians, and let them intersect at O .

Join AO ,

and produce it to meet BC in X .

It is required to shew that AX is the remaining median of the Δ .

Through C draw CK parallel to BY :


produce AX to meet CK at K .

Join BK .

In the ΔAKC ,

because Y is the middle point of AC , and YO is parallel to CK ,

$\therefore O$ is the middle point of AK .



Ex. 1, p. 96.

Again in the ΔABK ,

since Z and O are the middle points of AB , AK ,

$\therefore ZO$ is parallel to BK ,

Ex. 2, p. 96.

that is, OC is parallel to BK :

\therefore the figure $BKCO$ is a par^m .

But the diagonals of a par^m bisect one another, Ex. 5, p. 64.

$\therefore X$ is the middle point of BC .

That is, AX is a median of the Δ .

Hence the three medians meet at the point O . Q.E.D.

COROLLARY. *The three medians of a triangle cut one another at a point of trisection, the greater segment in each being towards the angular point.*

For in the above figure it has been proved that

$AO = OK$,

also that OX is half of OK ;

$\therefore OX$ is half of OA ;

that is, OX is one third of AX .

Similarly OY is one third of BY ,


and OZ is one third of CZ .

Q.E.D.

By means of this Corollary it may be shewn that in any triangle the shorter median bisects the greater side.

[The point of intersection of the three medians of a triangle is called the **centroid**. It is shewn in mechanics that a thin triangular plate will balance in any position about this point: therefore the centroid of a triangle is also its centre of gravity.]

5. The perpendiculars drawn from the vertices of a triangle to the opposite sides are concurrent.



Let ABC be a Δ , and AD , BE , CF the three perp^s drawn from the vertices to the opposite sides:

then shall these perp^s be concurrent.

Through A , B , and C draw straight lines MN , NL , LM parallel to the opposite sides of the Δ .

Then the figure $BAMC$ is a par^m.

Def. 26.

$\therefore AB = MC$.

I. 34.

Also the figure $BACL$ is a par^m.

$\therefore AB = LC$,

$\therefore LC = CM$;

that is, C is the middle point of LM .

So also A and B are the middle points of MN and NL .

Hence AD , BE , CF are the perp^s to the sides of the Δ LMN from their middle points. Ex. 3, p. 54.

But these perp^s meet in a point: Ex. 1, p. 103.

that is, the perp^s drawn from the vertices of the Δ ABC to the opposite sides meet in a point. Q.E.D.

[For another proof see Theorems and Examples on Book III.]

DEFINITIONS.

(i) The intersection of the perpendiculars drawn from the vertices of a triangle to the opposite sides is called its **orthocentre**.

(ii) The triangle formed by joining the feet of the perpendiculars is called the **pedal triangle**.


VII. ON THE CONSTRUCTION OF TRIANGLES WITH GIVEN PARTS.

No general rules can be laid down for the solution of problems in this section; but in a few typical cases we give constructions, which the student will find little difficulty in adapting to other questions of the same class.

1. Construct a right-angled triangle, having given the hypotenuse and the sum of the remaining sides.

[It is required to construct a rt. angled Δ , having its hypotenuse equal to the given straight line K , and the sum of its remaining sides equal to AB .

From A draw AE making with BA an \angle equal to half a rt. \angle . From centre B , with radius equal to K , describe a circle cutting AE in the points C, C' .




From C and C' draw perp^s $CD, C'D'$ to AB ; and join $CB, C'B$. Then either of the Δ^s $CDB, C'D'B$ will satisfy the given conditions.

NOTE. If the given hypotenuse K be greater than the perpendicular drawn from B to AE , there will be *two* solutions. If the line K be equal to this perpendicular, there will be *one* solution; but if less, the problem is *impossible*.]

2. Construct a right-angled triangle, having given the hypotenuse and the difference of the remaining sides.

3. Construct an isosceles right-angled triangle, having given the sum of the hypotenuse and one side.

4. Construct a triangle, having given the perimeter and the angles at the base.



[Let AB be the perimeter of the required Δ , and X and Y the \angle^s at the base.

From A draw AP , making the $\angle BAP$ equal to half the $\angle X$.
 From B draw BP , making the $\angle ABP$ equal to half the $\angle Y$.
 From P draw PQ , making the $\angle APQ$ equal to the $\angle BAP$.
 From P draw PR , making the $\angle BPR$ equal to the $\angle ABP$.
 Then shall PQR be the required Δ .]

5. Construct a right-angled triangle, having given the perimeter and one acute angle.

6. Construct an isosceles triangle of given altitude, so that its base may be in a given straight line, and its two equal sides may pass through two fixed points. [See Ex. 7, p. 49.]


7. Construct an equilateral triangle, having given the length of the perpendicular drawn from one of the vertices to the opposite side.

8. Construct an isosceles triangle, having given the base, and the difference of one of the remaining sides and the perpendicular drawn from the vertex to the base. [See Ex. 1, p. 88.]

9. Construct a triangle, having given the base, one of the angles at the base, and the sum of the remaining sides.

10. Construct a triangle, having given the base, one of the angles at the base, and the difference of the remaining sides.

11. Construct a triangle, having given the base, the difference of the angles at the base, and the difference of the remaining sides.



[Let AB be the given base, X the difference of the \angle 's at the base, and K the difference of the remaining sides.

Draw BE , making the $\angle ABE$ equal to half the $\angle X$.

From centre A , with radius equal to K , describe a circle cutting BE in D and D' . Let D be the point of intersection nearer to B .

Join AD and produce it to C .

Draw BC , making the $\angle DBC$ equal to the $\angle BDC$.

Then shall CAB be the Δ required. Ex. 7, p. 101.

NOTE. This problem is possible only when the given difference K is greater than the perpendicular drawn from A to BE .]

12. Construct a triangle, having given the base, the difference of the angles at the base, and the sum of the remaining sides.

13. Construct a triangle, having given the perpendicular from the vertex on the base, and the difference between each side and the adjacent segment of the base.

14. Construct a triangle, having given two sides and the median which bisects the remaining side. [See Ex. 18, p. 102.]

15. Construct a triangle, having given one side, and the medians which bisect the two remaining sides.

[See Fig. to Ex. 4, p. 105.]

Let BC be the given side. Take two-thirds of each of the given medians; hence construct the triangle BOC . The rest of the construction follows easily.]

16. Construct a triangle, having given its three medians.

[See Fig. to Ex. 4, p. 105.]

Take two-thirds of each of the given medians, and construct the triangle OKC . The rest of the construction follows easily.]

VIII. ON AREAS.

See Propositions 35—48.

It must be understood that throughout this section the word *equal* as applied to rectilinear figures will be used as denoting *equality of area* unless otherwise stated.

1. Shew that a parallelogram is bisected by any straight line which passes through the middle point of one of its diagonals. [I. 29, 26.]

2. Bisect a parallelogram by a straight line drawn through a given point.

3. Bisect a parallelogram by a straight line drawn perpendicular to one of its sides.

4. Bisect a parallelogram by a straight line drawn parallel to a given straight line.

5. $ABCD$ is a trapezium in which the side AB is parallel to DC . Shew that its area is equal to the area of a parallelogram formed by drawing through X , the middle point of BC , a straight line parallel to AD . [I. 29, 26.]

6. A trapezium is equal to a parallelogram whose base is half the sum of the parallel sides of the given figure, and whose altitude is equal to the perpendicular distance between them.

7. $ABCD$ is a trapezium in which the side AB is parallel to DC ; shew that it is double of the triangle formed by joining the extremities of AD to X , the middle point of BC .

8. Shew that a trapezium is bisected by the straight line which joins the middle points of its parallel sides. [I. 38.]

In the following group of Exercises the proofs depend chiefly on Propositions 37 and 38, and the two converse theorems.

9. If two straight lines AB , CD intersect at X , and if the straight lines AC and BD , which join their extremities are parallel, shew that the triangle AXD is equal to the triangle BXC .

10. If two straight lines AB , CD intersect at X , so that the triangle AXD is equal to the triangle XCB , then AC and BD are parallel.

11. $ABCD$ is a parallelogram, and X any point in the diagonal AC produced; shew that the triangles XBC , XDC are equal. [See Ex. 13, p. 64.]

12. ABC is a triangle, and R , Q the middle points of the sides AB , AC ; shew that if BQ and CR intersect in X , the triangle BXC is equal to the quadrilateral $AQXR$. [See Ex. 5, p. 73.]

13. If the middle points of the sides of a quadrilateral be joined in order, the *parallelogram* so formed [see Ex. 9, p. 97] is equal to half the given figure.

14. Two triangles of equal area stand on the same base but on opposite sides of it: shew that the straight line joining their vertices is bisected by the base, or by the base produced.

15. The straight line which joins the middle points of the diagonals of a trapezium is parallel to each of the two parallel sides.

16. (i) *A triangle is equal to the sum or difference of two triangles on the same base (or on equal bases), if the altitude of the former is equal to the sum or difference of the altitudes of the latter.*

(ii) *A triangle is equal to the sum or difference of two triangles of the same altitude if the base of the former is equal to the sum or difference of the bases of the latter.*

Similar statements hold good of parallelograms.

17. $ABCD$ is a parallelogram, and O is any point outside it; shew that the sum or difference of the triangles OAB , OCD is equal to half the parallelogram. Distinguish between the two cases.

On the following proposition depends an important theorem in Mechanics: we give a proof of the first case, leaving the second case to be deduced by a similar method.


18. (i) $ABCD$ is a parallelogram, and O is any point without the angle BAD and its opposite vertical angle; shew that the triangle OAC is equal to the sum of the triangles OAD , OAB .

(ii) If O is within the angle BAD or its opposite vertical angle, the triangle OAC is equal to the difference of the triangles OAD , OAB .

CASE I. If O is without the $\angle DAB$ and its opp. vert. \angle , then OA is without the par^m $ABCD$: therefore the perp. drawn from C to OA is equal to the sum of the perp^s drawn from B and D to OA . [See Ex. 20, p. 99.]

Now the Δ^s OAC , OAD , OAB are upon the same base OA ; and the altitude of the Δ OAC with respect to this base has been shewn to be equal to the sum of the altitudes of the Δ^s OAD , OAB .


Therefore the Δ OAC is equal to the sum of the Δ^s OAD , OAB . [See Ex. 16, p. 110.]



19. $ABCD$ is a parallelogram, and, through O , any point within it, straight lines are drawn parallel to the sides of the parallelogram; shew that the difference of the parallelograms DO , BO is double of the triangle AOC . [See preceding theorem (ii).]

20. The area of a quadrilateral is equal to the area of a triangle, having two of its sides equal to the diagonals of the given figure, and the included angle equal to either of the angles between the diagonals.

21. ABC is a triangle, and D is any point in AB : it is required to draw through D a straight line DE to meet BC produced in E , so that the triangle DBE may be equal to the triangle ABC .



[Join DC . Through A draw AE parallel to DC . I. 31.
Join DE .

The Δ EBD shall be equal to the Δ ABC . I. 37.]

22. On a base of given length describe a triangle equal to a given triangle and having an angle equal to an angle of the given triangle.

23. Construct a triangle equal in area to a given triangle, and having a given altitude.

24. On a base of given length construct a triangle equal to a given triangle, and having its vertex on a given straight line.

25. On a base of given length describe (i) an isosceles triangle; (ii) a right-angled triangle, equal to a given triangle.

26. Construct a triangle equal to the sum or difference of two given triangles. [See Ex. 16, p. 110.]

27. ABC is a given triangle, and X a given point: describe a triangle equal to ABC , having its vertex at X , and its base in the same straight line as BC .

28. $ABCD$ is a quadrilateral: on the base AB construct a triangle equal in area to $ABCD$, and having the angle at A common with the quadrilateral.

[Join BD . Through C draw CX parallel to BD , meeting AD produced in X ; join BX .]

29. Construct a rectilineal figure equal to a given rectilineal figure, and having fewer sides by one than the given figure.

Hence shew how to construct a triangle equal to a given rectilineal figure.

30. $ABCD$ is a quadrilateral: it is required to construct a triangle equal in area to $ABCD$, having its vertex at a given point X in DC , and its base in the same straight line as AB .

31. Construct a rhombus equal to a given parallelogram.

32. Construct a parallelogram which shall have the same area and perimeter as a given triangle.

33. Bisect a triangle by a straight line drawn through one of its angular points.

34. Trisect a triangle by straight lines drawn through one of its angular points. [See Ex. 19, p. 102, and i. 38.]

35. Divide a triangle into any number of equal parts by straight lines drawn through one of its angular points. [See Ex. 19, p. 99, and i. 38.]

36. *Bisect a triangle by a straight line drawn through a given point in one of its sides.*


[Let ABC be the given Δ , and P the given point in the side AB .

Bisect AB at Z ; and join CZ , CP .
Through Z draw ZQ parallel to CP .

Join PQ .

Then shall PQ bisect the Δ .

See Ex. 21, p. 111.]



37. *Trisect a triangle by straight lines drawn from a given point in one of its sides.*


[Let ABC be the given Δ , and X the given point in the side BC .

Trisect BC at the points P , Q . Ex. 19, p. 99.
Join AX , and through P and Q draw PH and QK parallel to AX .

Join XH , XK .

These straight lines shall trisect the Δ ; as may be shewn by joining AP , AQ .

See Ex. 21, p. 111.]



38. *Cut off from a given triangle a fourth, fifth, sixth, or any part required by a straight line drawn from a given point in one of its sides.*
[See Ex. 19, p. 99, and Ex. 21, p. 111.]


39. *Bisect a quadrilateral by a straight line drawn through an angular point.*

[Two constructions may be given for this problem: the first will be suggested by Exercises 28 and 33, p. 112.

The second method proceeds thus.

Let $ABCD$ be the given quadrilateral, and A the given angular point.

Join AC , BD , and bisect BD in X .
Through X draw PXQ parallel to AC , meeting BC in P ; join AP .
Then shall AP bisect the quadrilateral.
Join AX , CX , and use I. 37, 38.]



40. *Cut off from a given quadrilateral a third, a fourth, a fifth, or any part required, by a straight line drawn through a given angular point.*
[See Exercises 28 and 35, p. 112.]

[The following Theorems depend on 1. 47.]

41. In the figure of 1. 47, shew that

- (i) the sum of the squares on AB and AE is equal to the sum of the squares on AC and AD .
- (ii) the square on EK is equal to the square on AB with four times the square on AC .
- (iii) the sum of the squares on EK and FD is equal to five times the square on BC .

42. If a straight line be divided into any two parts the square on the straight line is greater than the squares on the two parts.

43. If the square on one side of a triangle is less than the squares on the remaining sides, the angle contained by these sides is acute; if greater, obtuse.

44. ABC is a triangle, right-angled at A ; the sides AB , AC are intersected by a straight line PQ , and BQ , PC are joined: shew that the sum of the squares on BQ , PC is equal to the sum of the squares on BC , PQ .

45. In a right-angled triangle four times the sum of the squares on the medians which bisect the sides containing the right angle is equal to five times the square on the hypotenuse.

46. Describe a square whose area shall be three times that of a given square.

47. Divide a straight line into two parts such that the sum of their squares shall be equal to a given square.

IX. ON LOCUS.

It is frequently required in the course of Plane Geometry to find the position of a point which satisfies given conditions. Now all problems of this type hitherto considered have been found to be capable of definite determination, though some admit of more than one solution: this however will not be the case if *only one* condition is given. For example, if we are asked to find a point which shall be at a given distance from a given point, we observe at once that the problem is *indeterminate*, that is, that it admits of an indefinite number of solutions; for the condition stated is satisfied by any point on the circumference of the circle described from the given point as centre, with a radius equal to the given distance: moreover this condition is satisfied by no other point within or without the circle.

Again, suppose that it is required to find a point at a given distance from a given straight line.

Here, too, it is obvious that there are an infinite number of such points, and that they lie on the two parallel straight lines which may be drawn on either side of the given straight line at the given distance from it: further, no point that is not on one or other of these parallels satisfies the given condition.

Hence we see that when one condition is assigned it is not sufficient to determine the position of a point absolutely, but it may have the effect of restricting it to some definite line or lines, straight or curved. This leads us to the following definition.

DEFINITION. The **Locus** of a point satisfying an assigned condition consists of the line, lines, or part of a line, to which the point is thereby restricted; provided that the condition is satisfied by every point on such line or lines, and by no other.

A locus is sometimes defined as the path traced out by a point which moves in accordance with an assigned law.

Thus the locus of a point, which is always at a given distance from a given point, is a circle of which the given point is the centre: and the locus of a point, which is always at a given distance from a given straight line, is a pair of parallel straight lines.

We now see that in order to infer that a certain line, or system of lines, is the locus of a point under a given condition, it is necessary to prove

- (i) that any point which fulfils the given condition is on the supposed locus;
- (ii) that every point on the supposed locus satisfies the given condition.

1. Find the locus of point which is always equidistant from two given points.

Let A, B be the two given points.

(a) Let P be any point equidistant from A and B, so that $AP = BP$.

Bisect AB at X, and join PX.

Then in the Δ^s AXP, BXP,

Because $\begin{cases} AX = BX, \\ \text{and } PX \text{ is common to both,} \\ \text{also } AP = BP, \end{cases}$

Constr. 

\therefore the \angle PXA = the \angle PXB;

Hyp.

$\simeq 8.$

and they are adjacent \angle^s ;

\therefore PX is perp. to AB.

Q

\therefore Any point which is equidistant from A and B is on the straight line which bisects AB at right angles.

(β) Also every point in this line is equidistant from A and B.

For let Q be any point in this line.

Join AQ, BQ.

Then in the Δ^s AXQ, BXQ,

AX = BX,

Because { and XQ is common to both;
also the \angle AXQ = the \angle BXQ, being rt. \angle^s ,


\therefore AQ = BQ.

I. 4.

That is, Q is equidistant from A and B.

Hence we conclude that the locus of the point equidistant from two given points A, B is the straight line which bisects AB at right angles.

2. To find the locus of the middle point of a straight line drawn from a given point to meet a given straight line of unlimited length.



Let A be the given point, and BC the given straight line of unlimited length.

(α) Let AX be any straight line drawn through A to meet BC, and let P be its middle point.

Draw AF perp. to BC, and bisect AF at E.

Join EF, and produce it indefinitely.

Since AFX is a Δ , and E, P the middle points of the two sides AF, AX,

\therefore EP is parallel to the remaining side FX. Ex. 2, p. 96.

\therefore P is on the straight line which passes through the fixed point E, and is parallel to BC.

(β) Again, every point in EP, or EP produced, fulfils the required condition.

For, in this straight line take any point Q.

Join AQ, and produce it to meet BC in Y.

Then FAY is a Δ , and through E, the middle point of the side AF, EQ is drawn parallel to the side FY,

\therefore Q is the middle point of AY.

Ex. 1, p. 96.

Hence the required locus is the straight line drawn parallel to BC, and passing through E, the middle point of the perp. from A to BC.

3. Find the locus of a point equidistant from two given intersecting straight lines. [See Ex. 3, p. 49.]

4. Find the locus of a point at a given radial distance from the circumference of a given circle.

5. Find the locus of a point which moves so that the sum of its distances from two given intersecting straight lines of unlimited length is constant.

6. Find the locus of a point when the differences of its distances from two given intersecting straight lines of unlimited length is constant.

7. A straight rod of given length slides between two straight rulers placed at right angles to one another: find the locus of its middle point. [See Ex. 2, p. 100.]

8. On a given base as hypotenuse right-angled triangles are described: find the locus of their vertices.

9. AB is a given straight line, and AX is the perpendicular drawn from A to any straight line passing through B : find the locus of the middle point of AX .

10. Find the locus of the vertex of a triangle, when the base and area are given.

11. Find the locus of the intersection of the diagonals of a parallelogram, of which the base and area are given.

12. Find the locus of the intersection of the medians of a triangle described on a given base and of given area.

X. ON THE INTERSECTION OF LOCI.

It appears from various problems which have already been considered, that we are often required to find a point, the position of which is subject to two given conditions. The method of loci is very useful in the solution of problems of this kind: for corresponding to each condition there will be a locus on which the required point must lie; hence all points which are common to these two loci, that is, all the points of intersection of the loci, will satisfy *both* the given conditions.

EXAMPLE 1. *To construct a triangle, having given the base, the altitude, and the length of the median which bisects the base.*

Let **AB** be the given base, and **P** and **Q** the lengths of the altitude and median respectively:

then the triangle is known if its *vertex* is known.

(i) Draw a straight line **CD** parallel to **AB**, and at a distance from it equal to **P**.

then the required vertex must lie on **CD**.

(ii) Again, from the middle point of **AB** as centre, with radius equal to **Q**, describe a circle:

then the required vertex must lie on this circle.

Hence any points which are common to **CD** and the circle, satisfy both the given conditions: that is to say, if **CD** intersect the circle in **E**, **F** each of the points of intersection might be the vertex of the required triangle. This supposes the length of the median **Q** to be greater than the altitude.

EXAMPLE 2. *To find a point equidistant from three given points **A**, **B**, **C**, which are not in the same straight line.*

(i) The locus of points equidistant from **A** and **B** is the straight line **PQ**, which bisects **AB** at right angles. Ex. 1, p. 115.

(ii) Similarly the locus of points equidistant from **B** and **C** is the straight line **RS** which bisects **BC** at right angles.

Hence the point common to **PQ** and **RS** must satisfy both conditions: that is to say, the point of intersection of **PQ** and **RS** will be equidistant from **A**, **B**, and **C**.

These principles may also be used to prove the theorems relating to concurrency already given on page 103.

EXAMPLE. *To prove that the bisectors of the angles of a triangle are concurrent.*

Let **ABC** be a triangle.
Bisect the \angle^s **ABC**, **BCA** by straight lines **BO**, **CO**: these must meet at some point **O**. Ex. 12.

Join **OA**.

Then shall **OA** bisect the \angle **BAC**.
Now **BO** is the locus of points equidistant from **BC**, **BA**; Ex. 3, p. 19.


\therefore **OP** = **OR**.

Similarly **CO** is the locus of points equidistant from **BC**, **CA**.


\therefore **OP** \perp **OQ**; hence **OR** = **OQ**.

\therefore **O** is on the locus of points equidistant from **AB** and **AC**:
that is **OA** is the bisector of the \angle **BAC**.

Hence the bisectors of the three \angle^s meet at the point **O**.



PROPOSITION 30. PROBLEM.

To bisect a given arc.

Let ADB be the given arc
it is required to bisect it.

Join AB ; and bisect it at C . I. 10.

At C draw CD at rt. angles to AB , meeting the given arc at D . I. 11.

Then shall the arc ADB be bisected at D ,

Join AD , BD .

Then in the $\triangle ACD$, BCD ,

Because $\left\{ \begin{array}{l} AC = BC, \\ \text{and } CD \text{ is common;} \\ \text{and the } \angle ACD = \text{the } \angle BCD, \text{ being rt. angles;} \end{array} \right.$ Constr.

$\therefore AD = BD$. I. 4.

And since in the $\triangle ADB$, the chords AD , BD are equal,
 \therefore the arcs cut off by them are equal, the minor arc equal
to the minor, and the major arc to the major: III. 28.

and the arcs AD , BD are both minor arcs,
for each is less than a semi-circumference, since DC , bisecting
the chord AB at rt. angles, must pass through the centre
of the circle. III. 4 Cor.

\therefore the arc $AD =$ the arc BD :
that is, the arc ADB is bisected at D . Q. E. F.

EXERCISES.

1. If a tangent to a circle is parallel to a chord, the point of contact will bisect the arc cut off by the chord.

2. Trisect a quadrant, or the fourth part of the circumference, of a circle.

BOOK II.

Book II. deals with the areas of rectangles and squares.

DEFINITIONS.

1. A **Rectangle** is a parallelogram which has one of its angles a right angle.

It should be remembered that if a parallelogram has *one* right angle, *all* its angles are right angles. [Ex. 1, p. 64.]

2. A rectangle is said to be **contained** by any two of its sides which form a right angle: for it is clear that both the form and magnitude of a rectangle are fully determined when the lengths of two such sides are given.

Thus the rectangle ACDB is said to be contained by AB, AC; or by CD, DB: and if X and Y are two straight lines equal respectively to AB and AC, then the rectangle contained by X and Y is equal to the rectangle contained by AB, AC.

[See Ex. 12, p. 61.]




After Proposition 3, we shall use the abbreviation *rect.* AB, AC to denote the *rectangle contained by AB and AC*.

3. In any parallelogram the figure formed by either of the parallelograms about a diagonal together with the two complements is called a **gnomon**.

Thus the shaded portion of the annexed figure, consisting of the parallelogram EH together with the complements AK, KC is the *gnomon* AHF.

The other gnomon in the figure is that which is made up of AK, GF and FH, namely the *gnomon* AFH.



INTRODUCTORY.


Pure Geometry makes no use of *number* to estimate the magnitude of the lines, angles, and figures with which it deals: hence it requires no *units of magnitude* such as the student is familiar with in Arithmetic.

For example, though Geometry is concerned with the relative lengths of straight lines, it does not seek to express those lengths in terms of *yards, feet, or inches*: similarly it does not ask how many *square yards* or *square feet* a given figure contains, nor how many *degrees* there are in a given angle.

This constitutes an essential difference between the method of Pure Geometry and that of Arithmetic and Algebra; at the same time a close connection exists between the results of these two methods.

In the case of Euclid's Book II., this connection rests upon the fact that *the number of units of area in a rectangular figure is found by multiplying together the numbers of units of length in two adjacent sides*.

For example, if the two sides AB, AD of the rectangle ABCD are respectively *four* and *three* inches long, and if through the points of division parallels are drawn as in the annexed figure, it is seen that the rectangle is divided into *three* rows, each containing *four* square inches, or into *four* columns, each containing *three* square inches.



Hence the whole rectangle contains 3×4 , or 12, square inches.

Similarly if AB and AD contain m and n units of length respectively, it follows that the rectangle ABCD will contain mn units of area: further, if AB and AD are equal, each containing m units of length, the rectangle becomes a square, and contains m^2 units of area.

[It must be understood that this explanation implies that the lengths of the straight lines AB, AD are **commensurable**, that is, that they can be expressed *exactly* in terms of some common unit.

This however is not always the case: for example, it may be proved that the side and diagonal of a square are so related, that it is impossible to divide either of them into equal parts, *of which the other contains an exact number*. Such lines are said to be **incommen-**

surable. Hence if the adjacent sides of a rectangle are incommensurable, we cannot choose any linear unit in terms of which these sides may be *exactly* expressed; and thus it will be impossible to subdivide the rectangle into squares of unit area, as illustrated in the figure of the preceding page. We do not here propose to enter further into the subject of incommensurable quantities: it is sufficient to point out that further knowledge of them will convince the student that the area of a rectangle may be expressed *to any required degree of accuracy* by the product of the lengths of two adjacent sides, whether those lengths are commensurable or not.]

From the foregoing explanation we conclude that *the rectangle contained by two straight lines* in Geometry corresponds to *the product of two numbers* in Arithmetic or Algebra; and that *the square described on a straight line* corresponds to *the square of a number*. Accordingly it will be found in the course of Book II. that several theorems relating to the areas of rectangles and squares are analogous to well-known algebraical formulæ.


In view of these principles the rectangle contained by two straight lines AB , BC is sometimes expressed in the form of a product, as $AB \cdot BC$, and the square described on AB as AB^2 . This notation, together with the signs $+$ and $-$, will be employed in the additional matter appended to this book; *but it is not admitted into Euclid's text* because it is desirable in the first instance to emphasize the distinction between geometrical magnitudes themselves and the numerical equivalents by which they may be expressed arithmetically.

PROPOSITION I. THEOREM.

If there are two straight lines, one of which is divided into any number of parts, the rectangle contained by the two straight lines is equal to the sum of the rectangles contained by the undivided straight line and the several parts of the divided line.

Let P and AB be two straight lines, and let AB be divided into any number of parts AC , CD , DB :

then shall the rectangle contained by P , AB be equal to the sum of the rectangles contained by P , AC , by P , CD , and by P , DB .



From A draw AF perp. to AB; I. 11.
 and make AG equal to P. I. 3.
 Through G draw GH par^l to AB; I. 31.
 and through C, D, B draw CK, DL, BH par^l to AG.

Now the fig. AH is made up of the figs. AK, CL, DH :
 and of these,

the fig. AH is the rectangle contained by P, AB;
 for the fig. AH is contained by AG, AB; and AG = P :
 and the fig. AK is the rectangle contained by P, AC;
 for the fig. AK is contained by AG, AC; and AG = P :
 also the fig. CL is the rectangle contained by P, CD;
 for the fig. CL is contained by CK, CD;

and CK = the opp. side AG, and AG = P: I. 34.
 similarly the fig. DH is the rectangle contained by P, DB.

∴ the rectangle contained by P, AB is equal to the
 sum of the rectangles contained by P, AC, by P, CD, and
 by P, DB. Q.E.D.

CORRESPONDING ALGEBRAICAL FORMULA.

In accordance with the principles explained on page 122, the result
 of this proposition may be written thus:

$$P \cdot AB = P \cdot AC + P \cdot CD + P \cdot DB.$$

Now if the line P contains p units of length, and if AC, CD, DB
 contain a , b , c units respectively,

$$\text{then } AB = a + b + c,$$

and we have
$$p(a + b + c) = pa + pb + pc.$$

PROPOSITION 2. THEOREM.

If a straight line is divided into any two parts, the square on the whole line is equal to the sum of the rectangles contained by the whole line and each of the parts.

Let the straight line AB be divided at C into the two parts AC , CB :

then shall the sq. on AB be equal to the sum of the rectx. contained by AB , AC , and by AB , BC .

On AB describe the square $ADEB$. I. 46.

Through C draw CF par^l to AD . I. 31.

Now the fig. AE is made up of the figs. AF , CE :
and of these,

the fig. AE is the sq. on AB : *Constr.*
and the fig. AF is the rectangle contained by AB , AC ;
for the fig. AF is contained by AD , AC ; and $AD = AB$;
also the fig. CE is the rectangle contained by AB , BC ;
for the fig. CE is contained by BE , BC ; and $BE = AB$.

\therefore the sq. on AB – the sum of the rectx. contained by AB , AC , and by AB , BC . Q.E.D.

CORRESPONDING ALGEBRAICAL FORMULA.

The result of this proposition may be written

$$AB^2 = AB \cdot AC + AB \cdot BC.$$


Let AC contain a units of length, and let CB contain b units,

$$\text{then } AB = a + b,$$

and we have $(a + b)^2 = (a + b) a + (a + b) b.$

PROPOSITION 3. THEOREM.

If a straight line is divided into any two parts, the rectangle contained by the whole and one of the parts is equal to the square on that part together with the rectangle contained by the two parts.



Let the straight line AB be divided at C into the two parts AC , CB :

then shall the rect. contained by AB , AC be equal to the sq. on AC together with the rect. contained by AC , CB .

On AC describe the square $AFDC$; I. 46.
and through B draw BE par^l to AF , meeting FD produced in E .
I. 31.

Now the fig. AE is made up of the figs. AD , CE ;
and of these,

the fig. AE = the rect. contained by AB , AC ;
for $AF = AC$;

and the fig. AD is the sq. on AC ; Constr.
also the fig. CE is the rect. contained by AC , CB ;
for $CD = AC$.

\therefore the rect. contained by AB , AC is equal to the sq. on AC together with the rect. contained by AC , CB . Q.E.D.

CORRESPONDING ALGEBRAICAL FORMULA.

This result may be written $AB \cdot AC = AC^2 + AC \cdot CB$.

Let AC , CB contain a and b units of length respectively,

then $AB = a + b$,


and we have

$$(a + b)a = a^2 + ab.$$

NOTE. It should be observed that Props. 2 and 3 are *special cases* of Prop. 1.

PROPOSITION 4. THEOREM.

If a straight line is divided into any two parts, the square on the whole line is equal to the sum of the squares on the two parts together with twice the rectangle contained by the two parts.



Let the straight line AB be divided at C into the two parts AC , CB :

then shall the sq. on AB be equal to the sum of the sqq. on AC , CB , together with twice the rect. AC , CB .

On AB describe the square $ADEB$; I. 46.
and join BD .

Through C draw CF par^l to BE , meeting BD in G . I. 31.

Through G draw HGK par^l to AB .

It is first required to shew that the fig. CK is the sq. on BC .

Because the straight line BGD meets the par^{ls} CG , AD ,

\therefore the ext. angle $CGB =$ the int. opp. angle ADB . I. 29.

But $AB = AD$, being sides of a square;

\therefore the angle $ADB =$ the angle ABD ; I. 5.

\therefore the angle $CGB =$ the angle CBG .

$\therefore CB = CG$. I. 6.

And the opp. sides of the par^m CK are equal; I. 34.

\therefore the fig. CK is equilateral;

and the angle CBK is a right angle; Def. 28.

$\therefore CK$ is a square, and it is described on BC . I. 46, Cor.

Similarly the fig. HF is the sq. on HG , that is, the sq. on AC ,

for $HG =$ the opp. side AC . I. 34.

Again, the complement AG = the complement GE. r. 43.

But the fig. AG = the rect. AC, CB ; for CG = CB.

∴ the two figs. AG, GE = twice the rect. AC, CB.

*Now the sq. on AB = the fig. AE

= the figs. HF, CK, AG, GE

= the sqq. on AC, CB together with
twice the rect. AC, CB.

∴ the sq. on AB = the sum of the sqq. on AC, CB with
twice the rect. AC, CB. Q.E.D.

* For the purpose of oral work, this step of the proof may conveniently be arranged as follows :

Now the sq. on AB is equal to the fig. AE,

that is, to the figs. HF, CK, AG, GE ;

that is, to the sqq. on AC, CB together
with twice the rect. AC, CB.

COROLLARY. *Parallelograms about the diagonals of a square are themselves squares.*

CORRESPONDING ALGEBRAICAL FORMULA.

The result of this important Proposition may be written thus :

$$AB^2 = AC^2 + CB^2 + 2AC \cdot CB.$$

Let $AC = a$, and $CB = b$;

then $AB = a + b$,

and we have $(a + b)^2 = a^2 + b^2 + 2ab$.

PROPOSITION 5. THEOREM.

If a straight line is divided equally and also unequally, the rectangle contained by the unequal parts, and the square on the line between the points of section, are together equal to the square on half the line.

A P Q B

Let the straight line AB be divided equally at P , and unequally at Q :

then the rect. AQ , QB and the sq. on PQ shall be together equal to the sq. on PB .

On PB describe the square $PCDB$. I. 46.

Join BC .

Through Q draw QE par^l to BD , cutting BC in F . I. 31.

Through F draw $LFHG$ par^l to AB .

Through A draw AG par^l to BD .

Now the complement $PF =$ the complement FD : I. 43.

to each add the fig. QL ;

then the fig. $PL =$ the fig. QD .

But the fig. $PL =$ the fig. AH , for they are par^{ms} on equal bases and between the same par^{ls}. I. 36.

\therefore the fig. $AH =$ the fig. QD .

To each add the fig. PF ;

then the fig. $AF =$ the gnomon PLE .

Now the fig. $AF =$ the rect. AQ , QB , for $QB = QF$;

\therefore the rect. AQ , $QB =$ the gnomon PLE .

To each add the sq. on PQ , that is, the fig. HE ; II. 4.
then the rect. AQ , QB with the sq. on PQ


the gnomon PLE with the fig. HE
the whole fig. PD ,
which is the sq. on PB .

That is, the rect. **AQ**, **QB** and the sq. on **PQ** are together equal to the sq. on **PB**. Q.E.D.

COROLLARY. From this Proposition it follows that *the difference of the squares on two straight lines is equal to the rectangle contained by their sum and difference.*

For let **X** and **Y** be the given st. lines, of which **X** is the greater.

Draw **AP** equal to **X**, and produce it to **B**, making **PB** equal to **AP**, that is to **X**.



From **PB** cut off **PQ** equal to **Y**.

Then **AQ** is equal to the sum of **X** and **Y**,
and **QB** is equal to the difference of **X** and **Y**.

Now because **AB** is divided equally at **P** and unequally at **Q**,
 \therefore the rect. **AQ**, **QB** with sq. on **PQ** = the sq. on **PB**; II. 5.
that is, the difference of the sqq. on **PB**, **PQ** = the rect. **AQ**, **QB**,
or, the difference of the sqq. on **X** and **Y** = the rect. contained by the sum and the difference of **X** and **Y**.

CORRESPONDING ALGEBRAICAL FORMULA.

This result may be written ,

$$\mathbf{AQ \cdot QB + PQ^2 = PB^2.}$$

Let $\mathbf{AB} = 2a$; and let $\mathbf{PQ} = b$;

then **AP** and **PB** each = a .

Also $\mathbf{AQ} = a + b$; and $\mathbf{QB} = a - b$.

Hence we have

$$(a + b) (a - b) + b^2 = a^2,$$

or

$$(a + b) (a - b) = a^2 - b^2.$$

EXERCISE.

*In the above figure shew that **AP** is half the sum of **AQ** and **QB** ; and that **PQ** is half their difference.*

PROPOSITION 6. THEOREM.

If a straight line is bisected and produced to any point, the rectangle contained by the whole line thus produced, and the part of it produced, together with the square on half the line bisected, is equal to the square on the straight line made up of the half and the part produced.

A P B Q

Let the straight line AB be bisected at P, and produced to Q :

then the rect. AQ, QB and the sq. on PB shall be together equal to the sq. on PQ.

On PQ describe the square PCDQ. I. 46.

Join QC.

Through B draw BE par^l to QD, meeting QC in F. 31.

Through F draw LFHG par^l to AQ.

Through A draw AG par^l to QD:

Now the complement PF = the complement FD. I. 43.

But the fig. PF = the fig. AH ; for they are par^{ms} on equal bases and between the same par^{ls}. I. 36.

∴ the fig. AH = the fig. FD.

To each add the fig. PL ;

then the fig. AL = the gnomon PLE.

Now the fig. AL = the rect. AQ, QB, for QB = QL ;

∴ the rect. AQ, QB = the gnomon PLE.

To each add the sq. on PB, that is, the fig. HE ;
then the rect. AQ, QB with the sq. on PB

= the gnomon PLE with the fig. HE

= the whole fig. PD,

which is the square on PQ.

That is, the rect. AQ, QB and the sq. on PB are together equal to the sq. on PQ. Q.E.D.

CORRESPONDING ALGEBRAICAL FORMULA.

This result may be written

$$AQ \cdot QB + PB^2 = PQ^2.$$

Let $AB = 2a$; and let $PQ = b$;

then AP and PB each $= a$.

Also $AQ = a + b$; and $QB = b - a$.

Hence we have

$$(a + b)(b - a) + a^2 = b^2,$$

or

$$(b + a)(b - a) = b^2 - a^2.$$

DEFINITION. If a point X is taken in a straight line AB , or in AB produced, the distances of the point of section from the extremities of AB are said to be the **segments** into which AB is divided at X .



In the former case AB is divided **internally**, in the latter case **externally**.

Thus in the annexed figures the segments into which AB is divided at X are the lines XA and XB .

This definition enables us to include Props. 5 and 6 in a single Enunciation.

If a straight line is bisected, and also divided (internally or externally) into two unequal segments, the rectangle contained by the unequal segments is equal to the difference of the squares on half the line, and on the line between the points of section.

EXERCISE.

Shew that the Enunciations of Props. 5 and 6 may take the following form:

The rectangle contained by two straight lines is equal to the difference of the squares on half their sum and on half their difference.

[See Ex., p. 129.]

PROPOSITION 7. THEOREM.

If a straight line is divided into any two parts, the sum of the squares on the whole line and on one of the parts is equal to twice the rectangle contained by the whole and that part, together with the square on the other part.

H

Let the straight line AB be divided at C into the two parts AC , CB :

then shall the sum of the sqq. on AB , BC be equal to twice the rect. AB , BC together with the sq. on AC .

On AB describe the square $AD\Xi B$. i. 46.

Join BD .

Through C draw CF par^l to BE , meeting BD in G . i. 31.

Through G draw HGK par^l to AB .

Now the complement AG = the complement GE ; i. 43.
to each add the fig. CK :

then the fig. AK = the fig. CE .

But the fig. AK = the rect. AB , BC ; for BK = BC .

\therefore the two figs. AK , CE = twice the rect. AB , BC .

But the two figs. AK , CE make up the gnomon AKF and the fig. CK :

\therefore the gnomon AKF with the fig. CK = twice the rect. AB , BC .

To each add the fig. HF , which is the sq. on AC :

then the gnomon AKF with the figs. CK , HF

= twice the rect. AB , BC with the sq. on AC .

Now the sqq. on AE , BC = the figs. AE , CK

– the gnomon AKF with the
figs. CK , HF

– twice the rect. AB , BC with
the sq. on AC .

CORRESPONDING ALGEBRAICAL FORMULA.

The result of this proposition may be written

$$AB^2 + BC^2 = 2AB \cdot BC + AC^2.$$

Let $AB = a$, and $BC = b$; then $AC = a + b$.

Hence we have $a^2 + b^2 = 2ab + (a + b)^2$,

or $(a - b)^2 = a^2 - 2ab + b^2$.


PROPOSITION 8. THEOREM.

If a straight line be divided into any two parts, four times the rectangle contained by the whole line and one of the parts, together with the square on the other part, is equal to the square on the straight line which is made up of the whole and that part.

[As this proposition is of little importance we merely give the figure, and the leading points in Euclid's proof.]

Let AB be divided at C .

Produce AB to D , making BD equal to BC .



On AD describe the square $AEDF$; and complete the construction as indicated in the figure.

Euclid then proves (i) that the figs. CK , BN , GR , KO are all equal.


(ii) that the figs. AG , MP , PL , RF are all equal.

Hence the eight figures named above are four times the sum of the figs. AG , CK ; that is, four times the fig. AK ; that is, four times the rect. AB , BC .

But the whole fig. AF is made up of these eight figures, together with the fig. XH , which is the sq. on AC :

hence the sq. on AD = four times the rect. AB , BC , together with the sq. on AC .


Q.E.D.



The accompanying figure will suggest a less cumbrous proof, which we leave as an Exercise to the student.

PROPOSITION 9. THEOREM.

If a straight line is divided equally and also unequally, the sum of the squares on the two unequal parts is twice the sum of the squares on half the line and on the line between the points of section.



Let the straight line AB be divided equally at P , and unequally at Q :

then shall the sum of the sqq. on AQ , QB be twice the sum of the sqq. on AP , PQ .

At P draw PC at rt. angles to AB ; 1. 11.

and make PC equal to AP or PB . 1. 3.

Join AC , BC .

Through Q draw QD par^l to PC ; 1. 31.

and through D draw DE par^l to AB .

Join AD .

Then since $PA = PC$, Constr.

\therefore the angle $PAC =$ the angle PCA . 1. 5.

And since, in the triangle APC , the angle APC is a rt. angle, Constr.

\therefore the sum of the angles PAC , PCA is a rt. angle : 1. 32.

hence each of the angles PAC , PCA is half a rt. angle.

So also, each of the angles PBC , PCB is half a rt. angle.

\therefore the whole angle ACB is a rt. angle.

Again, the ext. angle CED – the int. opp. angle CPB , 1. 29.

\therefore the angle CED is a rt. angle :

and the angle ECD is half a rt. angle. Proved.

\therefore also the angle EDC is half a rt. angle ; 1. 32.

\therefore the angle $ECD =$ the angle EDC ;

$\therefore EC = ED$. 1. 6.

Again, the ext. angle $\angle DQB =$ the int. opp. angle $\angle CPB$. I. 29.

\therefore the angle $\angle DQB$ is a rt. angle.

And the angle $\angle QBD$ is half a rt. angle; *Proved.*

\therefore also the angle $\angle QDB$ is half a rt. angle: I. 32.

\therefore the angle $\angle QBD =$ the angle $\angle QDB$;

$\therefore QD = QB$. I. 6.

Now the sq. on $AP =$ the sq. on PC ; for $AP = PC$. *Constr.*

But the sq. on $AC =$ the sum of the sqq. on AP, PC ,
for the angle $\angle APC$ is a rt. angle. I. 47.

\therefore the sq. on AC is twice the sq. on AP .

So also, the sq. on CD is twice the sq. on ED , that is, twice the sq. on the opp. side PQ . I. 34.

Now the sqq. on $AQ, QB =$ the sqq. on AQ, QD
 $=$ the sq. on AD , for $\angle AQD$ is a rt. angle; I. 47.
 $=$ the sum of the sqq. on AC, CD ,
for $\angle ACD$ is a rt. angle; I. 47.
 $=$ twice the sq. on AP with twice the sq. on PQ . *Proved.*

That is,
the sum of the sqq. on $AQ, QB =$ twice the sum of the sqq. on AP, PQ . Q.E.D.

CORRESPONDING ALGEBRAICAL FORMULA.

The result of this proposition may be written

$$AQ^2 + QB^2 = 2(AP^2 + PQ^2).$$

Let $AB = 2a$; and $PQ = b$;

then AP and PB each $= a$.


Also $AQ = a + b$; and $QB = a - b$.

Hence we have

$$(a + b)^2 + (a - b)^2 = 2(a^2 + b^2).$$

* PROPOSITION 10. THEOREM.

If a straight line is bisected and produced to any point, the sum of the squares on the whole line thus produced, and on the part produced, is twice the sum of the squares on half the line bisected and on the line made up of the half and the part produced.



Let the st. line AB be bisected at P , and produced to Q :
then shall the sum of the sqq. on AQ , QB be twice the
sum of the sqq. on AP , PQ .

At P draw PC at right angles to AB ; 1. 11.

and make PC equal to PA or PB . 1. 3.

Join AC , BC .

Through Q draw QD par^l to PC , to meet CB produced
in D ; 1. 31.

and through D draw DE par^l to AB , to meet CP produced
in E .

Join AD .

Then since $PA = PC$, (Constr.

\therefore the angle $PAC =$ the angle PCA . 1. 5.

And since in the triangle APC , the angle APC is a rt. angle,

\therefore the sum of the angles PAC , PCA is a rt. angle. 1. 32.

Hence each of the angles PAC , PCA is half a rt. angle.

So also, each of the angles PBC , PCB is half a rt. angle.

\therefore the whole angle ACB is a rt. angle.

Again, the ext. angle $CPB =$ the int. opp. angle CED : 1. 29.

\therefore the angle CED is a rt. angle :

and the angle ECD is half a rt. angle. *Proved.*

\therefore the angle EDC is half a rt. angle. 1. 32.

\therefore the angle $ECD =$ the angle EDC ;

$\therefore EC = ED$. 1. 6.

Again, the angle \mathbf{DQB} = the alt. angle \mathbf{CPB} . 1. 29.

\therefore the angle \mathbf{DQB} is a rt. angle.

Also the angle \mathbf{QBD} = the vert. opp. angle \mathbf{CBP} ; 1. 15.

that is, the angle \mathbf{QBD} is half a rt. angle.

\therefore the angle \mathbf{QDB} is half a rt. angle; 1. 32.

\therefore the angle \mathbf{QBD} = the angle \mathbf{QDB} ;

$\therefore \mathbf{QB} = \mathbf{QD}$. 4. 6.

Now the sq. on \mathbf{AP} = the sq. on \mathbf{PC} ; for $\mathbf{AP} = \mathbf{PC}$. *Constr.*

But the sq. on \mathbf{AC} = the sum of the sqq. on \mathbf{AP} , \mathbf{PC} ,
for the angle \mathbf{APC} is a rt. angle. 1. 47.

\therefore the sq. on \mathbf{AC} is twice the sq. on \mathbf{AP} .

So also, the sq. on \mathbf{CD} is twice the sq. on \mathbf{ED} , that is,
twice the sq. on the opp. side \mathbf{PQ} . 1. 34.

Now the sqq. on \mathbf{AQ} , \mathbf{QB} = the sqq. on \mathbf{AQ} , \mathbf{QD} •
the sq. on \mathbf{AD} , for \mathbf{AQD} is a rt.
angle; 1. 47.

the sum of the sqq. on \mathbf{AC} , \mathbf{CD} ,
for \mathbf{ACD} is a rt. angle; 1. 17.

twice the sq. on \mathbf{AP} with twice
the sq. on \mathbf{PQ} . *Proved.*

That is,
the sum of the sqq. on \mathbf{AQ} , \mathbf{QB} is twice the sum of the sqq.
on \mathbf{AP} , \mathbf{PQ} . Q. E. D.

CORRESPONDING ALGEBRAICAL FORMULA.

The result of this proposition may be written

$$\mathbf{AQ}^2 + \mathbf{BQ}^2 = 2(\mathbf{AP}^2 + \mathbf{PQ}^2).$$

Let $\mathbf{AB} = 2a$; and $\mathbf{PQ} = b$;

then \mathbf{AP} and \mathbf{PB} each = a .

Also $\mathbf{AQ} = a + b$; and $\mathbf{BQ} = b - a$.

Hence we have

$$(a + b)^2 + (b - a)^2 = 2(a^2 + b^2).$$


EXERCISE.

Shew that the enunciations of Props. 9 and 10 may take the following form:

The sum of the squares on two straight lines is equal to twice the sum of the squares on half their sum and on half their difference.

PROPOSITION 11. PROBLEM.

To divide a given straight line into two parts, so that the rectangle contained by the whole and one part may be equal to the square on the other part.



Let AB be the given straight line.

It is required to divide it into two parts, so that the rectangle contained by the whole and one part may be equal to the square on the other part.

On AB describe the square $ACDB$. I. 46.

Bisect AC at E . I. 10.

Join EB .

Produce CA to F , making EF equal to EB . I. 3.

On AF describe the square $AFGH$. I. 46.

Then shall AB be divided at H , so that the rect. AB , BH is equal to the sq. on AH .

Produce GH to meet CD in K .

Then because CA is bisected at E , and produced to F ,
 \therefore the rect. CF , FA with the sq. on AE = the sq. on FE II. 6.
= the sq. on EB . Constr.

But the sq. on EB = the sum of the sqq. on AB , AE ,
for the angle EAB is a rt. angle. I. 47.
 \therefore the rect. CF , FA with the sq. on AE = the sum of the
sqq. on AB , AE .

From these take the sq. on AE :
then the rect. CF , FA = the sq. on AB .

But the rect. CF , FA = the fig. FK ; for $FA = FG$;
 and the sq. on AB = the fig. AD . *Constr.*
 \therefore the fig. FK = the fig. AD .

From these take the common fig. AK ,
 then the remaining fig. FH = the remaining fig. HD .


But the fig. HD = the rect. AB , BH ; for $BD = AB$;
 and the fig. FH is the sq. on AH .

\therefore the rect. AB , BH = the sq. on AH . Q.E.D.

DEFINITION. A straight line is said to be divided in **Medial Section** when the rectangle contained by the given line and one of its segments is equal to the square on the other segment.

The student should observe that this division may be *internal* or *external*.

Thus if the straight line AB is divided internally at H , and externally at H' , so that

- (i) $AB \cdot BH = AH^2$,
 (ii) $AB \cdot BH' = AH'^2$, 

we shall in either case consider that AB is divided in medial section.

The case of *internal* section is alone given in Euclid *ix*. 11; but a straight line may be divided *externally* in medial section by a similar process. See Ex. 21, p. 146.

ALGEBRAICAL ILLUSTRATION.

It is required to find a point H in AB , or AB produced, such that
 $AB \cdot BH = AH^2$.

Let AB contain a units of length, and let AH contain x units;

then $HB = a - x$;

and x must be such that $a(a - x) = x^2$,

or $x^2 + ax - a^2 = 0$.

Thus the construction for dividing a straight line in medial section corresponds to the algebraical solution of this quadratic equation.


EXERCISES.

In the figure of *ix*. 11, shew that

- (i) if CH is produced to meet BF at L , CL is at right angles to BF ;
- (ii) if BE and CH meet at O , AO is at right angles to CH ;
- (iii) the lines BG , DF , AK are parallel;
- (iv) CF is divided in medial section at A .

† PROPOSITION 12. THEOREM.

In an obtuse-angled triangle, if a perpendicular is drawn from either of the acute angles to the opposite side produced, the square on the side subtending the obtuse angle is greater than the squares on the sides containing the obtuse angle, by twice the rectangle contained by the side on which, when produced, the perpendicular falls, and the line intercepted without the triangle, between the perpendicular and the obtuse angle.



Let ABC be an obtuse-angled triangle, having the obtuse angle at C ; and let AD be drawn from A perp. to BC produced:

then shall the sq. on AB be greater than the sqq. on BC , CA , by twice the rect. BC , CD .

Because BD is divided into two parts at C ,
 \therefore the sq. on BD = the sum of the sqq. on BC , CD , with twice the rect. BC , CD . II. 4.

To each add the sq. on DA .

Then the sqq. on BD , DA = the sum of the sqq. on BC , CD , DA , with twice the rect. BC , CD .

But the sum of the sqq. on BD , DA = the sq. on AB ,
 for the angle at D is a rt. angle. I. 47.

Similarly the sum of the sqq. on CD , DA = the sq. on CA .



\therefore the sq. on AB = the sum of the sqq. on BC , CA , with twice the rect. BC , CD .

That is, the sq. on AB is greater than the sum of the sqq. on BC , CA by twice the rect. BC , CD . Q.E.D.

[For alternative Enunciations to Props. 12 and 13 and Exercises, see p. 142.]

PROPOSITION 13. THEOREM.

In every triangle the square on the side subtending an acute angle, is less than the squares on the sides containing that angle, by twice the rectangle contained by either of these sides, and the straight line intercepted between the perpendicular let fall on it from the opposite angle, and the acute angle.



Let ABC be any triangle having the angle at B an acute angle; and let AD be the perp. drawn from A to the opp. side BC :

then shall the sq. on AC be less than the sum of the sqq. on AB , BC , by twice the rect. CB , BD .

Now AD may fall within the triangle ABC , as in Fig. 1, or without it, as in Fig. 2.

Because (in Fig. 1. BC is divided into two parts at D ,
(in Fig. 2. BD is divided into two parts at C ,

\therefore in both cases,

the sum of the sqq. on CB , BD = twice the rect. CB , BD with the sq. on CD . II. 7.

To each add the sq. on DA .

Then the sum of the sqq. on CB , BD , DA = twice the rect. CB , BD with the sum of the sqq. on CD , DA .

But the sum of the sqq. on BD , DA = the sq. on AB ,

for the angle ADB is a rt. angle. I. 47.

Similarly the sum of the sqq. on CD , DA = the sq. on AC .

\therefore the sum of the sqq. on AB , BC , = twice the rect. CB , BD , with the sq. on AC .

That is, the sq. on AC is less than the sqq. on AB , BC by twice the rect. CB , BD . Q.E.D.

Obs. If the perpendicular **AD** coincides with **AC**, that is, if **ACB** is a right angle, it may be shewn that the proposition merely repeats the result of **r. 47**.

NOTE. The result of **Prop. 12** may be written

$$AB^2 = BC^2 + CA^2 + 2BC \cdot CD.$$

Remembering the definition of the **Projection** of a straight line given on page 97, the student will see that this proposition may be enunciated as follows:

In an obtuse-angled triangle the square on the side opposite the obtuse angle is greater than the sum of the squares on the sides containing the obtuse angle by twice the rectangle contained by either of those sides, and the projection of the other side upon it.

Prop. 13 may be written

$$AC^2 = AB^2 + BC^2 - 2CB \cdot BD,$$

and it may also be enunciated as follows:

In every triangle the square on the side subtending an acute angle, is less than the squares on the sides containing that angle, by twice the rectangle contained by either of these sides, and the projection of the other side upon it.

EXERCISES.

The following theorem should be noticed; it is proved by the help of **II. 1**.

1. If four points **A, B, C, D** are taken in order on a straight line, then will

$$AB \cdot CD + BC \cdot AD = AC \cdot BD.$$

ON **II. 12** AND **13**.


2. If from one of the base angles of an isosceles triangle a perpendicular is drawn to the opposite side, then twice the rectangle contained by that side and the segment adjacent to the base is equal to the square on the base.

3. If one angle of a triangle is one-third of two right angles, shew that the square on the opposite side is less than the sum of the squares on the sides forming that angle, by the rectangle contained by these two sides. [See **Ex. 10**, p. 101.]

4. If one angle of a triangle is two-thirds of two right angles, shew that the square on the opposite side is greater than the squares on the sides forming that angle, by the rectangle contained by these sides. [See **Ex. 10**, p. 101.]

PROPOSITION 14. PROBLEM.

To describe a square that shall be equal to a given rectilinear figure.



Let A be the given rectilinear figure.

It is required to describe a square equal to A.

Describe the par^m BCDE equal to the fig. A, and having the angle CBE a right angle. I. 45.

Then if BC = BE, the fig. BD is a square; and what was required is done.

But if not, produce BE to F, making EF equal to ED; I. 3.
and bisect BF at G. I. 10.

From centre G, with radius GF, describe the semicircle BHF; produce DE to meet the semicircle at H.

Then shall the sq. on EH be equal to the given fig. A.

Join GH.

Then because BF is divided equally at G and unequally at E,

\therefore the rect. BE, EF with the sq. on GE = the sq. on GF II. 5.
= the sq. on GH.

But the sq. on GH = the sum of the sq. on GE, EH; for the angle HEG is a rt. angle. I. 47.

\therefore the rect. BE, EF with the sq. on GE = the sum of the sq. on GE, EH.

From these take the sq. on GE:
then the rect. BE, EF = the sq. on HE.

But the rect. BE, EF = the fig. BD; for EF = ED; Constr.

and the fig. BD = the given fig. A. Constr.

\therefore the sq. on EH = the given fig. A. Q.E.F.

THEOREMS AND EXAMPLES ON BOOK II.

ON II. 4 AND 7.

1. *Shew by II. 4 that the square on a straight line is four times the square on half the line.*

[This result is constantly used in solving examples on Book II, especially those which follow from II. 12 and 13.]

2. *If a straight line is divided into any three parts, the square on the whole line is equal to the sum of the squares on the three parts together with twice the rectangles contained by each pair of these parts.*

Shew that the algebraical formula corresponding to this theorem is

$$(a + b + c)^2 = a^2 + b^2 + c^2 + 2bc + 2ca + 2ab.$$

3. *In a right-angled triangle, if a perpendicular is drawn from the right angle to the hypotenuse, the square on this perpendicular is equal to the rectangle contained by the segments of the hypotenuse.*

4. *In an isosceles triangle, if a perpendicular be drawn from one of the angles at the base to the opposite side, shew that the square on the perpendicular is equal to twice the rectangle contained by the segments of that side together with the square on the segment adjacent to the base.*

5. *Any rectangle is half the rectangle contained by the diagonals of the squares described upon its two sides.*

6. *In any triangle if a perpendicular is drawn from the vertical angle to the base, the sum of the squares on the sides forming that angle, together with twice the rectangle contained by the segments of the base, is equal to the square on the base together with twice the square on the perpendicular.*

ON II. 5 AND 6.

The student is reminded that these important propositions are both included in the following enunciation.


The difference of the squares on two straight lines is equal to the rectangle contained by their sum and difference.

7. *In a right-angled triangle the square on one of the sides forming the right angle is equal to the rectangle contained by the sum and difference of the hypotenuse and the other side. [I. 47 and II. 5.]*

8. *The difference of the squares on two sides of a triangle is equal to twice the rectangle contained by the base and the intercept between the middle point of the base and the foot of the perpendicular drawn from the vertical angle to the base.*

Let ABC be a triangle, and let P be the middle point of the base BC : let AQ be drawn perp. to BC .

Then shall $AB^2 - AC^2 = 2BC \cdot PQ$.



First, let AQ fall within the triangle.

$$\text{Now } AB^2 = BQ^2 + QA^2,$$

1. 17.

$$\text{also } AC^2 = QC^2 + QA^2,$$

$$\therefore AB^2 - AC^2 = BQ^2 - QC^2$$

Ax. 3.

$$= (BQ + QC)(BQ - QC)$$

II. 5.

$$= BC \cdot 2PQ$$

Ex. 1, p. 129.

$$= 2BC \cdot PQ.$$

Q.E.D.

The case in which AQ falls outside the triangle presents no difficulty.

9. *The square on any straight line drawn from the vertex of an isosceles triangle to the base is less than the square on one of the equal sides by the rectangle contained by the segments of the base.*

10. *The square on any straight line drawn from the vertex of an isosceles triangle to the base produced, is greater than the square on one of the equal sides by the rectangle contained by the segments into which the base is divided externally.*

11. *If a straight line is drawn through one of the angles of an equilateral triangle to meet the opposite side produced, so that the rectangle contained by the segments of the base is equal to the square on the side of the triangle; shew that the square on the line so drawn is double of the square on a side of the triangle.*

12. *If XY be drawn parallel to the base BC of an isosceles triangle ABC , then the difference of the squares on BY and CY is equal to the rectangle contained by BC , XY .* [See above, Ex. 8.]

13. *In a right-angled triangle, if a perpendicular be drawn from the right angle to the hypotenuse, the square on either side forming the right angle is equal to the rectangle contained by the hypotenuse and the segment of it adjacent to that side.*

ON II. 9 AND 10.

14. Deduce Prop. 9 from Props. 4 and 5, using also the theorem that the square on a straight line is four times the square on half the line.

15. Deduce Prop. 10 from Props. 7 and 6, using also the theorem mentioned in the preceding Exercise.

16. If a straight line is divided equally and also unequally, the squares on the two unequal segments are together equal to twice the rectangle contained by these segments together with four times the square on the line between the points of section.

ON II. 11.

17. If a straight line is divided internally in medial section, and from the greater segment a part be taken equal to the less: shew that the greater segment is also divided in medial section.

18. If a straight line is divided in medial section, the rectangle contained by the sum and difference of the segments is equal to the rectangle contained by the segments.

19. If AB is divided at H in medial section, and if X is the middle point of the greater segment AH , shew that a triangle whose sides are equal to AH , XH , BX respectively must be right-angled.

20. If a straight line AB is divided internally in medial section at H , prove that the sum of the squares on AB , BH is three times the square on AH .

21. Divide a straight line externally in medial section.


[Proceed as in II. 11, but instead of drawing EF , make EF' equal to EB in the direction remote from A ; and on AF' describe the square $AF'G'H'$ on the side remote from AB . Then AB will be divided externally at H' as required.]

ON II. 12 AND 13.

22. In a triangle ABC the angles at B and C are acute: if E and F are the feet of perpendiculars drawn from the opposite angles to the sides AC , AB , shew that the square on BC is equal to the sum of the rectangles AB , BF and AC , CE .

23. ABC is a triangle right-angled at C , and DE is drawn from a point D in AC perpendicular to AB : shew that the rectangle AB , AE is equal to the rectangle AC , AD .

24. *In any triangle the sum of the squares on two sides is equal to twice the square on half the third side together with twice the square on the median which bisects the third side.*



Let ABC be a triangle, and AP the median bisecting the side BC .
Then shall $AB^2 + AC^2 = 2BP^2 + 2AP^2$.

Draw AQ perp. to BC .

Consider the case in which AQ falls within the triangle, but does not coincide with AP .

Then of the angles APB , APC , one must be obtuse, and the other acute: let APB be obtuse

Then in the $\triangle APB$, $AB^2 = BP^2 + AP^2 + 2BP \cdot PQ$, II. 12.

Also in the $\triangle APC$, $AC^2 = CP^2 + AP^2 - 2CP \cdot PQ$. II. 13.

But $CP = BP$,

$\therefore CP^2 = BP^2$; and the rect. $BP, PQ =$ the rect. CP, PQ .

Hence adding the above results

$AB^2 + AC^2 = 2 \cdot BP^2 + 2 \cdot AP^2$. Q.E.D.

The student will have no difficulty in adapting this proof to the cases in which AQ falls without the triangle, or coincides with AP .

25. *The sum of the squares on the sides of a parallelogram is equal to the sum of the squares on the diagonals.*

26. *In any quadrilateral the squares on the diagonals are together equal to twice the sum of the squares on the straight lines joining the middle points of opposite sides.* [See Ex. 9, p. 97.]

27. *If from any point within a rectangle straight lines are drawn to the angular points, the sum of the squares on one pair of the lines drawn to opposite angles is equal to the sum of the squares on the other pair.*

28. *The sum of the squares on the sides of a quadrilateral is greater than the sum of the squares on its diagonals by four times the square on the straight line which joins the middle points of the diagonals.*

29. *O is the middle point of a given straight line AB , and from O as centre, any circle is described: if P be any point on its circumference, shew that the sum of the squares on AP , BP is constant.*

30. Given the base of a triangle, and the sum of the squares on the sides forming the vertical angle; find the locus of the vertex.

31. ABC is an isosceles triangle in which AB and AC are equal. AB is produced beyond the base to D , so that BD is equal to AB . Shew that the square on CD is equal to the square on AB together with twice the square on BC .

32. In a right-angled triangle the sum of the squares on the straight lines drawn from the right angle to the points of trisection of the hypotenuse is equal to five times the square on the line between the points of trisection.

33. Three times the sum of the squares on the sides of a triangle is equal to four times the sum of the squares on the medians.

34. ABC is a triangle, and O the point of intersection of its medians: shew that

$$AB^2 + BC^2 + CA^2 = 3(OA^2 + OB^2 + OC^2).$$

35. $ABCD$ is a quadrilateral, and X the middle point of the straight line joining the bisections of the diagonals; with X as centre any circle is described, and P is any point upon this circle: shew that $PA^2 + PB^2 + PC^2 + PD^2$ is constant, being equal to

$$XA^2 + XB^2 + XC^2 + XD^2 + 4XP^2.$$

36. The squares on the diagonals of a trapezium are together equal to the sum of the squares on its two oblique sides, with twice the rectangle contained by its parallel sides.

PROBLEMS.

37. Construct a rectangle equal to the difference of two squares.

38. Divide a given straight line into two parts so that the rectangle contained by them may be equal to the square described on a given straight line which is less than half the straight line to be divided.

39. Given a square and one side of a rectangle which is equal to the square, find the other side.

40. Produce a given straight line so that the rectangle contained by the whole line thus produced and the part produced, may be equal to the square on another given line.

41. Produce a given straight line so that the rectangle contained by the whole line thus produced and the given line shall be equal to the square on the part produced.

42. Divide a straight line AB into two parts at C , such that the rectangle contained by BC and another line X may be equal to the square on AC .


PART II.

BOOK III.

Book III. deals with the properties of Circles.

DEFINITIONS.

1. A **circle** is a plane figure bounded by one line, which is called the **circumference**, and is such that all straight lines drawn from a certain point within the figure to the circumference are equal to one another: this point is called the **centre** of the circle.



2. A **radius** of a circle is a straight line drawn from the centre to the circumference.

3. A **diameter** of a circle is a straight line drawn through the centre, and terminated both ways by the circumference.

4. A **semicircle** is the figure bounded by a diameter of a circle and the part of the circumference cut off by the diameter.

From these definitions we draw the following inferences:

(i) The distance of a point from the centre of a circle is less than the radius, if the point is within the circumference: and the distance of a point from the centre is greater than the radius, if the point is without the circumference.

(ii) A point is within a circle if its distance from the centre is less than the radius: and a point is without a circle if its distance from the centre is greater than the radius.

(iii) Circles of equal radius are equal in all respects; that is to say, their areas and circumferences are equal.

(iv) A circle is divided by any diameter into two parts which are equal in all respects.

5. Circles which have the same centre are said to be **concentric**.

6. An **arc** of a circle is any part of the circumference.

7. A **chord** of a circle is the straight line which joins any two points on the circumference.


From these definitions it may be seen that a chord of a circle, which does not pass through the centre, divides the circumference into two unequal arcs; of these, the greater is called the **major arc**, and the less the **minor arc**. Thus the major arc is *greater*, and the minor arc *less* than the semicircumference.

The major and minor arcs, into which a circumference is divided by a chord, are said to be **conjugate** to one another.




8. Chords of a circle are said to be **equidistant** from the centre, when the perpendiculars drawn to them from the centre are equal:


and one chord is said to be **further from the centre** than another, when the perpendicular drawn to it from the centre is greater than the perpendicular drawn to the other.




9. A **secant** of a circle is a straight line of indefinite length, which cuts the circumference in two points.



10. A **tangent** to a circle is a straight line which meets the circumference, but being produced, does not cut it. Such a line is said to **touch** the circle at a point; and the point is called the **point of contact**.




If a secant, which cuts a circle at the points **P** and **Q**, gradually changes its position in such a way that **P** remains fixed, the point **Q** will ultimately approach the fixed point **P**, until at length these points may be made to coincide. When the straight line **PQ** reaches this limiting position, it becomes the *tangent* to the circle at the point **P**.




Hence a tangent may be defined as a straight line which passes through *two coincident points* on the circumference.

11. Circles are said to **touch one another** when they meet, but do not cut one another.




When each of the circles which meet is *outside the other*, they are said to touch one another **externally**, or to have **external contact**; when one of the circles is *within the other*, they are said to touch one another **internally**, or to have **internal contact**.

12. A **segment** of a circle is the figure bounded by a chord and one of the two arcs into which the chord divides the circumference.




The chord of a segment is sometimes called its **base**.

13. An **angle in a segment** is one formed by two straight lines drawn from any point in the arc of the segment to the extremities of its chord.



[It will be shewn in Proposition 21, that all angles in the same segment of a circle are equal.]


14. An **angle at the circumference** of a circle is one formed by straight lines drawn from a point on the circumference to the extremities of an arc: such an angle is said to **stand upon** the arc, which it subtends.



15. **Similar segments** of circles are those which contain equal angles.



16. A **sector** of a circle is a figure bounded by two radii and the arc intercepted between them.




SYMBOLS AND ABBREVIATIONS.

In addition to the symbols and abbreviations given on page 10, we shall use the following.

\odot for circle, \odot^{ce} for circumference.

PROPOSITION I. PROBLEM.

To find the centre of a given circle.



Let ABC be a given circle:
it is required to find its centre.

In the given circle draw any chord AB ,
and bisect AB at D . I. 10.

From D draw DC at right angles to AB ; I. 11.
and produce DC to meet the circle at E and C .

Bisect EC at F . I. 10.

Then shall F be the centre of the circle ABC .

First, the centre of the circle must be in EC ;
for if not, let the centre be at a point G without EC .

Join AG , DG , BG .

Then in the \triangle^s GDA , GDB ,

Because $\left\{ \begin{array}{l} DA = DB, \quad \text{Constr.} \\ \text{and } GD \text{ is common;} \\ \text{and } GA = GB, \text{ for by supposition they are radii;} \end{array} \right.$
 \therefore the $\angle GDA =$ the $\angle GDB$; I. 8.

\therefore these angles, being adjacent, are rt. angles.

But the $\angle CDB$ is a rt. angle; Constr.

\therefore the $\angle GDB =$ the $\angle CDB$, Ax. 11.

the part equal to the whole, which is impossible.

$\therefore G$ is not the centre.

So it may be shewn that no point outside EC is the centre;
 \therefore the centre lies in EC .


$\therefore F$, the middle point of the diameter EC , must be the
centre of the circle ABC . Q.E.F.

COROLLARY. *The straight line which bisects a chord of
a circle at right angles passes through the centre.*

[For Exercises, see page 156.]

PROPOSITION 2. THEOREM.

If any two points are taken in the circumference of a circle, the chord which joins them falls within the circle.



Let ABC be a circle, and A and B any two points on its $circ$;

then shall the chord AB fall within the circle.

Find D , the centre of the $\odot ABC$; III. 1.

and in AB take any point E .

Join DA , DE , DB .

In the $\triangle DAB$, because $DA = DB$, III. Def. 1.

\therefore the $\angle DAB =$ the $\angle DBA$. I. 5.

But the ext. $\angle DEB$ is greater than the int. opp. $\angle DAE$; I. 16.

\therefore also the $\angle DEB$ is greater than the $\angle DBE$;

\therefore in the $\triangle DEB$, the side DB , which is opposite the greater angle, is greater than DE which is opposite the less: I. 19.

that is to say, DE is less than a radius of the circle;

$\therefore E$ falls within the circle.

So also any other point between A and B may be shewn to fall within the circle.


$\therefore AB$ falls within the circle. Q. E. D.

DEFINITION. A part of a curved line is said to be **concave** to a point when, *any* chord being taken in it, all straight lines drawn from the given point to the intercepted arc are cut by the chord: if, when any chord is taken, no straight line drawn from the given point to the intercepted arc is cut by the chord, the curve is said to be **convex** to that point.

Proposition 2 proves that the whole circumference of a circle is **concave** to its centre.

PROPOSITION 3. THEOREM.

*If a straight line drawn through the centre of a circle bisects a chord which does not pass through the centre, it shall cut it at right angles :
and, conversely, if it cut it at right angles, it shall bisect it.*



Let ABC be a circle; and let CD be a st. line drawn through the centre, and AB a chord which does not pass through the centre.

First.

Let CD bisect AB at F :

then shall CD cut AB at rt. angles.

Find E, the centre of the circle ; III. 1.

and join EA, EB.

Then in the \triangle^s AFE, BFE,

Because $\left\{ \begin{array}{l} AF = BF, \\ \text{and FE is common;} \\ \text{and AE = BE, being radii of the circle;} \end{array} \right. \quad \begin{array}{l} \text{Hyp.} \\ \\ \end{array}$

\therefore the \angle AFE = the \angle BFE; I. 8.

\therefore these angles, being adjacent, are rt. angles,

that is, DC cuts AB at rt. angles. Q. E. D.

Conversely. Let CD cut AB at rt. angles :

then shall CD bisect AB at F.

As before, find E the centre ; and join EA, EB.

In the \triangle EAB, because EA = EB, III. Def. 1.

\therefore the \angle EAB = the \angle EBA. I. 5.

Then in the \triangle^s EFA, EFB,

Because $\left\{ \begin{array}{l} \text{the } \angle \text{ EAF} = \text{the } \angle \text{ EBF,} \\ \text{and the } \angle \text{ EFA} = \text{the } \angle \text{ EFB, being rt. angles;} \\ \text{and EF is common;} \end{array} \right. \quad \begin{array}{l} \text{Proved.} \\ \text{Hyp.} \\ \end{array}$

\therefore AF = BF. I. 26.

Q. E. D.

[For Exercises, see page 156.]

EXERCISES.

ON PROPOSITION 1.

1. If two circles intersect at the points A , B , shew that the line which joins their centres bisects their common chord AB at right angles.

2. AB , AC are two equal chords of a circle; shew that the straight line which bisects the angle BAC passes through the centre.

3. Two chords of a circle are given in position and magnitude: find the centre of the circle.

4. Describe a circle that shall pass through three given points, which are not in the same straight line.

5. Find the locus of the centres of circles which pass through two given points.

6. Describe a circle that shall pass through two given points, and have a given radius.

ON PROPOSITION 2.

7. A straight line cannot cut a circle in more than two points.

ON PROPOSITION 3.

8. Through a given point within a circle draw a chord which shall be bisected at that point.

9. The parts of a straight line intercepted between the circumferences of two concentric circles are equal.

10. The line joining the middle points of two parallel chords of a circle passes through the centre.


11. Find the locus of the middle points of a system of parallel chords drawn in a circle.

12. If two circles cut one another, any two parallel straight lines drawn through the points of intersection to cut the circles, are equal.

13. PQ and XY are two parallel chords in a circle: shew that the points of intersection of PX , QY , and of PY , QX lie on the straight line which passes through the middle points of the given chords.

PROPOSITION 1. THEOREM.

If in a circle two chords cut one another, which do not both pass through the centre, they cannot both be bisected at their point of intersection.



Let $ABCD$ be a circle, and AC , BD two chords which intersect at E , but do not both pass through the centre:

then AC and BD shall not be *both* bisected at E .

CASE I. If *one* chord passes through the centre, it is a diameter, and the centre is its middle point;

\therefore it cannot be bisected by the other chord, which by hypothesis does not pass through the centre.

CASE II. If neither chord passes through the centre then, if possible, let E be the middle point of *both*;

that is, let $AE = EC$; and $BE = ED$.

Find F , the centre of the circle: III. 1.

• Join EF .

Then, because FE , which passes through the centre, bisects the chord AC , Hyp.

\therefore the $\angle FEC$ is a rt. angle. III. 3.

And because FE , which passes through the centre, bisects the chord BD , Hyp.

\therefore the $\angle FED$ is a rt. angle.

\therefore the $\angle FEC =$ the $\angle FED$,


the whole equal to its part, which is impossible.

$\therefore AC$ and BD are not *both* bisected at E . Q. E. D.

[For Exercises, see page 158.]

PROPOSITION 5. THEOREM.

If two circles cut one another, they cannot have the same centre.



Let the two \odot^s AGC, BFC cut one another at C:
then, they shall not have the same centre.

For, if possible, let the two circles have the same centre;
and let it be called E.

Join EC;

and from E draw any st. line to meet the \odot^s at F and G.

Then, because E is the centre of the \odot AGC, *Hyp.*

$\therefore EG = EC.$

And because E is also the centre of the \odot BFC, *Hyp.*

$\therefore EF = EC.$

$\therefore EG = EF,$

the whole equal to its part, which is impossible.

\therefore the two circles have not the same centre.

Q. E. D.

EXERCISES.

ON PROPOSITION 4.

1. If a parallelogram can be inscribed in a circle, the point of intersection of its diagonals must be at the centre of the circle.


2. Rectangles are the only parallelograms that can be inscribed in a circle.

ON PROPOSITION 5.

3. Two circles, which intersect at one point, must also intersect at another.

PROPOSITION 6. THEOREM.

If two circles touch one another internally, they cannot have the same centre.



Let the two \odot^s ABC, DEC touch one another internally at C:

then they shall not have the same centre.

For, if possible, let the two circles have the same centre;
and let it be called F.

Join FC;

and from F draw any st. line to meet the \odot^s at E and B.

Then, because F is the centre of the \odot ABC, *Hyp.*

$$\therefore FB = FC.$$

And because F is the centre of the \odot DEC, *Hyp.*

$$\therefore FE = FC.$$

$$\therefore FB = FE;$$

the whole equal to its part, which is impossible.

\therefore the two circles have not the same centre. Q.E.D.

NOTE. From Propositions 5 and 6 it is seen that circles, whose circumferences have any point in common, cannot be concentric, unless they coincide entirely.


Conversely, the circumferences of concentric circles can have no point in common.

PROPOSITION 7. THEOREM

If from any point within a circle which is not the centre, straight lines are drawn to the circumference, the greatest is that which passes through the centre; and the least is that which, when produced backwards, passes through the centre:

and of all other such lines, that which is nearer to the greatest is always greater than one more remote:

also two equal straight lines, and only two, can be drawn from the given point to the circumference, one on each side of the diameter.



Let ABCD be a circle, within which any point F is taken, which is not the centre: let FA, FB, FC, FG be drawn to the \bigcirc^e , of which FA passes through E the centre, and FB is nearer than FC to FA, and FC nearer than FG: and let FD be the line which, when produced backwards, passes through the centre: then of all these st. lines

- (i) FA shall be the greatest;
- (ii) FD shall be the least;
- (iii) FB shall be greater than FC, and FC greater than FG;

(iv) also two, and only two, equal st. lines can be drawn from F to the \bigcirc^e .

Join EB, EC, EG.

(i) Then in the $\triangle FEB$, the two sides FE, EB are together greater than the third side FB. I. 20.

But EB = EA, being radii of the circle;

\therefore FE, EA are together greater than FE;
that is, FA is greater than FB.

Similarly FA may be shewn to be greater than any other st. line drawn from F to the \bigcirc^{co} ;

$\therefore FA$ is the greatest of all such lines.

(ii) In the $\triangle EFG$, the two sides EF, FG are together greater than EG ; I. 20.

and $EG = ED$, being radii of the circle,

$\therefore EF, FG$ are together greater than ED .

Take away the common part EF ;

then FG is greater than FD .

Similarly any other st. line drawn from F to the \bigcirc^{co} may be shewn to be greater than FD .

$\therefore FD$ is the least of all such lines.

(iii) In the $\triangle^s BEF, CEF$, III. Def. 1.
Because $\left\{ \begin{array}{l} BE = CE, \\ \text{and } EF \text{ is common;} \end{array} \right.$

but the $\angle BEF$ is greater than the $\angle CEF$;

$\therefore FB$ is greater than FC . I. 24.

Similarly it may be shewn that FC is greater than FG .

(iv) At E in FE make the $\angle FEH$ equal to the $\angle FEG$. I. 23.

Join FH .

Then in the $\triangle^s GEF, HEF$,

Because $\left\{ \begin{array}{l} GE = HE, \\ \text{and } EF \text{ is common;} \end{array} \right.$ III. Def. 1.
also the $\angle GEF = \text{the } \angle HEF$; Constr.
 $\therefore FG = FH$. I. 4.

And besides FH no other straight line can be drawn from F to the \bigcirc^{co} equal to FG .

For, if possible, let $FK = FG$.

Then, because $FH = FG$, Proved.

$\therefore FK = FH$,

that is, a line nearer to FA , the greatest, is equal to a line which is more remote; which is impossible. Proved.


\therefore two, and only two, equal st. lines can be drawn from F to the \bigcirc^{co} . Q. E. D.

PROPOSITION 8. THEOREM.

If from any point without a circle straight lines are drawn to the circumference, of those which fall on the concave circumference, the greatest is that which passes through the centre; and of others, that which is nearer to the greatest is always greater than one more remote:

further, of those which fall on the convex circumference, the least is that which, when produced, passes through the centre; and of others that which is nearer to the least is always less than one more remote:

** lastly, from the given point there can be drawn to the circumference two, and only two, equal straight lines, one on each side of the shortest line.*



Let BGD be a circle of which C is the centre; and let A be any point outside the circle: let ABD, AEH, AFG, be st. lines drawn from A, of which AD passes through C, the centre, and AH is nearer than AG to AD:

then of st. lines drawn from A to the concave \odot^{ce} ,

(i) AD shall be the greatest, and (ii) AH greater than AG:

and of st. lines drawn from A to the convex \odot^{ce} ,

(iii) AB shall be the least, and (iv) AE less than AF.

(v) Also two, and only two, equal st. lines can be drawn from A to the \odot^{ce} .

Join CH, CG, CF, CE.

(i) Then in the $\triangle ACH$, the two sides AC, CH are together greater than AH: I. 20.

but CH = CD, being radii of the circle;

\therefore AC, CD are together greater than AH:

that is, AD is greater than AH.

Similarly AD may be shewn to be greater than any other st. line drawn from A to the concave \odot^{ce} ;

\therefore AD is the greatest of all such lines.

(ii) In the $\triangle HCA, GCA$,
 Because $\left\{ \begin{array}{l} HC = GC, \quad \text{III. Def. 1.} \\ \text{and } CA \text{ is common;} \\ \text{but the } \angle HCA \text{ is greater than the } \angle GCA; \end{array} \right.$
 $\therefore AH$ is greater than AG . I. 24.

(iii) In the $\triangle AEC$, the two sides AE, EC are together greater than AC : I. 20.

but $EC = BC$; III. Def. 1
 \therefore the remainder AE is greater than the remainder AB .

Similarly any other st. line drawn from A to the convex \circ^e may be shewn to be greater than AB ;

$\therefore AB$ is the least of all such lines.

(iv) In the $\triangle AFC$, because AE, EC are drawn from the extremities of the base to a point E within the triangle,

$\therefore AF, FC$ are together greater than AE, EC . I. 21.

But $FC = EC$, III. Def. 1.
 \therefore the remainder AF is greater than the remainder AE .

(v) At C , in $\triangle AC$, make the $\angle ACM$ equal to the $\angle ACE$.
 Join AM .

Then in the two $\triangle ECA, MCA$,

Because $\left\{ \begin{array}{l} EC = MC, \quad \text{III. Def. 1.} \\ \text{and } CA \text{ is common;} \\ \text{also the } \angle ECA = \text{the } \angle MCA; \end{array} \right.$ Constr.
 $\therefore AE = AM$; I. 4.

and besides AM , no st. line can be drawn from A to the \circ^e , equal to AE .

For, if possible, let $AK = AE$:

then because $AM = AE$, Proved.
 $\therefore AM = AK$;

that is, a line nearer to the shortest line is equal to a line which is more remote; which is impossible. Proved.


\therefore two, and only two, equal st. lines can be drawn from A to the \circ^e . Q.E.D.

Where are the limits of that part of the circumference which is concave to the point A ?

Obs. Of the following proposition Euclid gave two distinct proofs, the first of which has the advantage of being *direct*.

PROPOSITION 9. THEOREM. [FIRST PROOF].

If from a point within a circle more than two equal straight lines can be drawn to the circumference, that point is the centre of the circle.



Let ABC be a circle, and D a point within it, from which more than two equal st. lines are drawn to the \bigcirc^e , namely DA, DB, DC :

then D shall be the centre of the circle.

Join AB, BC :

and bisect AB, BC at E and F respectively. I. 10.

Join DE, DF .

Then in the \triangle^s DEA, DEB ,

Because $\begin{cases} EA = EB, & \text{Constr.} \\ \text{and } DE \text{ is common;} \\ \text{and } DA = DB; & \text{Hyp.} \end{cases}$

\therefore the $\angle DEA =$ the $\angle DEB$; I. 8.

\therefore these angles, being adjacent, are rt. angles.


Hence ED , which bisects the chord AB at rt. angles, must pass through the centre. III. 1. Cor.

Similarly it may be shewn that FD passes through the centre.

$\therefore D$, which is the only point common to ED and FD , must be the centre. Q.E.D.

PROPOSITION 9. THEOREM. [SECOND PROOF.]

If from a point within a circle more than two equal straight lines can be drawn to the circumference, that point is the centre of the circle.



Let ABC be a circle, and D a point within it, from which more than two equal st. lines are drawn to the \odot^e , namely DA, DB, DC :

then D shall be the centre of the circle.

• For, if not, suppose E to be the centre.

Join DE , and produce it to meet the \odot^e at F, G .

Then because D is a point within the circle, not the centre, and because DF passes through the centre E ;

$\therefore DA$, which is nearer to DF , is greater than DB , which is more remote: • • III. 7.

but this is impossible, since by hypothesis, DA, DB , are equal.

$\therefore E$ is not the centre of the circle.

*And wherever we suppose the centre E to be, otherwise than at D , two at least of the st. lines DA, DB, DC may be shewn to be unequal, which is contrary to hypothesis.

$\therefore D$ is the centre of the $\odot ABC$.


Q.E.D.

* NOTE. For example, if the centre E were supposed to be within the angle BDC , then DB would be greater than DA ; if within the angle ADB , then DB would be greater than DC ; if on one of the three straight lines, as DB , then DB would be greater than both DA and DC .

Obs. Two proofs of Proposition 10, both indirect, were given by Euclid.

PROPOSITION 10. THEOREM. [FIRST PROOF.]

One circle cannot cut another at more than two points.



If possible, let $DABC$, $EABC$ be two circles, cutting one another at more than two points, namely at A , B , C .

Join AB , BC .

Draw FH , bisecting AB at rt. angles; I. 10, 11.
and draw GH bisecting BC at rt. angles.

Then because AB is a chord of *both* circles, and FH bisects it at rt. angles,

\therefore the centre of both circles lies in FH . III. 1. *Cor.*

Again, because BC is a chord of both circles, and GH bisects it at right angles,

\therefore the centre of both circles lies in GH . III. 1. *Cor.*

Hence H , the only point common to FH and GH , is the centre of both circles;

which is impossible, for circles which cut one another cannot have a common centre. III. 5.

\therefore one circle cannot cut another at more than two points.

Q.E.D.


COROLLARIES. (i) *Two circles cannot meet in three points without coinciding entirely.*

(ii) *Two circles cannot have a common arc without coinciding entirely.*

(iii) *Only one circle can be described through three points, which are not in the same straight line*

PROPOSITION 10. THEOREM. [SECOND PROOF.]

One circle cannot cut another at more than two points.



If possible, let **DABC**, **EABC** be two circles, cutting one another at more than two points, namely at **A**, **B**, **C**.

Find **H**, the centre of the \odot **DABC**,
and join **HA**, **HB**, **HC**. III. 1.

Then since **H** is the centre of the \odot **DABC**,

\therefore **HA**, **HB**, **HC** are all equal. III. Def. 1.

And because **H** is a point within the \odot **EABC**, from which more than two equal st. lines, namely **HA**, **HB**, **HC** are drawn to the \odot^{ce} ,

\therefore **H** is the centre of the \odot **EABC**: III. 9.


that is to say, the two circles have a common centre **H**,
but this is impossible, since they cut one another. III. 5.

Therefore one circle cannot cut another in more than two points. Q.E.D.

NOTE. This proof is imperfect, because it assumes that the centre of the circle **DABC** must fall within the circle **EABC**; whereas it may be conceived to fall either without the circle **EABC**, or on its circumference. Hence to make the proof complete, two additional cases are required.

PROPOSITION 11. THEOREM.

If two circles touch one another internally, the straight line which joins their centres, being produced, shall pass through the point of contact.



Let ABC and ADE be two circles which touch one another internally at A ; let F be the centre of the $\odot ABC$, and G the centre of the $\odot ADE$:

then shall FG produced pass through A .

If not, let it pass otherwise, as $FGEH$.

Join FA , GA .

Then in the $\triangle FGA$, the two sides FG , GA are together greater than FA : I. 20.

but $FA = FH$, being radii of the $\odot ABC$: Hyp.

$\therefore FG$, GA are together greater than FH .

Take away the common part FG ;

then GA is greater than GH .

But $GA = GE$, being radii of the $\odot ADE$: Hyp.

$\therefore GE$ is greater than GH ,

the part greater than the whole; which is impossible.

$\therefore FG$, when produced, must pass through A .

Q. E. D.


EXERCISES.

1. If the distance between the centres of two circles is equal to the difference of their radii, then the circles must meet in one point, but in no other; that is, they must touch one another.

2. If two circles whose centres are A and B touch one another internally, and a straight line be drawn through their point of contact, cutting the circumferences at P and Q ; shew that the radii AP and BQ are parallel.

PROPOSITION 12. THEOREM.

If two circles touch one another externally, the straight line which joins their centres shall pass through the point of contact.



Let ABC and ADE be two circles which touch one another externally at A ; let F be the centre of the $\odot ABC$, and G the centre of the $\odot ADE$

then shall FG pass through A .

If not, let FG pass otherwise, as $FHKG$.

Join FA , GA .

Then in the $\triangle FAG$, the two sides FA , GA are together greater than FG ;

I. 20.

but $FA = FH$, being radii of the $\odot ABC$;

Hyp.

and $GA = GK$, being radii of the $\odot ADE$;

Hyp.

$\therefore FH$ and GK are together greater than FG ;

which is impossible.

$\therefore FG$ must pass through A .

Q.E.D.

EXERCISES.

1. Find the locus of the centres of all circles which touch a given circle at a given point.


2. Find the locus of the centres of all circles of given radius, which touch a given circle.

3. If the distance between the centres of two circles is equal to the sum of their radii, then the circles meet in one point, but in no other; that is, they touch one another.

4. If two circles whose centres are A and B touch one another externally, and a straight line be drawn through their point of contact cutting the circumferences at P and Q ; shew that the radii AP and BQ are parallel.

PROPOSITION 13. THEOREM.

Two circles cannot touch one another at more than one point, whether internally or externally.



If possible, let **ABC**, **EDF** be two circles which touch one another at more than one point, namely at **B** and **D**.

Join **BD**;

and draw **GF**, bisecting **BD** at rt. angles. I. 10, 11.

Then, whether the circles touch one another internally, as in Fig. 1, or externally as in Fig. 2,

because **B** and **D** are on the *circles* of both circles,

\therefore **BD** is a chord of both circles :

\therefore the centres of both circles lie in **GF**, which bisects **BD** at rt. angles. III. 1. *Cor.*

Hence **GF** which joins the centres must pass through a point of contact ; III. 11, and 12.

which is impossible, since **B** and **D** are without **GF**.

\therefore two circles cannot touch one another at more than one point.

Q. E. D.

NOTE. It must be observed that the proof here given applies, by virtue of Propositions 11 and 12, to *both* the above figures: we have therefore omitted the separate discussion of Fig. 2, which finds a place in most editions based on Simson's text.

EXERCISES ON PROPOSITIONS 1- 13.

1. Describe a circle to pass through two given points and have its centre on a given straight line. When is this impossible?

2. All circles which pass through a fixed point, and have their centres on a given straight line, pass also through a second fixed point.

3. Describe a circle of given radius to touch a given circle at a given point. How many solutions will there be? When will there be only one solution?

4. From a given point as centre describe a circle to touch a given circle. How many solutions will there be?

5. Describe a circle to pass through a given point, and touch a given circle at a given point. [See Ex. 1, p. 169 and Ex. 5, p. 156.] When is this impossible?

6. Describe a circle of given radius to touch two given circles. [See Ex. 2, p. 169.] How many solutions will there be?

7. Two parallel chords of a circle are six inches and eight inches in length respectively, and the perpendicular distance between them is one inch: find the radius.

8. If two circles touch one another externally, the straight lines, which join the extremities of parallel diameters towards opposite parts, must pass through the point of contact.


9. Find the greatest and least straight lines which have one extremity on each of two given circles, which do not intersect.

10. In any segment of a circle, of all straight lines drawn at right angles to the chord and intercepted between the chord and the arc, the greatest is that which passes through the middle point of the chord; and of others that which is nearer the greatest is greater than one more remote.

11. If from any point on the circumference of a circle straight lines be drawn to the circumference, the greatest is that which passes through the centre; and of others, that which is nearer to the greatest is greater than one more remote; and from this point there can be drawn to the circumference two, and only two, equal straight lines.

PROPOSITION 14. THEOREM.

*Equal chords in a circle are equidistant from the centre
and, conversely, chords which are equidistant from the
centre are equal.*



Let ABC be a circle, and let AB and CD be chords,
of which the perp. distances from the centre are EF
and EG.

First, Let $AB = CD$:
then shall AB and CD be equidistant from the centre E.

Join EA, EC.

Then, because EF, which passes through the centre, is
perp. to the chord AB; *Hyp.*

\therefore EF bisects AB; III. 3.

that is, AB is double of FA.

For a similar reason, CD is double of GC.

But $AB = CD$, *Hyp.*

\therefore FA = GC.

Now $EA = EC$, being radii of the circle;

\therefore the sq. on EA = the sq. on EC.

But the sq. on EA = the sqq. on EF, FA;

for the \angle EFA is a rt. angle. I. 17.

And the sq. on EC = the sqq. on EG, GC;

for the \angle EGC is a rt. angle.

\therefore the sqq. on EF, FA = the sqq. on EG, GC.

Now of these, the sq. on FA = the sq. on GC; for $FA = GC$.

\therefore the sq. on EF = the sq. on EG,

\therefore EF = EG;

that is, the chords AB, CD are equidistant from the centre.

Q.E.D.

Conversely. Let AB and CD be equidistant from the centre E ;

that is, let $EF = EG$;

• then shall $AB = CD$. •

For, the same construction being made, it may be shewn as before that AB is double of FA , and CD double of GC ;

and that the sqq. on EF , $FA =$ the sqq. on EG , GC .

Now of these, the sq. on $EF =$ the sq. on EG ,

for $EF = EG$;

Hyp

∴ the sq. on $FA =$ the square on GC ;

∴ $FA = GC$;

and doubles of these equals are equal,

that is, $AB = CD$.

Q. E. D.


EXERCISES.

1. Find the locus of the middle points of equal chords of a circle.
2. If two chords of a circle cut one another, and make equal angles with the straight line which joins their point of intersection to the centre, they are equal.
3. If two equal chords of a circle intersect, shew that the segments of the one are equal respectively to the segments of the other.
4. In a given circle draw a chord which shall be equal to one given straight line (not greater than the diameter) and parallel to another.
5. PQ is a fixed chord in a circle, and AB is any diameter: shew that the sum or difference of the perpendiculars let fall from A and B on PQ is constant, that is, the same for all positions of AB .

PROPOSITION 15. THEOREM.

*The diameter is the greatest chord in a circle ;
and of others, that which is nearer to the centre is greater
than one more remote.*

*conversely, the greater chord is nearer to the centre than
the less.*



- Let ABCD be a circle, of which AD is a diameter, and E the centre ; and let BC and FG be any two chords, whose perp. distances from the centre are EH and EK :
then (i) AD shall be greater than BC :
(ii) if EH is less than EK, BC shall be greater than FG :
(iii) if BC is greater than FG, EH shall be less than EK.

- (i) Join EB, EC.
Then in the $\triangle BEC$, the two sides BE, EC are together greater than BC, I. 20.
but BE = AE, III. Def. 1.
and EC = ED ;
 \therefore AE and ED together are greater than BC ;
that is, AD is greater than BC.
Similarly AD may be shewn to be greater than any other chord, not a diameter.

- (ii) Let EH be less than EK ;
then BC shall be greater than FG.
Join EF.

Since EH, passing through the centre, is perp. to the chord BC,
 \therefore EH bisects BC, III. 3.

that is, BC is double of HB .

For a similar reason FG is double of KF .

Now $EB = EF$,

III. *Def.* 1.

\therefore the sq. on $EB =$ the sq. on EF .

But the sq. on $EB =$ the sqq. on EH, HB ;

for the $\angle EHB$ is a rt. angle;

I. 47.

also the sq. on $EF =$ the sqq. on EK, KF ;

for the $\angle EKF$ is a rt. angle.

\therefore the sqq. on $EH, HB =$ the sqq. on EK, KF .

But the sq. on EH is less than the sq. on EK ,

for EH is less than EK ;

Hyp.

\therefore the sq. on HB is greater than the sq. on KF ;

$\therefore HB$ is greater than KF ;

hence BC is greater than FG .

(iii) Let BC be greater than FG ;
then EH shall be less than EK .

For since BC is greater than FG ,

Hyp.

$\therefore HB$ is greater than KF ;

\therefore the sq. on HB is greater than the sq. on KF .

But the sqq. on $EH, HB =$ the sqq. on EK, KF : *Proved.*

\therefore the sq. on EH is less than the sq. on EK ;

$\therefore EH$ is less than EK .

Q.E.D.

EXERCISES.

1. Through a given point within a circle draw the least possible chord.

2. AB is a fixed chord of a circle, and XY any other chord having its middle point Z on AB : what is the greatest, and what the least length that XY may have?

Shew that XY increases, as Z approaches the middle point of AB .

3. In a given circle draw a chord of given length, having its middle point on a given chord.


When is this problem impossible?

Obs. Of the following proofs of Proposition 16, the second (by reductio ad absurdum) is that given by Euclid; but the first is to be preferred, as it is *direct*, and not less simple than the other.

✓ PROPOSITION 16. THEOREM. [ALTERNATIVE PROOF.]

The straight line drawn at right angles to a diameter of a circle at one of its extremities is a tangent to the circle;

and every other straight line drawn through this point cuts the circle.



Let AKB be a circle, of which E is the centre, and AB a diameter; and through B let the st. line CBD be drawn at rt. angles to AB:

- then (i) CBD shall be a tangent to the circle;
 (ii) any other st. line through B, as BF, shall cut the circle.

(i) In CD take any point G, and join EG.

Then, in the $\triangle EBG$, the $\angle EBG$ is a rt. angle; *Hyp.*

\therefore the $\angle EGB$ is less than a rt. angle; I. 17.

\therefore the $\angle EBG$ is greater than the $\angle EGB$;

\therefore EG is greater than EB: I. 19.

that is, EG is greater than a radius of the circle;

\therefore the point G is without the circle.

Similarly any other point in CD, except B, may be shewn to be outside the circle:


hence CD meets the circle at B, but being produced, does not cut it;

that is, CD is a tangent to the circle. III. Def. 10.

- (ii) Draw \square EH perp. to BF . I. 12.
 Then in the $\triangle EHB$, because the $\angle EHB$ is a rt. angle,
 \therefore the $\angle EBH$ is less than a rt. angle; I. 17.
 $\therefore EB$ is greater than EH ; I. 19.
 that is, EH is less than a radius of the circle;
 $\therefore H$, a point in BF , is within the circle;
 $\therefore BF$ must cut the circle. Q. E. D.

PROPOSITION 16. THEOREM. [EUCLID'S PROOF.]

*The straight line drawn at right angles to a diameter of a circle at one of its extremities, is a tangent to the circle;
 and no other straight line can be drawn through this point so as not to cut the circle.*



Let ABC be a circle, of which D is the centre, and AB a diameter; let AE be drawn at rt. angles to BA , at its extremity A :

- (i) then shall AE be a tangent to the circle.

For, if not, let AE cut the circle at C .
 Join DC .

Then in the $\triangle DAC$, because $DA = DC$, III. Def. 1.
 \therefore the $\angle DAC =$ the $\angle DCA$.


But the $\angle DAC$ is a rt. angle; Hyp.
 \therefore the $\angle DCA$ is a rt. angle;

that is, two angles of the $\triangle DAC$ are together equal to two rt. angles; which is impossible. I. 17.

Hence AE meets the circle at A , but being produced, does not cut it;

that is, AE is a tangent to the circle. III. Def. 10.

(ii) Also through A no other straight line but AE can be drawn so as not to cut the circle.



For, if possible, let AF be another st. line drawn through A so as not to cut the circle.

From D draw DG perp. to AF ; I. 12.
and let DG meet the C^e at H .

Then in the $\triangle DAG$, because the $\angle DGA$ is a rt. angle,
 \therefore the $\angle DAG$ is less than a rt. angle; I. 17.

$\therefore DA$ is greater than DG . I. 19.

But $DA = DH$, III. Def. 1.

$\therefore DH$ is greater than DG ,

the part greater than the whole, which is impossible.

\therefore no st. line can be drawn from the point A , so as not to cut the circle, except AE .

COROLLARIES. (i) *A tangent touches a circle at one point only.*

(ii) *There can be but one tangent to a circle at a given point.*

PROPOSITION 17. PROBLEM.

To draw a tangent to a circle from a given point either on, or without the circumference.

Fig. 1





Fig. 2



Let BCD be the given circle, and A the given point:
it is required to draw from A a tangent to the \odot CDB.

CASE I. If the given point A is on the \odot^{ce} .

Find E, the centre of the circle. III. 1.

Join EA.

At A draw AK at rt. angles to EA. I. 11.

Then AK being perp. to a diameter at one of its extremities,
is a tangent to the circle. III. 16.

CASE II. If the given point A is without the \odot^{ce} .

Find E, the centre of the circle; III. 1.

and join AE, cutting the \odot BCD at D.

From centre E, with radius EA, describe the \odot AFG.

At D, draw GDF at rt. angles to EA, cutting the \odot AFG at
F and G. I. 11.

Join EF, EG, cutting the \odot BCD at B and C.

Join AB, AC.


Then both AB and AC shall be tangents to the \odot CDB.

For in the \triangle^s AEB, FED,

Because $\left\{ \begin{array}{l} AE = FE, \text{ being radii of the } \odot \text{ GAF;} \\ \text{and } EB = ED, \text{ being radii of the } \odot \text{ BDC;} \\ \text{and the included angle } AEF \text{ is common;} \end{array} \right.$
 \therefore the \angle ABE = the \angle FDE. I. 4.

PROPOSITION 18. THEOREM.

The straight line drawn from the centre of a circle to the point of contact of a tangent is perpendicular to the tangent.



Let ABC be a circle, of which F is the centre;
and let the st. line DE touch the circle at C ;
then shall FC be perp. to DE .

For, if not, suppose FG to be perp. to DE , I. 12
and let it meet the circ^e at B .

Then in the $\triangle FCG$, because the $\angle FGC$ is a rt. angle, *Hyp.*

\therefore the $\angle FCG$ is less than a rt. angle: I. 17.

\therefore the $\angle FGC$ is greater than the $\angle FCG$;

$\therefore FC$ is greater than FG : I. 19.

but $FC = FB$; — — —

$\therefore FB$ is greater than FG ,

the part greater than the whole, which is impossible.

$\therefore FC$ cannot be otherwise than perp. to DE :

that is, FC is perp. to DE . Q.E.D.

EXERCISES.

1. Draw a tangent to a circle (i) parallel to, (ii) at right angles to a given straight line.

2. Tangents drawn to a circle from the extremities of a diameter are parallel.


3. Circles which touch one another internally or externally have a common tangent at their point of contact.

4. In two concentric circles any chord of the outer circle which touches the inner, is bisected at the point of contact.

5. In two concentric circles, all chords of the outer circle which touch the inner, are equal.

PROPOSITION 19. THEOREM

The straight line drawn perpendicular to a tangent to a circle from the point of contact passes through the centre.



Let ABC be a circle, and DE a tangent to it at the point C ;
and let CA be drawn perp. to DE :

then shall CA pass through the centre.

For if not, suppose the centre to be outside CA , as at F .

Join CF .

Then because DE is a tangent to the circle, and FC is drawn from the centre F to the point of contact,

\therefore the $\angle FCE$ is a rt. angle. III. 18.

But the $\angle ACE$ is a rt. angle ; *Hyp.*

\therefore the $\angle FCE =$ the $\angle ACE$;

the part equal to the whole, which is impossible.

\therefore the centre cannot be otherwise than in CA ;

that is, CA passes through the centre.

Q.E.D.

EXERCISES ON THE TANGENT.

PROPOSITIONS 16, 17, 18, 19.

1. *The centre of any circle which touches two intersecting straight lines must lie on the bisector of the angle between them.*

2. AB and AC are two tangents to a circle whose centre is O ;
shew that AO bisects the chord of contact BC at right angles.

3. If two circles are concentric all tangents drawn from points on the circumference of the outer to the inner circle are equal.

4. The diameter of a circle bisects all chords which are parallel to the tangent at either extremity.

5. Find the locus of the centres of all circles which touch a given straight line at a given point.

6. Find the locus of the centres of all circles which touch each of two parallel straight lines.

7. Find the locus of the centres of all circles which touch each of two intersecting straight lines of unlimited length.

8. Describe a circle of given radius to touch two given straight lines.

9. Through a given point, within or without a circle, draw a chord equal to a given straight line.

In order that the problem may be possible, between what limits must the given line lie, when the given point is (i) without the circle, (ii) within it?

10. Two parallel tangents to a circle intercept on any third tangent a segment which subtends a right angle at the centre.

11. In any quadrilateral circumscribed about a circle, the sum of one pair of opposite sides is equal to the sum of the other pair.

12. Any parallelogram which can be circumscribed about a circle, must be equilateral.

13. If a quadrilateral be described about a circle, the angles subtended at the centre by any two opposite sides are together equal to two right angles.

14. AB is any chord of a circle, AC the diameter through A, and AD the perpendicular on the tangent at B: shew that AB bisects the angle DAC.

15. Find the locus of the extremities of tangents of fixed length drawn to a given circle.

16. In the diameter of a circle produced, determine a point such that the tangent drawn from it shall be of given length.

17. In the diameter of a circle produced, determine a point such that the two tangents drawn from it may contain a given angle.


18. Describe a circle that shall pass through a given point, and touch a given straight line at a given point. [See page 183. Ex. 5.]

19. Describe a circle of given radius, having its centre on a given straight line, and touching another given straight line.

20. Describe a circle that shall have a given radius, and touch a given circle and a given straight line. How many such circles can be drawn?

PROPOSITION 20. THEOREM.

The angle at the centre of a circle is double of an angle at the circumference, standing on the same arc.



Let ABC be a circle, of which E is the centre; and let BEC be an angle at the centre, and BAC an angle at the C^{th} , standing on the same arc BC :

then shall the $\angle BEC$ be double of the $\angle BAC$.

Join AE , and produce it to F .

CASE I. When the centre E is within the angle BAC .

Then in the $\triangle EAB$, because $EA = EB$,

\therefore the $\angle EAB =$ the $\angle EBA$; I. 5.

\therefore the sum of the $\angle^s EAB, EBA =$ twice the $\angle EAB$.

But the ext. $\angle BCF =$ the sum of the $\angle^s EAB, EBA$; I. 32.

\therefore the $\angle BEF =$ twice the $\angle EAB$.

Similarly the $\angle FEC =$ twice the $\angle EAC$.

\therefore the sum of the $\angle^s BEF, FEC =$ twice the sum of

the $\angle^s EAB, EAC$;

that is, the $\angle BEC =$ twice the $\angle BAC$.

CASE II. When the centre E is without the $\angle BAC$.

As before, it may be shewn that the $\angle FEB =$ twice the $\angle FAB$;


also the $\angle FEC =$ twice the $\angle FAC$;

\therefore the difference of the $\angle^s FEC, FEB =$ twice the difference of the $\angle^s FAC, FAB$;

that is, the $\angle BEC =$ twice the $\angle BAC$.


Q. E. D.

NOTE. If the arc BFC , on which the angles stand, is greater than a semi-circumference, it is clear that the angle BEC at the centre will be reflex: but it may still be shewn as, in Case I., that the reflex $\angle BEC$ is double of the $\angle BAC$ at the \odot^{ce} , standing on the same arc BFC .



PROPOSITION 21. THEOREM.

Angles in the same segment of a circle are equal.



Let $ABCD$ be a circle, and let BAD , BED be angles in the same segment $BAED$:

then shall the $\angle BAD =$ the $\angle BED$.

Find F , the centre of the circle. III. 1.

CASE I. When the segment $BAED$ is greater than a semicircle


Join BF , DF .

Then the $\angle BFD$ at the centre = twice the $\angle BAD$ at the \odot^{ce} , standing on the same arc BD : III. 20.

and similarly the $\angle BFD =$ twice the $\angle BED$. III. 20.

\therefore the $\angle BAD =$ the $\angle BED$.

CASE II. When the segment $BAED$ is not greater than a semicircle.



Join AF , and produce it to meet the \odot^c at C .

Join EC .

Then since $AEDC$ is a semicircle;

\therefore the segment $BAEC$ is greater than a semicircle;

\therefore the $\angle BAC =$ the $\angle BEC$, in this segment. *Case 1.*

Similarly the segment $CAED$ is greater than a semicircle;

\therefore the $\angle CAD =$ the $\angle CED$, in this segment.

\therefore the sum of the $\angle^s BAC, CAD =$ the sum of the $\angle^s BEC, CED$;

that is, the $\angle BAD =$ the $\angle BED$. Q. E. D.

EXERCISES.

1. P is any point on the arc of a segment of which AB is the chord. Shew that the sum of the angles PAB, PBA is constant.

2. PQ and RS are two chords of a circle intersecting at X : prove that the triangles PXS, RXQ are equiangular.

3. Two circles intersect at A and B ; and through A any straight line PAQ is drawn terminated by the circumferences: shew that PQ subtends a constant angle at B .


4. Two circles intersect at A and B ; and through A any two straight lines PAQ, XAY are drawn terminated by the circumferences: shew that the arcs PX, QY subtend equal angles at B .

5. P is any point on the arc of a segment whose chord is AB : and the angles PAB, PBA are bisected by straight lines which intersect at O . Find the locus of the point O .

NOTE. If the extension of Proposition 20, given in the note on page 185, is adopted, a separate treatment of the second case of the present proposition is unnecessary.

For, as in Case I.,

the reflex $\angle BFD = \text{twice the } \angle BAD$; III. 20.
also the reflex $\angle BFD = \text{twice the } \angle BED$;
 \therefore the $\angle BAD = \text{the } \angle BED$.



The converse of Proposition 21 is very important. For the construction used in its proof, viz. To describe a circle about a given triangle, the student is referred to Book IV, Proposition 5. [Or see Theorems and Examples on Book I. Page 103, No. 1.]

CONVERSE OF PROPOSITION 21.

Equal angles standing on the same base, and on the same side of it, have their vertices on an arc of a circle, of which the given base is the chord.

Let $\angle BAC, \angle BDC$ be two equal angles standing on the same base BC :

then shall the vertices A and D lie upon a segment of a circle having BC as its chord.

Describe a circle about the $\triangle BAC$: IV. 5.
then this circle shall pass through D .

For, if not, it must cut BD , or BD produced, at some other point E .

Join EC .

Then the $\angle BAC = \text{the } \angle BEC$, in the same segment: III. 21.

but the $\angle BAC = \text{the } \angle BDC$, by hypothesis;

\therefore the $\angle BEC = \text{the } \angle BDC$;

that is, an ext. angle of a triangle = an int. opp. angle;
which is impossible. I. 16.


\therefore the circle which passes through B, A, C , cannot pass otherwise than through D .

That is, the vertices A and D are on an arc of a circle of which the chord is BC . Q. E. D.

The following corollary is important.


All triangles drawn on the same base, and with equal vertical angles, have their vertices on an arc of a circle, of which the given base is the chord.

OR, *The locus of the vertices of triangles drawn on the same base with equal vertical angles is an arc of a circle.*



PROPOSITION 22. THEOREM.

The opposite angles of any quadrilateral inscribed in a circle are together equal to two right angles.



Let $ABCD$ be a quadrilateral inscribed in the $\odot ABC$;
then shall, (i) the $\angle^s ADC, ABC$ together = two rt. angles;
(ii) the $\angle^s BAD, BCD$ together = two rt. angles.

Join AC, BD .

Then the $\angle ADB$ = the $\angle ACB$, in the segment $ADCB$; III. 21.
also the $\angle CDB$ = the $\angle CAB$, in the segment $CDAB$.

\therefore the $\angle ADC$ = the sum of the $\angle^s ACB, CAB$.

To each of these equals add the $\angle ABC$:

then the two $\angle^s ADC, ABC$ together = the three $\angle^s ACB, CAB, ABC$.

But the $\angle^s ACB, CAB, ABC$, being the angles of a triangle, together = two rt. angles. I. 32.

\therefore the $\angle^s ADC, ABC$ together = two rt. angles.

Similarly it may be shewn that

the $\angle^s BAD, BCD$ together = two rt. angles.

Q. E. D.

6

EXERCISES.

1. If a circle can be described about a parallelogram, the parallelogram must be rectangular.

2. ABC is an isosceles triangle, and XY is drawn parallel to the base BC : shew that the four points B, C, X, Y lie on a circle.

3. If one side of a quadrilateral inscribed in a circle is produced, the exterior angle is equal to the opposite interior angle of the quadrilateral.

PROPOSITION 22. [Alternative Proof.]

Let $ABCD$ be a quadrilateral inscribed in the $\odot ABC$;
then shall the $\angle^s ADC, ABC$ together = two rt. angles.

Join FA, FC .


Then the $\angle AFC$ at the centre = twice the $\angle ADC$ at the C^e , standing on the same arc ABC . III. 20.

Also the reflex angle AFC at the centre = twice the $\angle ABC$ at the C^e , standing on the same arc ADC . III. 20.

Hence the $\angle^s ADC, ABC$ are together half the sum of the $\angle AFC$ and the reflex angle AFC ;

but these make up four rt. angles: I. 15. Cor. 2.

\therefore the $\angle^s ADC, ABC$ together = two rt. angles. Q. E. D.



DEFINITION. Four or more points through which a circle may be described are said to be **concyelic**.

CONVERSE OF PROPOSITION 22.

If a pair of opposite angles of a quadrilateral are together equal to two right angles, its vertices are concyclic.

Let $ABCD$ be a quadrilateral, in which the opposite angles at B and D together = two rt. angles;

then shall the four points A, B, C, D be concyclic.

Through the three points A, B, C describe a circle: IV. 5.

then this circle must pass through D .

For, if not, it will cut AD , or AD produced, at some other point E .

Join EC .

Then since the quadrilateral $ABCE$ is inscribed in a circle,

\therefore the $\angle^s ABC, AEC$ together = two rt. angles. III. 22.

But the $\angle^s ABC, ADC$ together = two rt. angles; Hyp.

hence the $\angle^s ABC, AEC$ = the $\angle^s ABC, ADC$.


Take from these equals the $\angle ABC$;

then the $\angle AEC$ = the $\angle ADC$;

that is, an ext. angle of a triangle = an int. opp. angle;
which is impossible. I. 16.

\therefore the circle which passes through A, B, C cannot pass otherwise than through D ;


that is the four vertices A, B, C, D are concyclic. Q. E. D.



DEFINITION. Similar segments of circles are those which contain equal angles.

PROPOSITION 23. THEOREM.

On the same chord and on the same side of it, there cannot be two similar segments of circles, not coinciding with one another.



If possible, on the same chord AB , and on the same side of it, let there be two similar segments of circles ACB , ADB , not coinciding with one another.

Then since the arcs ADB , ACB intersect at A and B ,

\therefore they cannot cut one another again; III. 10.

\therefore one segment falls within the other.

In the outer arc take any point D ;

join AD , cutting the inner arc at C ;

join CB , DB .

Then because the segments are similar,

\therefore the $\angle ACB =$ the $\angle ADB$; III. Def.

that is, an ext. angle of a triangle = an int. opp. angle;

which is impossible. I. 16.

Hence the two similar segments ACB , ADB , on the same chord AB and on the same side of it, must coincide.

Q. E. D.

EXERCISES ON PROPOSITION 22.

1. The straight lines which bisect any angle of a quadrilateral figure inscribed in a circle and the opposite exterior angle, meet on the circumference.

2. A triangle is inscribed in a circle: shew that the sum of the angles in the three segments exterior to the triangle is equal to four right angles.

3. Divide a circle into two segments, so that the angle contained by the one shall be double of the angle contained by the other.

PROPOSITION 24. THEOREM.

Similar segments of circles on equal chords are equal to one another.

B

D

Let AEB and CFD be similar segments on equal chords AB, CD:

then shall the segment ABE = the segment CDF.

For if the segment ABE be applied to the segment CDF, so that A falls on C, and AB falls along CD;

then since AB = CD,

\therefore B must coincide with D.

\therefore the segment AEB must coincide with the segment CFD; for if not, on the same chord and on the same side of it there would be two similar segments of circles, not coinciding with one another: which is impossible. III. 23.

\therefore the segment AEB = the segment CDF. Q. E. D.

EXERCISES.

1. Of two segments standing on the same chord, the greater segment contains the smaller angle.

2. A segment of a circle stands on a chord AB, and P is any point on the same side of AB as the segment: shew that the angle APB is greater or less than the angle in the segment, according as P is within or without the segment.


3. P, Q, R are the middle points of the sides of a triangle, and X is the foot of the perpendicular let fall from one vertex on the opposite side: shew that the four points P, Q, R, X are concyclic.

[See page 96, Ex. 2; also page 100, Ex. 2.]

4. Use the preceding exercise to shew that the middle points of the sides of a triangle and the feet of the perpendiculars let fall from the vertices on the opposite sides, are concyclic.

PROPOSITION 25. PROBLEM*.

An arc of a circle being given, to describe the whole circumference of which the given arc is a part.



Let ABC be an arc of a circle;
it is required to describe the whole \cup^{ce} of which the arc ABC is a part.

In the given arc take any three points A, B, C .

Join AB, BC .

- Draw DF bisecting AB at rt. angles, I. 10. 11.
and draw EF bisecting BC at rt. angles.

Then because DF bisects the chord AB at rt. angles,

\therefore the centre of the circle lies in DF . III. 1. *Cor*

Again, because EF bisects the chord BC at rt. angles,

\therefore the centre of the circle lies in EF . III. 1. *Cor*.

\therefore the centre of the circle is F , the only point common to DF, EF .

Hence the \cup^{ce} of a circle described from centre F , with radius FA , is that of which the given arc is a part. Q. E. F.

* NOTE. Euclid gave this proposition a somewhat different form, as follows:

A segment of a circle being given, to describe the circle of which it is a segment.

Let ABC be the given segment standing on the chord AC .

Draw DB , bisecting AC at rt. angles. I. 10.

Join AB .


At A , in BA , make the $\angle BAE$ equal to the $\angle ABD$. I. 23.

Let AE meet BD , or BD produced, at E .

Then E shall be the centre of the required circle.


[Join EC ; and prove (i) $EA = EC$; I. 4.

(ii) $EA = EB$. I. 6.]



PROPOSITION 26. THEOREM.

In equal circles the arcs which subtend equal angles, whether at the centres or at the circumferences, shall be equal.



Let ABC, DEF be equal circles and let the $\angle^s BGC, EHF$, at the centres be equal, and consequently the $\angle^s BAC, EDF$ at the \odot^s equal: III. 20.

then shall the arc BKC = the arc ELF .

Join BC, EF .

Then because the $\odot^s ABC, DEF$ are equal,
 \therefore their radii are equal.

Hence in the $\triangle^s BGC, EHF$,

Because $\left\{ \begin{array}{l} BG = EH, \\ \text{and } GC = HF, \\ \text{and the } \angle BGC = \text{the } \angle EHF; \end{array} \right. \quad \begin{array}{l} \text{Hyp.} \\ \text{I. 4.} \end{array}$
 $\therefore BC = EF$.

Again, because the $\angle BAC = \text{the } \angle EDF$, Hyp.
 \therefore the segment BAC is similar to the segment EDF ;
 III. Def. 15.

and they are on equal chords BC, EF ;

\therefore the segment $BAC = \text{the segment } EDF$. III. 24.

But the whole $\odot ABC = \text{the whole } \odot DEF$;


\therefore the remaining segment $BKC = \text{the remaining segment } ELF$,
 \therefore the arc $BKC = \text{the arc } ELF$.

Q. E. D.

[For an Alternative Proof and Exercises see pp. 197, 198.]

PROPOSITION 27. THEOREM.

In equal circles the angles, whether at the centres or the circumferences, which stand on equal arcs, shall be equal.



Let ABC , DEF be equal circles,
and let the arc BC = the arc EF ;
then shall the $\angle BGC$ = the $\angle EHF$, at the centres ;
and also the $\angle BAC$ = the $\angle EDF$, at the \odot^{cs} .
If the \angle^s BGC , EHF are not equal, one must be the greater.

If possible, let the $\angle BGC$ be the greater.
At G , in BG , make the $\angle BGK$ equal to the $\angle EHF$. I. 23.

Then because in the equal \odot^{cs} ABC , DEF ,
the $\angle BGK$ = the $\angle EHF$, at the centres ; *Constr.*
 \therefore the arc BK = the arc EF . III. 26.

But the arc BC = the arc EF , *Hyp.*
 \therefore the arc BK = the arc BC ,
 \therefore a part equal to the whole, which is impossible.

\therefore the $\angle BGC$ is not unequal to the $\angle EHF$;
that is, the $\angle BGC$ = the $\angle EHF$.


And since the $\angle BAC$ at the \odot^{ce} is half the $\angle BGC$ at the centre, III. 20,

and likewise the $\angle EDF$ is half the $\angle EHF$,
 \therefore the $\angle BAC$ = the $\angle EDF$. Q. E. D.

[For Exercises see pp. 197, 198.]

PROPOSITION 28. THEOREM.

In equal circles the arcs, which are cut off by equal chords, shall be equal, the major arc equal to the major arc, and the minor to the minor.



Let ABC , DEF be two equal circles,
and let the chord BC = the chord EF :
then shall the major arc BAC = the major arc EDF ,
and the minor arc BGC = the minor arc EHF .

Find K and L the centres of the \odot s ABC , DEF : III. 1.
and join BK , KC , EL , LF .

Then because the \odot s ABC , DEF are equal,
 \therefore their radii are equal.

Hence in the \triangle s BKC , ELF ,


Because $\begin{cases} BK = EL, \\ KC = LF, \\ \text{and } BC = EF; \end{cases}$ Hyp.
 \therefore the $\angle BKC$ = the $\angle ELF$; I. 8.
 \therefore the arc BGC = the arc EHF ; III. 26.
and these are the minor arcs.

But the whole $\odot^{\text{cc}} ABGC$ = the whole $\odot^{\text{cc}} DEHF$; Hyp.
 \therefore the remaining arc BAC = the remaining arc EDF :
and these are the major arcs. Q.E.D.

[For Exercises see pp. 197, 198.]

PROPOSITION 29. THEOREM.

In equal circles the chords, which cut off equal arcs, shall be equal.



Let ABC , DEF be equal circles,
and let the arc BGC = the arc EHF :
then shall the chord BC = the chord EF .

Find K , L the centres of the circles. III. 1.

Join BK , KC , EL , LF .

Then in the equal \odot^s ABC , DEF ,
because the arc BGC = the arc EHF ,

\therefore the $\angle BKC$ = the $\angle ELF$. III. 27.

Hence in the \triangle^s BKC , ELF ,

Because $\left\{ \begin{array}{l} BK = EL, \text{ being radii of equal circles;} \\ KC = LF, \text{ for the same reason,} \\ \text{and the } \angle BKC = \text{the } \angle ELF; \end{array} \right. \quad \text{Proved.}$

$\therefore BC = EF$. I. 4.

Q. E. D.

EXERCISES

ON PROPOSITIONS 26, 27.

1. *If two chords of a circle are parallel, they intercept equal arcs.*
2. *The straight lines, which join the extremities of two equal arcs of a circle towards the same parts, are parallel.*
3. *In a circle, or in equal circles, sectors are equal if their angles at the centres are equal.*

4. If two chords of a circle intersect at right angles, the opposite arcs are together equal to a semicircumference.

5. If two chords intersect within a circle, they form an angle equal to that subtended at the circumference by the sum of the arcs they cut off.

6. If two chords intersect without a circle, they form an angle equal to that subtended at the circumference by the difference of the arcs they cut off.

7. If AB is a fixed chord of a circle, and P any point on one of the arcs cut off by it, then the bisector of the angle APB cuts the conjugate arc in the same point, whatever be the position of P .

8. Two circles intersect at A and B ; and through these points straight lines are drawn from any point P on the circumference of one of the circles: shew that when produced they intercept on the other circumference an arc which is constant for all positions of P .

9. A triangle ABC is inscribed in a circle, and the bisectors of the angles meet the circumference at X, Y, Z . Find each angle of the triangle XYZ in terms of those of the original triangle.

ON PROPOSITIONS 28, 29.

10. The straight lines which join the extremities of parallel chords in a circle (i) towards the same parts, (ii) towards opposite parts, are equal.

11. Through A , a point of intersection of two equal circles two straight lines PAQ, XAY are drawn: shew that the chord PX is equal to the chord QY .

12. Through the points of intersection of two circles two parallel straight lines are drawn terminated by the circumferences: shew that the straight lines which join their extremities towards the same parts are equal.

13. Two equal circles intersect at A and B ; and through A any straight line PAQ is drawn terminated by the circumferences: shew that $BP = BQ$.

14. ABC is an isosceles triangle inscribed in a circle, and the bisectors of the base angles meet the circumference at X and Y . Shew that the figure $BXAYC$ must have four of its sides equal.


What relation must subsist among the angles of the triangle ABC , in order that the figure $BXAYC$ may be equilateral?

NOTE. We have given Euclid's demonstrations of Propositions 26, 27, 28, 29; but it should be noticed that all these propositions also admit of direct proof by the method of *superposition*.

• To illustrate this method we will apply it to Proposition 26.

PROPOSITION 26. [Alternative Proof.]

In equal circles, the arcs which subtend equal angles whether at the centres or circumferences, shall be equal.



• Let ABC , DEF be equal circles, and let the \angle 's BGC , EHF at the centres be equal, and consequently the \angle 's BAC , EDF at the cs equal: III. 20.

then shall the arc BKC = the arc ELF .

For if the $\odot ABC$ be applied to the $\odot DEF$, so that the centre G may fall on the centre H ,

then because the circles are equal, *Hyp.*

\therefore their cs must coincide,

hence by revolving the upper circle about its centre, the lower circle remaining fixed,

B may be made to coincide with E ,

and consequently GB with HE .

And because the \angle BGC , the \angle EHF ,

$\therefore GC$ must coincide with HF :

and since $GC = HF$,

$\therefore C$ must fall on F . *Hyp.*

Now B coinciding with E , and C with F ,

and the cs of the $\odot ABC$ with the cs of the $\odot DEF$,

\therefore the arc BKC must coincide with the arc ELF .

\therefore the arc $BKC =$ the arc ELF .

Q.E.D.

19. A segment of a circle is described on the chord AB : find a point C on its arc so that the sum of AC , BC may be a maximum.

20. Of all triangles that can be inscribed in a circle that which has the greatest perimeter is equilateral.

21. Of all triangles that can be inscribed in a given circle that which has the greatest area is equilateral.

22. Of all triangles that can be inscribed in a given triangle that which has the least perimeter is the triangle formed by joining the feet of the perpendiculars drawn from the vertices on opposite sides.

23. Of all rectangles of given area, the square has the least perimeter.

24. Describe the triangle of maximum area, having its angles equal to those of a given triangle, and its sides passing through three given points.

VI. HARDER MISCELLANEOUS EXAMPLES

1. AB is a diameter of a given circle; and AC , BD , two chords on the same side of AB , intersect at E : shew that the circle which passes through D , E , C cuts the given circle orthogonally.

2. Two circles whose centres are C and D intersect at A and B , and a straight line PAQ is drawn through A and terminated by the circumferences: prove that

(i) the angle PBQ = the angle CAD

(ii) the angle BPC = the angle BQD .

3. Two chords AB , CD of a circle whose centre is O intersect at right angles at P : shew that

(i) $PA^2 + PB^2 + PC^2 + PD^2 = 4$ (radius) 2 .

(ii) $AB^2 + CD^2 + 4OP^2 = 8$ (radius) 2 .

4. Two parallel tangents to a circle intercept on any third tangent a portion which is so divided at its point of contact that the rectangles contained by its two parts is equal to the square on the radius.


5. Two equal circles move between two straight lines placed at right angles, so that each straight line is touched by one circle, and the two circles touch one another: find the locus of the point of contact.

6. AB is a given diameter of a circle, and CD is any parallel chord: if any point X in AB is joined to the extremities of CD , shew that

$$XC^2 + XD^2 = XA^2 + XB^2.$$

PROPOSITION 31. THEOREM.

*The angle in a semicircle is a right angle ;
the angle in a segment greater than a semicircle is less
than a right angle ;
and the angle in a segment less than a semicircle is
greater than a right angle.*



Let ABCD be a circle, of which BC is a diameter, and E the centre; and let AC be a chord dividing the circle into the segments ABC, ADC, of which the segment ABC is greater, and the segment ADC less than a semicircle:
then (i) the angle in the semicircle BAC shall be a rt. angle;
(ii) the angle in the segment ABC shall be less than a rt. angle;
(iii) the angle in the segment ADC shall be greater than a rt. angle.

In the arc ADC take any point D;
Join BA, AD, DC, AE; and produce BA to F.

(i) Then because $EA = EB$, III. Def. 1.
 \therefore the $\angle EAB =$ the $\angle EBA$. I. 5.

And because $EA = EC$,
 \therefore the $\angle EAC =$ the $\angle ECA$.

\therefore the whole $\angle BAC =$ the sum of the $\angle^s EBA, ECA$;
but the ext. $\angle FAC =$ the sum of the two int. $\angle^s CBA, BCA$;
 \therefore the $\angle BAC =$ the $\angle FAC$;

\therefore these angles, being adjacent, are rt. angles.
 \therefore the $\angle BAC$, in the semicircle BAC, is a rt. angle.

(ii) In the $\triangle ABC$, because the two \angle^s ABC , BAC are together less than two rt. angles; I. 17.

and of these, the $\angle BAC$ is a rt. angle; *Proved.*
 \therefore the $\angle ABC$, which is the angle in the segment ABC , is less than a rt. angle.

(iii) Because $ABCD$ is a quadrilateral inscribed in the $\odot ABC$,

\therefore the \angle^s ABC , ADC together = two rt. angles; III. 22.
 and of these, the $\angle ABC$ is less than a rt. angle: *Proved.*
 \therefore the $\angle ADC$, which is the angle in the segment ADC , is greater than a rt. angle. Q. E. D.

EXERCISES.

1. A circle described on the hypotenuse of a right-angled triangle as diameter, passes through the opposite angular point.

2. A system of right-angled triangles is described upon a given straight line as hypotenuse: find the locus of the opposite angular points.

3. A straight rod of given length slides between two straight rulers placed at right angles to one another: find the locus of its middle point.

4. Two circles intersect at A and B ; and through A two diameters AP , AQ are drawn, one in each circle: shew that the points P , B , Q are collinear. [See Def. p. 102.]

5. A circle is described on one of the equal sides of an isosceles triangle as diameter. Shew that it passes through the middle point of the base.

6. Of two circles which have internal contact, the diameter of the inner is equal to the radius of the outer. Shew that any chord of the outer circle, drawn from the point of contact, is bisected by the circumference of the inner circle.

7. Circles described on any two sides of a triangle as diameters intersect on the third side, or the third side produced.

8. Find the locus of the middle points of chords of a circle drawn through a fixed point.

Distinguish between the cases when the given point is within, on, or without the circumference.

9. Describe a square equal to the difference of two given squares.


10. Through one of the points of intersection of two circles draw a chord of one circle which shall be bisected by the other.

11. On a given straight line as base a system of equilateral four-sided figures is described: find the locus of the intersection of their diagonals.

NOTE 1. The extension of Proposition 20 to *straight* and *reflex* angles furnishes a simple alternative proof of the first theorem contained in Proposition 31, viz.

- The angle in a semicircle is a right angle.

For, in the adjoining figure, the angle at the centre, standing on the arc BHC , is double the angle at the \circ^o , standing on the same arc.



Now the angle at the centre is the *straight angle* BEC ;

\therefore the $\angle BAC$ is half of the *straight angle* BEC ;

and a straight angle two rt. angles;

- \therefore the $\angle BAC$ — one half of two rt. angles,
— one rt. angle.

Q.E.D.

NOTE 2. From Proposition 31 we may derive a simple practical solution of Proposition 17, namely,

- To draw a tangent to a circle from a given external point.

Let BCD be the given circle, and A the given external point:

it is required to draw from A a tangent to the $\circ BCD$.

Find E , the centre of the circle, and join AE .

On AE describe the semicircle ABE , to cut the given circle at B .

Join AB .

Then AB shall be a tangent to the $\circ BCD$.


For the $\angle ABE$, being in a semicircle, is a rt. angle. III. 31.

$\therefore AB$ is drawn at rt. angles to the radius EB , from its extremity B ;

$\therefore AB$ is a tangent to the circle. III. 16.


Q.E.F.

Since the semicircle might be described on either side of AE , it is clear that there will be a second solution of the problem, as shewn by the dotted lines of the figure.



• PROPOSITION 32. THEOREM.

If a straight line touch a circle, and from the point of contact a chord be drawn, the angles which this chord makes with the tangent shall be equal to the angles in the alternate segments of the circle.



Let EF touch the given $\odot ABC$ at B , and let BD be a chord drawn from B , the point of contact:

then shall (i) the $\angle DBF$ the angle in the alternate segment BAD :

(ii) the $\angle DBE$ the angle in the alternate segment BCD .

From B draw BA perp. to EF . 1. 11.

Take any point C in the arc BD ,
and join AD , DC , CB .

(i) Then because BA is drawn perp. to the tangent EF , at its point of contact B ,

$\therefore BA$ passes through the centre of the circle: III. 19.


\therefore the $\angle ADB$, being in a semicircle, is a rt. angle: III. 31.

\therefore in the $\angle ABD$, the other $\angle^s ABD$, BAD together = a rt. angle; I. 32.

that is, the $\angle^s ABD$, BAD together = the $\angle ABF$.

From these equals take the common $\angle ABD$;

\therefore the $\angle DBF$ = the $\angle BAD$, which is in the alternate segment.



(ii) Because ABCD is a quadrilateral inscribed in a circle,

\therefore the \angle^s BCD, BAD together = two rt. angles: III. 22.

but the \angle^s DBE, DBF together = two rt. angles; I. 13.

\therefore the \angle^s DBE, DBF together = the \angle^s BCD, BAD:

and of these the \angle DBF = the \angle BAD; *Proved.*


\therefore the \angle DBE = the \angle DCB, which is in the alternate segment, Q. E. D.

EXERCISES.

1. State and prove the converse of this proposition.
2. Use this Proposition to shew that the tangents drawn to a circle from an external point are equal.
3. If two circles touch one another, any straight line drawn through the point of contact cuts off similar segments.
Prove this for (i) internal, (ii) external contact.
4. If two circles touch one another, and from A, the point of contact, two chords APQ, AXY are drawn: then PX and QY are parallel.
Prove this for (i) internal, (ii) external contact.
5. Two circles intersect at the points A, B: and one of them passes through O, the centre of the other: prove that OA bisects the angle between the common chord and the tangent to the first circle at A.
6. Two circles intersect at A and B; and through P, any point on the circumference of one of them, straight lines PAC, PBD are drawn to cut the other circle at C and D: shew that CD is parallel to the tangent at P.
7. If from the point of contact of a tangent to a circle, a chord be drawn, the perpendiculars dropped on the tangent and chord from the middle point of either arc cut off by the chord are equal.

PROPOSITION 33. PROBLEM.

On a given straight line to describe a segment of a circle which shall contain an angle equal to a given angle.



Let AB be the given st. line, and C the given angle: it is required to describe on AB a segment of a circle which shall contain an angle equal to C .

At A in BA , make the $\angle BAD$ equal to the $\angle C$. I. 23.

From A draw AE at rt. angles to AD . I. 11.

Bisect AB at F ; I. 10.

and from F draw FG at rt. angles to AB , cutting AE at G .

Join GB .

Then in the \triangle^s AFG , BFG .

$AF = BF$, *Constr.*

Because FG is common,

(and the $\angle AFG = \angle BFG$, being rt. angles,
 $\therefore GA = GB$: I. 4.

\therefore the circle described from centre G , with radius GA , will pass through B .

Describe this circle, and call it ABH :

then the segment AHB shall contain an angle equal to C .

Because AD is drawn at rt. angles to the radius GA from its extremity A ,

$\therefore AD$ is a tangent to the circle: III. 16.

and from A , its point of contact, a chord AB is drawn;


\therefore the $\angle BAD =$ the angle in the alt. segment AHB . III. 32.

But the $\angle BAD = \angle C$: *Constr.*

\therefore the angle in the segment $AHB = \angle C$.

$\therefore AHB$ is the segment required. Q. E. F.

NOTE. In the particular case when the given angle C is a rt. angle, the segment required will be the semicircle described on the given st line AB ; for the angle in a semicircle is a rt. angle. III. 31.



EXERCISES.

[The following exercises depend on the corollary to Proposition 21 given on page 187, namely

The locus of the vertices of triangles which stand on the same base and have a given vertical angle, is the arc of the segment standing on this base, and containing an angle equal to the given angle.

Exercises 1 and 2 afford good illustrations of the method of finding required points by the *Intersection of Loci*. See page 117.]

1. Describe a triangle on a given base, having a given vertical angle, and having its vertex on a given straight line.

2. Construct a triangle, having given the base, the vertical angle and

- (i) one other side,
- (ii) the altitude,
- (iii) the length of the median which bisects the base,
- (iv) the point at which the perpendicular from the vertex meets the base.

3. Construct a triangle having given the base, the vertical angle, and the point at which the base is cut by the bisector of the vertical angle.

[Let AB be the base, X the given point in it, and K the given angle. On AB describe a segment of a circle containing an angle equal to K ; complete the co^e by drawing the arc APB . Bisect the arc APB at P ; join PX , and produce it to meet the co^e at C . Then ABC shall be the required triangle.]


4. Construct a triangle having given the base, the vertical angle, and the sum of the remaining sides.

[Let AB be the given base, K the given angle, and H the given line equal to the sum of the sides. On AB describe a segment containing an angle equal to K , also another segment containing an angle equal to half the $\angle K$. From centre A , with radius H , describe a circle cutting the last drawn segment at X and Y . Join AX (or AY) cutting the first segment at C . Then ABC shall be the required triangle.]

5. Construct a triangle having given the base, the vertical angle, and the difference of the remaining sides.

PROPOSITION 31. PROBLEM.

From a given circle to cut off a segment which shall contain an angle equal to a given angle.



Let ABC be the given circle, and D the given angle: it is required to cut off from the $\odot ABC$ a segment which shall contain an angle equal to D .

Take any point B on the \odot ,
and at B draw the tangent EBF . III. 17.

At B , in FB , make the $\angle FBC$ equal to the $\angle D$. I. 23.
Then the segment BAC shall contain an angle equal to D .

Because EF is a tangent to the circle, and from B , its point of contact, a chord BC is drawn,

\therefore the $\angle FBC$ = the angle in the alternate segment BAC . III. 32.

But the $\angle FBC$ = the $\angle D$; Constr.

\therefore the angle in the segment BAC = the $\angle D$.

Hence from the given $\odot ABC$ a segment BAC has been cut off, containing an angle equal to D . Q. E. F.


EXERCISES.

1. The chord of a given segment of a circle is produced to a fixed point: on this straight line so produced draw a segment of a circle similar to the given segment.

2. Through a given point without a circle draw a straight line that will cut off a segment capable of containing an angle equal to a given angle.

PROPOSITION 35. THEOREM.

If two chords of a circle cut one another, the rectangle contained by the segments of one shall be equal to the rectangle contained by the segments of the other.



Let AB, CD, two chords of the \odot ACBD, cut one another at E:

then shall the rect. AE, EB = the rect. CE, ED.

Find F the centre of the \odot ACB: III. 1.

From F draw FG, FH perp. respectively to AB, CD. I. 12.

Join FA, FE, FD.

Then because FG is drawn from the centre F perp. to AB,
 \therefore AB is bisected at G. III. 3.

For a similar reason CD is bisected at H.

Again, because AB is divided equally at G, and unequally at E,
 \therefore the rect. AE, EB with the sq. on EG = the sq. on AG. II. 5.

To each of these equals add the sq. on GF;
 then the rect. AE, EB with the sqq. on EG, GF = the sum of
 the sqq. on AG, GF.

But the sqq. on EG, GF = the sq. on FE: I. 17.

and the sqq. on AG, GF = the sq. on AF:

for the angles at G are rt. angles.

\therefore the rect. AE, EB with the sq. on FE = the sq. on AF.

Similarly it may be shewn that

the rect. CE, ED with the sq. on FE = the sq. on FD.

But the sq. on AF = the sq. on FD; for AF = FD.

\therefore the rect. AE, EB with the sq. on FE = the rect. CE, ED
 with the sq. on FE.

From these equals take the sq. on FE:

then the rect. AE, EB = the rect. CE, ED. Q. E. D.

COROLLARY. *If through a fixed point within a circle any number of chords are drawn, the rectangles contained by their segments are all equal.*

NOTE. The following special cases of this proposition deserve notice.

- (i) when the given chords both pass through the centre:
- (ii) when one chord passes through the centre, and cuts the other at right angles:
- (iii) when one chord passes through the centre, and cuts the other obliquely.


In each of these cases the general proof requires some modification, which may be left as an exercise to the student.

EXERCISES.

1. Two straight lines AB , CD intersect at E , so that the rectangle AE , EB is equal to the rectangle CE , ED : shew that the four points A , B , C , D are concyclic.
2. The rectangle contained by the segments of any chord drawn through a given point within a circle is equal to the square on half the shortest chord which may be drawn through that point.
3. ABC is a triangle right-angled at C ; and from C a perpendicular CD is drawn to the hypotenuse: shew that the square on CD is equal to the rectangle AD , DB .
4. ABC is a triangle; and AP , BQ the perpendiculars dropped from A and B on the opposite sides, intersect at O : shew that the rectangle AO , OP is equal to the rectangle BO , OQ .
5. Two circles intersect at A and B , and through any point in AB their common chord two chords are drawn, one in each circle; shew that their four extremities are concyclic.
6. A and B are two points within a circle such that the rectangle contained by the segments of any chord drawn through A is equal to the rectangle contained by the segments of any chord through B : shew that A and B are equidistant from the centre.
7. If through E , a point without a circle, two secants EAB , ECD are drawn; shew that the rectangle EA , EB is equal to the rectangle EC , ED .
[Proceed as in III. 35, using II. 6.]
8. Through A , a point of intersection of two circles, two straight lines CAE , DAF are drawn, each passing through a centre and terminated by the circumferences: shew that the rectangle CA , AE is equal to the rectangle DA , AF .

PROPOSITION 36. THEOREM.

If from any point without a circle a tangent and a secant be drawn, then the rectangle contained by the whole secant and the part of it without the circle shall be equal to the square on the tangent.



Let ABC be a circle; and from D a point without it, let there be drawn the secant DCA , and the tangent DB :

then the rect. DA, DC shall be equal to the sq. on DB .

Find E , the centre of the $\odot ABC$: III. 1

and from E , draw EF perp. to AD . I. 12.

Join EB, EC, ED .

Then because EF , passing through the centre, is perp. to the chord AC ,

$\therefore AC$ is bisected at F . III. 3.

And since AC is bisected at F and produced to D ,

\therefore the rect. DA, DC with the sq. on FC = the sq. on FD . II. 6.

To each of these equals add the sq. on EF :

then the rect. DA, DC with the sqq. on EF, FC = the sqq. on EF, FD .

But the sqq. on EF, FC = the sq. on EC ; for EFC is a rt. angle;
= the sq. on EB .

And the sqq. on EF, FD = the sq. on ED ; for EFD is a rt. angle;
the sq. on EB, BD ; for EBD is a
rt. angle. III. 18.

\therefore the rect. DA, DC with the sq. on EB = the sq. on EB, BD .

From these equals take the sq. on EB :


then the rect. DA, DC = the sq. on DB . Q. E. D.

NOTE. This proof may easily be adapted to the case when the secant passes through the centre of the circle.

COROLLARY. *If from a given point without a circle any number of secants are drawn, the rectangles contained by the whole secants and the parts of them without the circle are all equal; for each of these rectangles is equal to the square on the tangent drawn from the given point to the circle.*

For instance, in the adjoining figure, each of the rectangles PB, PA and PD, PC and PF, PE is equal to the square on the tangent PQ :

∴ the rect. PB, PA
 the rect. PD, PC
 the rect. PF, PE .



NOTE. Remembering that the segments into which the chord AB is divided at P , are the lines PA, PB , (see Part I. page 131) we are enabled to include the corollaries of Propositions 35 and 36 in a single enunciation.

If any number of chords of a circle are drawn through a given point, within or without a circle, the rectangles contained by the segments of the chords are equal.

EXERCISES.

1. Use this proposition to shew that tangents drawn to a circle from an external point are equal.

2. If two circles intersect, tangents drawn to them from any point in their common chord produced are equal.

3. If two circles intersect at A and B , and PQ is a tangent to both circles; shew that AB produced bisects PQ .


4. If P is any point on the straight line AB produced, shew that the tangents drawn from P to all circles which pass through A and B are equal.

5. ABC is a triangle right-angled at C , and from any point P in AC , a perpendicular PQ is drawn to the hypotenuse; shew that the rectangle AC, AP is equal to the rectangle AB, AQ .

6. ABC is a triangle right-angled at C , and from C a perpendicular CD is drawn to the hypotenuse; shew that the rect. AB, AD is equal to the square on AC .

PROPOSITION 37. THEOREM.

If from a point without a circle there be drawn two straight lines, one of which cuts the circle, and the other meets it, and if the rectangle contained by the whole line which cuts the circle and the part of it without the circle be equal to the square on the line which meets the circle, then the line which meets the circle shall be a tangent to it.



Let ABC be a circle; and from D , a point without it, let there be drawn two st. lines DCA and DB , of which DCA cuts the circle at C and A , and DB meets it; and let the rect. DA, DC = the sq. on DB :

then shall DB be a tangent to the circle.

From D draw DE to touch the $\odot ABC$: I. 17.

let E be the point of contact.

Find the centre F , and join FB, FD, FE . III. 1.

Then since DCA is a secant, and DE a tangent to the circle,

\therefore the rect. DA, DC = the sq. on DE , III. 36.

But, by hypothesis, the rect. DA, DC = the sq. on DB ;

\therefore the sq. on DE = the sq. on DB ,

$\therefore DE = DB$.

Hence in the $\triangle^s DBF, DEF$.

Because $\begin{cases} DB = DE, \\ \text{and } BF = EF; \\ \text{and } DF \text{ is common;} \end{cases}$ Proved.
III. Def. 1.

\therefore the $\angle DBF$ = the $\angle DEF$. I. 8.

But DEF is a rt. angle; III. 18.

$\therefore DBF$ is also a rt. angle;

and since BF is a radius,

$\therefore DB$ touches the $\odot ABC$ at the point B .

Q. E. D.

NOTE ON THE METHOD OF LIMITS AS APPLIED TO TANGENCY.

Euclid defines a tangent to a circle as *a straight line which meets the circumference, but being produced, does not cut it* and from this definition he deduces the fundamental theorem that a tangent is perpendicular to the radius drawn to the point of contact. Prop. 16.

But this result may also be established by the Method of Limits, which regards the tangent as *the ultimate position of a secant when its two points of intersection with the circumference are brought into coincidence* [See Note on page 151]; and it may be shown that every theorem relating to the tangent may be derived from some more general proposition relating to the secant, by considering the ultimate case when the two points of intersection coincide.

1. *To prove by the Method of Limits that a tangent to a circle is at right angles to the radius drawn to the point of contact.*

Let ABD be a circle, whose centre is C ; and $PABQ$ a secant cutting the circle in A and B ; and let $P'AQ'$ be the limiting position of PQ when the point B is brought into coincidence with A ; then shall CA be perp. to $P'Q'$.


Bisect AB at E and join CE :

then CE is perp. to PQ . III. 3.

Now let the secant $PABQ$ change its position in such a way that while the point A remains fixed, the point B continually approaches A , and ultimately coincides with it;

then, *however near B approaches to A* , the st. line CE is always perp. to PQ , since it joins the centre to the middle point of the chord AB .

But in the limiting position, when B coincides with A , and the secant PQ becomes the tangent $P'Q'$, it is clear that the point E will also coincide with A ; and the perpendicular CE becomes the radius CA . Hence CA is perp. to the tangent $P'Q'$ at its point of contact A . Q. E. D.



NOTE. It follows from Proposition 2 that *a straight line cannot cut the circumference of a circle at more than two points*. Now when the two points in which a secant cuts a circle move towards coincidence, the secant ultimately becomes a tangent to the circle: we infer therefore that a tangent cannot meet a circle otherwise than at its point of contact. Thus Euclid's definition of a tangent may be deduced from that given by the Method of Limits.

2. By this Method Proposition 32 may be derived as a special case from Proposition 21.

For let A and B be two points on the \odot^{ce} of the $\odot ABC$;
and let BCA , BPA be any two angles in the segment $BCPA$:
then the $\angle BPA =$ the $\angle BCA$. III. 21.

Produce PA to Q .

Now let the point P continually approach the fixed point A , and ultimately coincide with it;

then, however near P may approach to A ,
the $\angle BPQ =$ the $\angle BCA$. III. 21.


But in the limiting position when P coincides with A ,

and the secant PAQ becomes the tangent AQ' ,

it is clear that BP will coincide with BA ,

and the $\angle BPQ$ becomes the $\angle BAQ'$.

Hence the $\angle BAQ' =$ the $\angle BCA$, in the alternate segment. Q. E. D.



The contact of circles may be treated in a similar manner by adopting the following definition.

DEFINITION. If one or other of two intersecting circles alters its position in such a way that the two points of intersection continually approach one another, and ultimately coincide; in the limiting position they are said to **touch** one another, and the point in which the two points of intersection ultimately coincide is called the **point of contact**.

EXAMPLES ON LIMITS.

1. Deduce Proposition 19 from the Corollary of Proposition 1 and Proposition 3.

2. Deduce Propositions 11 and 12 from Ex. 1, page 156.

3. Deduce Proposition 6 from Proposition 5.

4. Deduce Proposition 13 from Proposition 10.

5. Shew that a straight line cuts a circle in two different points, two coincident points, or not at all, according as its distance from the centre is less than, equal to, or greater than a radius.

6. Deduce Proposition 32 from Ex. 3, page 188.

7. Deduce Proposition 36 from Ex. 7, page 209.

8. The angle in a semi-circle is a right angle.

To what Theorem is this statement reduced, when the vertex of the right angle is brought into coincidence with an extremity of the diameter?

9. From Ex. 1, page 190, deduce the corresponding property of a triangle inscribed in a circle.


THEOREMS AND EXAMPLES, ON BOOK III.

1. ON THE CENTRE AND CHORDS OF A CIRCLE.

See Propositions 1, 3, 14, 15, 25.

1. *All circles which pass through a fixed point, and have their centres on a given straight line, pass also through a second fixed point.*

Let AB be the given st. line, and P the given point.



From P draw PR perp. to AB ;
and produce PR to P' , making RP' equal to PR .

Then all circles which pass through P , and have their centres on AB , shall pass also through P' .

For let C be the centre of *any one* of these circles.

Join CP , CP' .

Then in the Δ^s CRP , CRP'

Because $\left\{ \begin{array}{l} CR \text{ is common,} \\ \text{and } RP = RP', \\ \text{and the } \angle CRP = \text{the } \angle CRP', \text{ being rt. angles;} \end{array} \right.$ Constr.

$\therefore CP = CP'$; I. 4.

\therefore the circle whose centre is C , and which passes through P , must pass also through P' .

But C is the centre of *any* circle of the system;

\therefore all circles, which pass through P , and have their centres in AB , pass also through P' . Q. E. D.

2. *Describe a circle that shall pass through three given points not in the same straight line.*

3. Describe a circle that shall pass through two given points and have its centre in a given straight line. When is this impossible?

4. Describe a circle of given radius to pass through two given points. When is this impossible?

5. ABC is an isosceles triangle; and from the vertex A , as centre a circle is described cutting the base, or the base produced, at X and Y . Shew that $BX = CY$.

6. If two circles which intersect are cut by a straight line parallel to the common chord, shew that the parts of it intercepted between the circumferences are equal.

7. If two circles cut one another, any two straight lines drawn through a point of section, making equal angles with the common chord, and terminated by the circumferences, are equal. [Ex. 12, p. 156.]

8. If two circles cut one another, of all straight lines drawn through a point of section and terminated by the circumferences, the greatest is that which is parallel to the line joining the centres.

9. Two circles, whose centres are C and D , intersect at A , B , and through A a straight line PAQ is drawn terminated by the circumferences: if PC , QD intersect at X , shew that the angle PXQ is equal to the angle CAD .

10. Through a point of section of two circles which cut one another draw a straight line terminated by the circumferences and bisected at the point of section.

11. AB is a fixed diameter of a circle, whose centre is C ; and from P , any point on the circumference, PQ is drawn perpendicular to AB ; shew that the bisector of the angle CPQ always intersects the circle in one or other of two fixed points.

12. Circles are described on the sides of a quadrilateral as diameters: shew that the common chord of any two consecutive circles is parallel to the common chord of the other two. [Ex. 9, p. 97.]

13. Two equal circles touch one another externally, and through the point of contact two chords are drawn, one in each circle, at right angles to each other: shew that the straight line joining their other extremities is equal to the diameter of either circle.

14. Straight lines are drawn from a given external point to the circumference of a circle: find the locus of their middle points. [Ex. 3, p. 97.]

15. Two equal segments of circles are described on opposite sides of the same chord AB ; and through O , the middle point of AB , any straight line POQ is drawn, intersecting the arcs of the segments at P and Q : shew that $OP = OQ$.

II. ON THE TANGENT AND THE CONTACT OF CIRCLES.

See Propositions 11, 12, 16, 17, 18, 19.

1. All equal chords placed in a given circle, touch a fixed concentric circle.

2. If from an external point two tangents are drawn to a circle, the angle contained by them is double the angle contained by the chord of contact and the diameter drawn through one of the points of contact.

3. Two circles touch one another externally, and through the point of contact a straight line is drawn terminated by the circumferences: shew that the tangents at its extremities are parallel.

4. Two circles intersect, and through one point of section any straight line is drawn terminated by the circumferences: shew that the angle between the tangents at its extremities is equal to the angle between the tangents at the point of section.

5. Shew that two parallel tangents to a circle intercept on any third tangent a segment which subtends a right angle at the centre.

6. Two tangents are drawn to a given circle from a fixed external point A , and any third tangent cuts them produced at P and Q : shew that PQ subtends a constant angle at the centre of the circle.

7. In any quadrilateral circumscribed about a circle, the sum of one pair of opposite sides is equal to the sum of the other pair.

8. If the sum of one pair of opposite sides of a quadrilateral is equal to the sum of the other pair, shew that a circle may be inscribed in the figure.

[Bisect two adjacent angles of the figure, and so describe a circle to touch three of its sides. Then prove indirectly by means of the last exercise that this circle must also touch the fourth side.]

9. Two circles touch one another internally: shew that of all chords of the outer circle which touch the inner, the greatest is that which is perpendicular to the straight line joining the centres.

10. In any triangle, if a circle is described from the middle point of one side as centre and with a radius equal to half the sum of the other two sides, it will touch the circles described on these sides as diameters.

11. Through a given point, draw a straight line to cut a circle, so that the part intercepted by the circumference may be equal to a given straight line.

In order that the problem may be possible, between what limits must the given line lie, when the given point is (i) without the circle, (ii) within it?

12. A series of circles touch a given straight line at a given point: shew that the tangents to them at the points where they cut a given parallel straight line all touch a fixed circle, whose centre is the given point.

13. If two circles touch one another internally, and any third circle be described touching both; then the sum of the distances of the centre of this third circle from the centres of the two given circles is constant.

14. Find the locus of points such that the pairs of tangents drawn from them to a given circle contain a constant angle.

15. Find a point such that the tangents drawn from it to two given circles may be equal to two given straight lines. When is this impossible?

16. If three circles touch one another two and two; prove that the tangents drawn to them at the three points of contact are concurrent and equal.


THE COMMON TANGENTS TO TWO CIRCLES.

17. To draw a common tangent to two circles.

First, if the given circles are external to one another, or if they intersect.

Let A be the centre of the greater circle, and B the centre of the less.

From A , with radius equal to the diff^e of the radii of the given circles, describe a circle: and from B draw BC to touch the last drawn circle. Join AC , and produce it to meet the greater of the given circles at D .



Through B draw the radius BE par^l to AD , and in the same direction.

Join DE :

then DE shall be a common tangent to the two given circles.

For since AC -- the diff^e between AD and BE ,

Constr.

CD -- BE :

and CD is par^l to BE ;

Constr.

$\therefore DE$ is equal and par^l to CB .

1. 33.

But since BC is a tangent to the circle at C ,

\therefore the $\angle ACB$ is a rt. angle;

III. 18.

hence each of the angles at D and E is a rt. angle:

1. 29.

$\therefore DE$ is a tangent to both circles.

Q.E.F.

It follows from hypothesis that the point B is outside the circle used in the construction:

∴ two tangents such as BC may always be drawn to it from B ; hence *two* common tangents may always be drawn to the given circles by the above method. These are called the **direct common tangents**.

When the given circles are external to one another and do not intersect, two more common tangents may be drawn.

For, from centre A , with a radius equal to the *sum* of the radii of the given circles, describe a circle.

From B draw a tangent to this circle;

and proceed as before, but draw BE in the direction *opposite* to AD .

It follows from hypothesis that B is external to the circle used in the construction;

∴ two tangents may be drawn to it from B .

Hence *two* more common tangents may be drawn to the given circles: these will be found to pass between the given circles, and are called the **transverse common tangents**.

Thus, in general, *four* common tangents may be drawn to two given circles.

The student should investigate for himself the number of common tangents which may be drawn in the following special cases, noting in each case where the general construction fails, or is modified:-

- (i) When the given circles intersect;
- (ii) When the given circles have external contact;
- (iii) When the given circles have internal contact;
- (iv) When one of the given circles is wholly within the other.

18. Draw the direct common tangents to two equal circles.

19. If the two direct, or the two transverse, common tangents are drawn to two circles, the parts of the tangents intercepted between the points of contact are equal.

20. If four common tangents are drawn to two circles external to one another; shew that the two direct, and also the two transverse, tangents intersect on the straight line which joins the centres of the circles.

21. Two given circles have external contact at A , and a direct common tangent is drawn to touch them at P and Q : shew that PQ subtends a right angle at the point A .

22. Two circles have external contact at A , and a direct common tangent is drawn to touch them at P and Q : shew that a circle described on PQ as diameter is touched at A by the straight line which joins the centres of the circles.

23. Two circles whose centres are C and C' have external contact at A , and a direct common tangent is drawn to touch them at P and Q : shew that the bisectors of the angles PCA , $QC'A$ meet at right angles in PQ . And if R is the point of intersection of the bisectors, shew that RA is also a common tangent to the circles.

24. Two circles have external contact at A , and a direct common tangent is drawn to touch them at P and Q : shew that the square on PQ is equal to the rectangle contained by the diameters of the circles.

25. Draw a tangent to a given circle, so that the part of it intercepted by another given circle may be equal to a given straight line. Which is this impossible?

26. Draw a secant to two given circles, so that the parts of it intercepted by the circumferences may be equal to two given straight lines.

PROBLEMS ON TANGENCY.

The following exercises are solved by the Method of Intersection of Loci, explained on page 117.

The student should begin by making himself familiar with the following loci.

(i) *The locus of the centres of circles which pass through two given points.*

(ii) *The locus of the centres of circles which touch a given straight line at a given point.*

(iii) *The locus of the centres of circles which touch a given circle at a given point.*

(iv) *The locus of the centres of circles which touch a given straight line, and have a given radius.*

(v) *The locus of the centres of circles which touch a given circle, and have a given radius.*

(vi) *The locus of the centres of circles which touch two given straight lines.*

In each exercise the student should investigate the limits and relations among the data, in order that the problem may be possible.

27. Describe a circle to touch three given straight lines.

28. Describe a circle to pass through a given point and touch a given straight line at a given point.

29. Describe a circle to pass through a given point, and touch a given circle at a given point.

30. Describe a circle of given radius to pass through a given point, and touch a given straight line.

31. Describe a circle of given radius to touch two given circles.

32. Describe a circle of given radius to touch two given straight lines.

33. Describe a circle of given radius to touch a given circle and a given straight line.

34. Describe two circles of given radii to touch one another and a given straight line, on the same side of it.

35. If a circle touches a given circle and a given straight line, shew that the points of contact and an extremity of the diameter of the given circle at right angles to the given line are collinear.

36. *To describe a circle to touch a given circle, and also to touch a given straight line at a given point.*

Let DEB be the given circle, PQ the given st. line, and A the given point in it:

it is required to describe a circle to touch the $\odot DEB$, and also to touch PQ at A .

At A draw AF perp. to PQ . III. 11.
then the centre of the required circle must lie in AF . III. 19.

Find C , the centre of the $\odot DEB$, III. 1.
and draw a diameter BD perp. to PQ :

join A to one extremity D , cutting the \odot at E .


Join CE , and produce it to cut AF at F .

Then F is the centre, and FA the radius of the required circle.

[Supply the proof: and shew that a second solution is obtained by joining AB , and producing it to meet the \odot : also distinguish between the nature of the contact of the circles, when PQ cuts, touches, or is without the given circle.]

37. Describe a circle to touch a given straight line, and to touch a given circle at a given point.

38. Describe a circle to touch a given circle, have its centre in a given straight line, and pass through a given point in that straight



[For other problems of the same class see page 235.]

ORTHOGONAL CIRCLES.

DEFINITION. Circles which intersect at a point, so that the two tangents at that point are at right angles to one another, are said to be **orthogonal**, or **to cut one another orthogonally**.

39. In two intersecting circles the angle between the tangents at one point of intersection is equal to the angle between the tangents at the other.

40. *If two circles cut one another orthogonally, the tangent to each circle at a point of intersection will pass through the centre of the other circle.*

41. *If two circles cut one another orthogonally, the square on the distance between their centres is equal to the sum of the squares on their radii.*

42. Find the locus of the centres of all circles which cut a given circle orthogonally at a given point.

43. Describe a circle to pass through a given point and cut a given circle orthogonally at a given point.

III. ON ANGLES IN SEGMENTS, AND ANGLES AT THE CENTRES AND CIRCUMFERENCES OF CIRCLES.

See Propositions 20, 21, 22; 26, 27, 28, 29; 31, 32, 33, 34.

1. *If two chords intersect within a circle, they form an angle equal to that at the centre, subtended by half the sum of the arcs they cut off.*

Let AB and CD be two chords, intersecting at E within the given circle $ADBC$:
then shall the $\angle AEC$ be equal to the angle at the centre, subtended by half the sum of the arcs AC , BD .

Join AD .


Then the ext. $\angle AEC =$ the sum of the int. opp. $\angle^s EDA, EAD$;

that is, the sum of the $\angle^s CDA, BAD$.

But the $\angle^s CDA, BAD$ are the angles at the circ^e subtended by the arcs AC, BD ;

\therefore their sum $=$ half the sum of the angles at the centre subtended by the same arcs;

or, the $\angle AEC =$ the angle at the centre subtended by half the sum of the arcs AC, BD .



Q. E. D.

2. *If two chords when produced intersect outside a circle, they form an angle equal to that at the centre subtended by half the difference of the arcs they cut off.*

3. *The sum of the arcs cut off by two chords of a circle at right angles to one another is equal to the semi-circumference.*

4. *AB , AC are any two chords of a circle and P , Q are the middle points of the minor arcs cut off by them. If PQ is joined, cutting AB and AC at X , Y , shew that $AX = AY$.*

5. *If one side of a quadrilateral inscribed in a circle is produced, the exterior angle is equal to the opposite interior angle.*

6. *If two circles intersect, and any straight lines are drawn, one through each point of section, terminated by the circumferences; shew that the chords which join their extremities towards the same parts are parallel.*

7. *$ABCD$ is a quadrilateral inscribed in a circle, and the opposite sides AB , DC are produced to meet at P , and CB , DA to meet at Q ; if the circles circumscribed about the triangles PBC , QAB intersect at R , shew that the points P , R , Q are collinear.*

8. *If a circle is described on one of the sides of a right-angled triangle, then the tangent drawn to it at the point where it cuts the hypotenuse bisects the other side.*

9. *Given three points not in the same straight line: shew how to find any number of points on the circle which passes through them, without finding the centre.*

10. *Through any one of three given points not in the same straight line, draw a tangent to the circle which passes through them, without finding the centre.*

11. *Of two circles which intersect at A and B , the circumference of one passes through the centre of the other: from A any straight line is drawn to cut the first at C , the second at D ; shew that $CB = CD$.*

12. *Two tangents AP , AQ are drawn to a circle, and B is the middle point of the arc PQ , convex to A . Shew that PB bisects the angle APQ .*

13. *Two circles intersect at A and B ; and at A tangents are drawn, one to each circle, to meet the circumferences at C and D : if CB , BD are joined, shew that the triangles ABC , DBA are equiangular to one another.*

14. *Two segments of circles are described on the same chord and on the same side of it; the extremities of the common chord are joined to any point on the arc of the exterior segment: shew that the arc intercepted on the interior segment is constant.*

15. If a series of triangles are drawn standing on a fixed base, and having a given vertical angle, shew that the bisectors of the vertical angles all pass through a fixed point.

16. ABC is a triangle inscribed in a circle, and E the middle point of the arc subtended by BC on the side remote from A ; if through E a diameter ED is drawn, shew that the angle DEA is half the difference of the angles at B and C . [See Ex. 7, p. 101.]

17. If two circles touch each other internally at a point A , any chord of the exterior circle which touches the interior is divided at its point of contact into segments which subtend equal angles at A .

18. If two circles touch one another internally, and a straight line is drawn to cut them, the segments of it intercepted between the circumferences subtend equal angles at the point of contact.

THE ORTHOCENTRE OF A TRIANGLE

* 19. *The perpendiculars drawn from the vertices of a triangle to the opposite sides are concurrent.*

In the $\triangle ABC$, let AD , BE be the perp^s drawn from A and B to the opposite sides; and let them intersect at O . Join CO , and produce it to meet AB at F .

It is required to shew that CF is perp to AB .

Join DE .

Then, because the \angle 's OEC , ODC are rt. angles,

Hyp.

\therefore the points O , E , C , D are concyclic.

the $\angle DEC =$ the $\angle DOC$, in the same segment;

\therefore the vert. opp. $\angle FOA$.

Again, because the \angle 's AEB , ADB are rt. angles,

Hyp.

\therefore the points A , E , D , B are concyclic:

\therefore the $\angle DEB =$ the $\angle DAB$, in the same segment.

the sum of the \angle 's FOA , $FAO =$ the sum of the \angle 's DEC , DEB

$=$ a rt. angle:


Hyp.

the remaining $\angle AFO =$ a rt. angle:

I. 32.

that is, CF is perp. to AB .

Hence the three perp^s AD , BE , CF meet at the point O . Q. E. D.



[For an Alternative Proof see page 106.]

DEFINITIONS.

(i) The intersection of the perpendiculars drawn from the vertices of a triangle to the opposite sides is called its **orthocentre**.


(ii) The triangle formed by joining the feet of the perpendiculars is called the **pedal** or **orthocentric triangle**.

20. In an acute-angled triangle the perpendiculars drawn from the vertices to the opposite sides bisect the angles of the pedal triangle through which they pass.

In the acute-angled $\triangle ABC$, let AD , BE , CF be the perp^s drawn from the vertices to the opposite sides, meeting at the orthocentre O , and let DEF be the pedal triangle.

then shall AD , BE , CF bisect respectively the \angle 's FDE , DEF , EFD .

For, as in the last theorem, it may be shewn that the points O , D , C , E are concyclic;



\therefore the \angle ODE = the \angle OCE , in the same segment.

Similarly the points O , D , B , F are concyclic,

\therefore the \angle ODF = the \angle OBF , in the same segment.

But the \angle OCE = the \angle OBF , each being the comp^t of the \angle BAC .
 \therefore the \angle ODE = the \angle ODF .

Similarly it may be shewn that the \angle 's DEF , EFD are bisected by BE and CF . Q. E. D.

COROLLARY. (i) Every two sides of the pedal triangle are equally inclined to that side of the original triangle in which they meet.

For the \angle EDC = the comp^t of the \angle ODE
 $=$ the comp^t of the \angle OCE
 $=$ the \angle BAC .

Similarly it may be shewn that the \angle FDB = the \angle BAC ,
 \therefore the \angle EDC = the \angle FDB = the \angle A .

In like manner it may be proved that


the \angle DEC = the \angle FEA = the \angle B ,
 and the \angle DFB = the \angle EFA = the \angle C .

COROLLARY. (ii) The triangles DEC , AEF , DBF are equiangular to one another and to the triangle ABC .

NOTE. If the angle BAC is obtuse, then the perpendiculars BE , CF bisect externally the corresponding angles of the pedal triangle.

21. *In any triangle, if the perpendiculars drawn from the vertices on the opposite sides are produced to meet the circumscribed circle, then each side bisects that portion of the line perpendicular to it which lies between the orthocentre and the circumference.*

Let ABC be a triangle in which the perpendiculars AD , BE are drawn, intersecting at O the orthocentre; and let AD be produced to meet the $circ$ of the circumscribing circle at G
then shall $DO = DG$.



Join BG .

Then in the two \triangle^s OEA , ODB ,
the $\angle OEA$, the $\angle ODB$, being rt. angles,
and the $\angle EOA$ the vert. opp. $\angle DOB$;
the remaining $\angle EAO$ the remaining $\angle DBO$. I. 32.

But the $\angle CAG$ = the $\angle CBG$, in the same segment;
the $\angle DBO$ the $\angle DBG$.

The \angle in the \triangle^s DBO , DBG .

Because $\left\{ \begin{array}{l} \text{the } \angle DBO = \text{the } \angle DBG, \\ \text{the } \angle BDO = \text{the } \angle BDG, \\ \text{and } BD \text{ is common;} \end{array} \right.$ *Proved.*
DO = DG. I. 26.

Q. E. D.

22. *In an acute-angled triangle the three sides are the external bisectors of the angles of the pedal triangle; and in an obtuse-angled triangle the sides containing the obtuse angle are the internal bisectors of the corresponding angles of the pedal triangle.*

23. *If O is the orthocentre of the triangle ABC , shew that the angles BOC , BAC are supplementary.*

24. *If O is the orthocentre of the triangle ABC , then any one of the four points O , A , B , C is the orthocentre of the triangle whose vertices are the other three.*

25. *The three circles which pass through two vertices of a triangle and its orthocentre are each equal to the circle circumscribed about the triangle.*

26. D , E are taken on the circumference of a semicircle described on a given straight line AB ; the chords AD , BE and AE , BD intersect (produced if necessary) at F and G : shew that FG is perpendicular to AB .

27. $ABCD$ is a parallelogram; AE and CE are drawn at right angles to AB , and CB respectively: shew that ED , if produced, will be perpendicular to AC .

28. ABC is a triangle, O is its orthocentre, and AK a diameter of the circumscribed circle: shew that $BOCK$ is a parallelogram.

29. The orthocentre of a triangle is joined to the middle point of the base, and the joining line is produced to meet the circumscribed circle: prove that it will meet it at the same point as the diameter which passes through the vertex.

30. The perpendicular from the vertex of a triangle on the base, and the straight line joining the orthocentre to the middle point of the base, are produced to meet the circumscribed circle at P and Q : shew that PQ is parallel to the base.

31. The distance of each vertex of a triangle from the orthocentre is double of the perpendicular drawn from the centre of the circumscribed circle on the opposite side.

32. Three circles are described each passing through the orthocentre of a triangle and two of its vertices: shew that the triangle formed by joining their centres is equal in all respects to the original triangle.

33. ABC is a triangle inscribed in a circle, and the bisectors of its angles which intersect at O are produced to meet the circumference in PQR : shew that O is the orthocentre of the triangle PQR .

34. Construct a triangle, having given a vertex, the orthocentre, and the centre of the circumscribed circle.

Locus

35. Given the base and vertical angle of a triangle, find the locus of its orthocentre.

Let BC be the given base, and X the given angle; and let BAC be any triangle on the base BC , having its vertical $\angle A$ equal to the $\angle X$.

Draw the perp^s BE , CF , intersecting at the orthocentre O .

It is required to find the locus of O .

Since the \angle 's OFA , OEA are rt. angles, the points O , F , A , E are concyclic, the $\angle FOE$ is the supplement of the $\angle A$;

the vert. opp. $\angle BOC$ is the supplement of the $\angle A$.

But the $\angle A$ is constant, being always equal to the $\angle X$;

\therefore its supplement is constant;

that is, the ΔBOC has a fixed base, and constant vertical angle; hence the focus of its vertex O is the arc of a segment of which BC is the chord.

F ,

E

B

C

III. 22.

[See p. 187.]

36. *Given the base and vertical angle of a triangle, find the locus of the intersection of the bisectors of its angles.*

Let BAC be any triangle on the given base BC , having its vertical angle equal to the given $\angle X$; and let AI , BI , CI be the bisectors of its angles: [see Ex. 2, p. 103.] it is required to find the locus of the point I .

Denote the angles of the $\triangle ABC$ by A, B, C ; and let the $\angle BIC$ be denoted by I .

Then from the $\triangle BIC$,

(i) $I + \frac{1}{2}B + \frac{1}{2}C = \text{two rt. angles}$, I. 32.
and from the $\triangle ABC$,

$A + B + C = \text{two rt. angles}$; I. 32.

(ii) so that $\frac{1}{2}A + \frac{1}{2}B + \frac{1}{2}C = \text{one rt. angle}$,

, taking the differences of the equals in (i) and (ii),

$I - \frac{1}{2}A = \text{one rt. angle}$;


or,

$I = \text{one rt. angle} + \frac{1}{2}A$.

But A is constant, being always equal to the $\angle X$,

$\therefore I$ is constant:

\therefore , since the base BC is fixed, the locus of I is the arc of a segment of which BC is the chord.



37. *Given the base and vertical angle of a triangle, find the locus of the centroid, that is, the intersection of the medians.*

Let BAC be any triangle on the given base BC , having its vertical angle equal to the given angle S ; let the medians AX , BY , CZ intersect at the centroid G [see Ex. 4, p. 105]:

it is required to find the locus of the point G .

Through G draw GP , GQ par^l to AB and AC respectively.

Then ZG is a third part of ZC ;

Ex. 4, p. 105.

and since GP is par^l to ZB ,

$\therefore BP$ is a third part of BC .

Ex. 19, p. 99.

Similarly QC is a third part of BC ;

$\therefore P$ and Q are fixed points.

Now since PG , GQ are par^l respectively to BA , AC ,

Constr.

\therefore the $\angle PGQ =$ the $\angle BAC$,


I. 29.

$=$ the $\angle S$,

that is, the $\angle PGQ$ is constant;

and since the base PQ is fixed,

\therefore the locus of G is the arc of a segment of which PQ is the chord.



Obs. In this problem the points A and G move on the arcs of similar segments.

38. Given the base and the vertical angle of a triangle; find the locus of the intersection of the bisectors of the exterior base angles.

39. Through the extremities of a given straight line AB any two parallel straight lines AP , BQ are drawn; find the locus of the intersection of the bisectors of the angles PAB , QBA .

40. Find the locus of the middle points of chords of a circle drawn through a fixed point.

Distinguish between the cases when the given point is within, on, or without the circumference.

41. Find the locus of the points of contact of tangents drawn from a fixed point to a system of concentric circles.

42. Find the locus of the intersection of straight lines which pass through two fixed points on a circle and intercept on its circumference an arc of constant length.

43. A and B are two fixed points on the circumference of a circle, and PQ is any diameter: find the locus of the intersection of PA and QB .

44. BAC is any triangle described on the fixed base BC and having a constant vertical angle; and BA is produced to P , so that BP is equal to the sum of the sides containing the vertical angle: find the locus of P .

45. AB is a fixed chord of a circle, and AC is a moveable chord passing through A ; if the parallelogram CB is completed, find the locus of the intersection of its diagonals.

46. A straight rod PQ slides between two rulers placed at right angles to one another, and from its extremities PX , QX are drawn perpendicular to the rulers: find the locus of X .

47. Two circles whose centres are C and D , intersect at A and B : through A , any straight line PAQ is drawn terminated by the circumferences; and PC , QD intersect at X : find the locus of X , and shew that it passes through B . [Ex. 9, p. 216.]

48. Two circles intersect at A and B , and through P , any point on the circumference of one of them, two straight lines PA , PB are drawn, and produced if necessary, to cut the other circle at X and Y : find the locus of the intersection of AY and BX .

49. Two circles intersect at A and B ; HAK is a fixed straight line drawn through A and terminated by the circumferences, and PAQ is any other straight line similarly drawn: find the locus of the intersection of HP and QK .

50. Two segments of circles are on the same chord AB and on the same side of it; and P and Q are any points one on each arc: find the locus of the intersection of the bisectors of the angles PAQ , PBQ .

51. Two circles intersect at A and B ; and through A any straight line PAQ is drawn terminated by the circumferences: find the locus of the middle point of PQ .

MISCELLANEOUS EXAMPLES ON ANGLES IN A CIRCLE.

52. ABC is a triangle, and circles are drawn through B, C , cutting the sides in P, Q, P', Q' : shew that $PQ, P'Q'$ are parallel to one another and to the tangent drawn at A to the circle circumscribed about the triangle.

53. Two circles intersect at B and C , and from any point A , on the circumference of one of them, AB, AC are drawn, and produced if necessary, to meet the other at D and E : shew that DE is parallel to the tangent at A .

54. A secant PAB and a tangent PT are drawn to a circle from an external point P ; and the bisector of the angle ATB meets AB at C : shew that PC is equal to PT .

55. From a point A on the circumference of a circle two chords AB, AC are drawn, and also the diameter AF : if AB, AC are produced to meet the tangent at F in D and E , shew that the triangles ABC, AED are equiangular to one another.

56. O is any point within a triangle ABC , and OD, OE, OF are drawn perpendicular to BC, CA, AB respectively: shew that the angle BOC is equal to the sum of the angles BAC, EDF .

57. If two tangents are drawn to a circle from an external point, shew that they contain an angle equal to the difference of the angles in the segments cut off by the chord of contact.

58. Two circles intersect, and through a point of section a straight line is drawn bisecting the angle between the diameters through that point: shew that this straight line cuts off similar segments from the two circles.

59. Two equal circles intersect at A and B ; and from centre A , with any radius less than AB a third circle is described cutting the given circles on the same side of AB at C and D : shew that the points B, C, D are collinear.

60. ABC and $A'B'C'$ are two triangles inscribed in a circle, so that AB, AC are respectively parallel to $A'B', A'C'$: shew that BC' is parallel to $B'C$.

61. Two circles intersect at A and B , and through A two straight lines HAK , PAQ are drawn terminated by the circumferences: if HP and KQ intersect at X , shew that the points H , B , K , X are concyclic.

62. Describe a circle touching a given straight line at a given point, so that tangents drawn to it from two fixed points in the given line may be parallel. [See Ex 10, p. 183]

63. C is the centre of a circle, and CA , CB two fixed radii: if from any point P on the arc AB perpendiculars PX , PY are drawn to CA and CB , shew that the distance XY is constant.

64. AB is a chord of a circle, and P any point in its circumference; PM is drawn perpendicular to AB , and AN is drawn perpendicular to the tangent at P : shew that MN is parallel to PB .

65. P is any point on the circumference of a circle of which AB is a fixed diameter, and PN is drawn perpendicular to AB , on AN and BN as diameters circles are described, which are cut by AP , BP at X and Y : shew that XY is a common tangent to these circles.

66. Upon the same chord and on the same side of it three segments of circles are described containing respectively a given angle, its supplement and a right angle: shew that the intercept made by the two former segments upon any straight line drawn through an extremity of the given chord is bisected by the latter segment.

67. Two straight lines of indefinite length touch a given circle, and any chord is drawn so as to be bisected by the chord of contact: if the former chord is produced, shew that the intercepts between the circumference and the tangents are equal.

68. Two circles intersect one another: through one of the points of contact draw a straight line of given length terminated by the circumferences.

69. On the three sides of any triangle equilateral triangles are described remote from the given triangle: shew that the circles described about them intersect at a point.

70. On BC , CA , AB the sides of a triangle ABC , any points P , Q , R are taken; shew that the circles described about the triangles AQR , BRP , CPQ meet in a point.

71. Find a point within a triangle at which the sides subtend equal angles.

72. Describe an equilateral triangle so that its sides may pass through three given points.

73. Describe a triangle equal in all respects to a given triangle, and having its sides passing through three given points.

SIMSON'S LINE.

74. If from any point on the circumference of the circle circumscribed about a triangle, perpendiculars are drawn to the three sides, the feet of these perpendiculars are collinear.

Let P be any point on the c^e of the circle circumscribed about the $\triangle ABC$; and let PD , PE , PF be the perps drawn from P to the three sides.

It is required to prove that the points D , E , F are collinear.

Join FD and DE :

then FD and DE shall be in the same st. line.

Join PB , PC

Because the \angle 's PDB , PFB are rt. angles, Hyp.
 \therefore the points P , D , B , F are concyclic;

the \angle PDF = the \angle PBF , in the same segment. III. 21.

But since $BACP$ is a quad^r inscribed in a circle, having one of its sides AB produced to F ,

the ext. \angle PBF = the opp. int. \angle ACP . Euc. 3, p. 188.

\therefore the \angle PDF = the \angle ACP .

To each add the \angle PDE :

then the \angle 's PDF , PDE = the \angle 's ACP , PDE ;

But since the \angle 's PDC , PEC are rt. angles,

\therefore the points P , D , E , C are concyclic;

the \angle 's ACP , PDE together = two rt. angles;

the \angle 's PDF , PDE together = two rt. angles;

\therefore FD and DE are in the same st. line; I. 14.


that is, the points D , E , F are collinear. Q.E.D.

[The line FDE is called the **Pedal** or **Simson's Line** of the triangle ABC for the point P ; though the tradition attributing the theorem to Robert Simson has been recently shaken by the researches of Dr. J. S. Mackay.]

75. ABC is a triangle inscribed in a circle; and from any point P on the circumference PD , PF are drawn perpendicular to BC and AB : if FD , or FD produced, cuts AC at E , shew that PE is perpendicular to AC .

76. Find the locus of a point which moves so that if perpendiculars are drawn from it to the sides of a given triangle, their feet are collinear.

77. ABC and $AB'C'$ are two triangles having a common vertical angle, and the circles circumscribed about them meet again at P : shew that the feet of perpendiculars drawn from P to the four lines AB , AC , BC , $B'C'$ are collinear.



78. A triangle is inscribed in a circle, and any point P on the circumference is joined to the orthocentre of the triangle: shew that this joining line is bisected by the pedal of the point P .

IV. ON THE CIRCLE IN CONNECTION WITH RECTANGLES.

See Propositions 35, 36, 37

1. If from any external point P two tangents are drawn to a given circle whose centre is O , and if OP meet the chord of contact at Q ; then the rectangle OP, OQ is equal to the square on the radius.

Let PH, PK be tangents, drawn from the external point P to the $\odot HAK$, whose centre is O , and let OP meet HK the chord of contact at Q , and the \odot at A ; then shall the rect. OP, OQ = the sq. on OA .


On HP as diameter describe a circle: this circle must pass through Q , since the $\angle HQP$ is a rt. angle. III. 31.

Join OH .

Then since PH is a tangent to the $\odot HAK$,
 \therefore the $\angle OHP$ is a rt. angle.

And since HP is a diameter of the $\odot HQP$,

$\therefore OH$ touches the $\odot HQP$ at H
 \therefore the rect. OP, OQ = the sq. on OH .
 = the sq. on OA .



III. 16.

III. 36.

Q. E. D.

2. ABC is a triangle, and AD, BE, CF the perpendiculars drawn from the vertices to the opposite sides, meeting in the orthocentre O : shew that the rect. AO, OD = the rect. BO, OE = the rect. CO, OF .

3. ABC is a triangle, and AD, BE the perpendiculars drawn from A and B on the opposite sides: shew that the rectangle CA, CE is equal to the rectangle CB, CD .

4. ABC is a triangle right-angled at C , and from D , any point in the hypotenuse AB , a straight line DE is drawn perpendicular to AB and meeting BC at E : shew that the square on DE is equal to the difference of the rectangles AD, DB and CE, EB .

5. From an external point P two tangents are drawn to a given circle whose centre is O , and OP meets the chord of contact at Q : shew that any circle which passes through the points P, Q will cut the given circle orthogonally. [See Def. p. 222.]

6. A series of circles pass through two given points, and from a fixed point in the common chord produced tangents are drawn to all the circles: shew that the points of contact lie on a circle which cuts all the given circles orthogonally.

7. All circles which pass through a fixed point, and cut a given circle orthogonally, pass also through a second fixed point.

8. Find the locus of the centres of all circles which pass through a given point and cut a given circle orthogonally.

9. Describe a circle to pass through two given points and cut a given circle orthogonally.

10. A, B, C, D are four points taken in order on a given straight line: find a point O between B and C such that the rectangle OA, OB may be equal to the rectangle OC, OD.

11. AB is a fixed diameter of a circle, and CD a fixed straight line of indefinite length cutting AB or AB produced at right angles: any straight line is drawn through A to cut CD at P and the circle at Q: shew that the rectangle AP, AQ is constant.

12. AB is a fixed diameter of a circle, and CD a fixed chord at right angles to AB: any straight line is drawn through A to cut CD at P and the circle at Q: shew that the rectangle AP, AQ is equal to the square on AC.

13. A is a fixed point and CD a fixed straight line of indefinite length: AP is any straight line drawn through A to meet CD at P, and in AP a point Q is taken such that the rectangle AP, AQ is constant: find the locus of Q.

14. Two circles intersect orthogonally, and tangents are drawn from any point on the circumference of one to touch the other: prove that the first circle passes through the middle point of the chord of contact of the tangents. [Ex 1, p 253.]

15. A semicircle is described on AB as diameter, and any two chords AC, BD are drawn intersecting at P: shew that

$$AB^2 = AC \cdot AP + BD \cdot BP.$$

16. Two circles intersect at B and C, and the two direct common tangents AE and DF are drawn: if the common chord is produced to meet the tangents at G and H, shew that $GH^2 = AE^2 + BC^2$.

17. If from a point P, without a circle, PM is drawn perpendicular to a diameter AB, and also a secant PCD, shew that

$$PM^2 = PC \cdot PD + AM \cdot MB.$$

18. Three circles intersect at D , and their other points of intersection are A, B, C ; AD cuts the circle BDC at E , and EB, EC cut the circles ADB, ADC respectively at F and G : shew that the points F, A, G are collinear, and F, B, C, G concyclic.

19. A semicircle is described on a given diameter BC , and from B and C any two chords BE, CF are drawn intersecting within the semicircle at O ; BF and CE are produced to meet at A : shew that the sum of the squares on AB, AC is equal to twice the square on the tangent from A together with the square on BC .

20. X and Y are two fixed points in the diameter of a circle equidistant from the centre C : through X any chord PXQ is drawn, and its extremities are joined to Y ; shew that the sum of the squares on the sides of the triangle PYQ is constant. [See p. 117, Ex. 21]

PROBLEMS ON TANGENCY

21. To describe a circle to pass through two given points and to touch a given straight line.

Let A and B be the given points, and CD the given st. line; it is required to describe a circle to pass through A and B and to touch CD .

Join BA , and produce it to meet CD at P .

Describe a square equal to the rect. PA, PB ; in 11.


and from PD (or PC) cut off PQ equal to a side of this square.

Through A, B and Q describe a circle. Ex. 1, p. 156.
Then since the rect. PA, PB = the sq. on PQ ,
 \therefore the $\circ ABQ$ touches CD at Q . II. 37.

Q. E. D.

NOTE. (i) Since PQ may be taken on either side of P , it is clear that there are in general two solutions of the problem.

(ii) When AB is parallel to the given line CD , the above method is not applicable. In this case a simple construction follows from III. 1, Cor. and III. 16 and it will be found that only one solution exists.



22. To describe a circle to pass through two given points and to touch a given circle.

Let A and B be the given points, and CRP the given circle:

it is required to describe a circle to pass through A and B , and to touch the $\odot CRP$.

Through A and B describe any circle to cut the given circle at P and Q .

Join AB , PQ , and produce them to meet at D .

From D draw DC to touch the given circle, and let C be the point of contact.

Then the circle described through A , B , C will touch the given circle.

For, from the $\odot ABQP$, the rect. DA , DB = the rect. DP , DQ :
and from the $\odot PQC$, the rect. DP , DQ = the sq. on DC , III. 36.

\therefore the rect. DA , DB = the sq. on DC .

$\therefore DC$ touches the $\odot ABC$ at C . III. 37.

But DC touches the $\odot PQC$ at C ; Constr.

\therefore the $\odot ABC$ touches the given circle, and it passes through the given points A and B . Q.E.D.

NOTE. (i) Since two tangents may be drawn from D to the given circle, it follows that there will be two solutions of the problem.

(ii) The general construction fails when the straight line bisecting AB at right angles passes through the centre of the given circle: the problem then becomes symmetrical, and the solution is obvious.

23. To describe a circle to pass through a given point and to touch two given straight lines.



Let P be the given point, and AB , AC the given straight lines: it is required to describe a circle to pass through P and to touch AB , AC .

Now the centre of every circle which touches AB and AC must lie on the bisector of the $\angle BAC$.

Ex. 7, p. 183.

Hence draw AE bisecting the $\angle BAC$.

From P draw PK perp. to AE , and produce it to P' , making KP' equal to PK .



Then every circle which has its centre in AE , and passes through P , must also pass through P' . Ex. 1, p. 215.

Hence the problem is now reduced to drawing a circle through P and P' to touch *either* AC or AB . Ex. 21, p. 235.

• Produce PP' to meet AC at S .

Describe a square equal to the rect. SP, SP' , II. 14.
and cut off SR equal to a side of the square.

• Describe a circle through the points P', P, R ;
then since the rect. SP, SP' = the sq. on SR , Constr.
 \therefore the circle touches AC at R ; III. 37.

and since its centre is in AE , the bisector of the $\angle BAC$,
it may be shewn also to touch AB . Q. 1. P'

NOTE (i) Since SR may be taken on either side of S , it follows that there will be two solutions of the problem.

(ii) If the given straight lines are parallel, the centre lies on the parallel straight line mid-way between them, and the construction proceeds as before.

21. To describe a circle to touch two given straight lines and a given circle.

Let AB, AC be the two given st. lines, and D the centre of the given circle:

it is required to describe a circle to touch AB, AC and the circle whose centre is D .

Draw EF, GH par^l to AB and AC respectively, on the sides remote from D , and at distances from them equal to the radius of the given circle.

Describe the $\odot MND$ to touch EF and GH at M and N , and to pass through D . Ex. 23, p. 236.

Let O be the centre of this circle.

Join OM, ON, OD meeting AB, AC and the given circle at P, Q and R .

Then a circle described from centre O with radius OP will touch AB, AC and the given circle.

For since O is the centre of the $\odot MND$,

$\therefore OM = ON = OD$.

But $PM = QN = RD$; Constr.


$\therefore OP = OQ = OR$.

\therefore a circle described from centre O , with radius OP , will pass through Q and R .

And since the \angle 's at M and N are rt. angles, III. 18.

\therefore the \angle 's at P and Q are rt. angles; I. 29.

\therefore the $\odot PQR$ touches AB and AC .



And since R , the point in which the circles meet, is on the line of centres OD ,

\therefore the $\odot PQR$ touches the given circle. Q. E. F.


NOTE. There will be two solutions of this problem, since two circles may be drawn to touch EF , GH and to pass through D .

25. To describe a circle to pass through a given point and touch a given straight line and a given circle

Let P be the given point, AB the given st. line, and DHE the given circle, of which C is the centre: it is required to describe a circle to pass through P , and to touch AB and the $\odot DHE$.

Through C draw $DCEF$ perp. to AB , cutting the circle at the points D and E , of which E is between C and AB .

Join DP ; and by describing a circle through F , E , and P , find a point K in DP (or DP produced) such that the rect. DE , $DF =$ the rect. DK , DP .



Describe a circle to pass through P , K and touch AB : Ex. 21, p. 235. This circle shall also touch the given $\odot DHE$.

For let G be the point at which this circle touches AB .

Join DG , cutting the given circle DHE at H .

Join HE .

Then the $\angle DHE$ is a rt. angle, being in a semicircle. III. 31.

also the angle at F is a rt. angle; Constr.

\therefore the points E , F , G , H are concyclic.

\therefore the rect. DE , $DF =$ the rect. DH , DG . III. 36.

but the rect. DE , $DF =$ the rect. DK , DP : Constr.

\therefore the rect. DH , $DG =$ the rect. DK , DP :

\therefore the point H is on the $\odot PKG$.

Let O be the centre of the $\odot PHG$.

Join OG , OH , CH .

Then OG and DF are paral. since they are both perp. to AB ; and DG meets them.

\therefore the $\angle OGD =$ the $\angle GDC$. I. 29.

But since $OG = OH$, and $CD = CH$,

\therefore the $\angle OGH =$ the $\angle OHG$; and the $\angle CDH =$ the $\angle CHD$:

\therefore the $\angle OHG =$ the $\angle CHD$;

\therefore OH and CH are in one st. line.

\therefore the $\odot PHG$ touches the given $\odot DHE$. Q. E. F.

NOTE. (i) Since two circles may be drawn to pass through P, K and to touch AB, it follows that there will be two solutions of the present problem.

(ii) Two more solutions may be obtained by joining PE, and proceeding as before.

The student should examine the nature of the contact between the circles in each case.

26. Describe a circle to pass through a given point, to touch a given straight line, and to have its centre on another given straight line.

27. Describe a circle to pass through a given point, to touch a given circle, and to have its centre on a given straight line.

28. Describe a circle to pass through two given points, and to intersect an arc of given length on a given circle.

29. Describe a circle to touch a given circle and a given straight line at a given point.

30. Describe a circle to touch two given circles and a given straight line.

V. ON MAXIMA AND MINIMA.

We gather from the Theory of Loci that the position of an angle, line or figure is capable under suitable conditions of gradual change; and it is usually found that change of *position* involves a corresponding and gradual change of *magnitude*.

Under these circumstances we may be required to note if any situations exist at which the magnitude in question, after increasing, begins to decrease; or after decreasing, to increase: in such situations the Magnitude is said to have reached a **Maximum** or a **Minimum** value; for in the former case it is greater, and in the latter case less than in adjacent situations on either side. In the geometry of the circle and straight line we only meet with such cases of continuous change as admit of one transition from an increasing to a decreasing state—or vice versa—so that in all the problems with which we have to deal (where a single circle is involved) there can be only one Maximum and one Minimum—the Maximum being the greatest, and the Minimum being the least value that the variable magnitude is capable of taking.

Thus a variable geometrical magnitude reaches its maximum or minimum value at a *turning point*, towards which the magnitude may mount or descend from either side: it is natural therefore to expect a maximum or minimum value to occur when, in the course of its change, the magnitude assumes a *symmetrical* form or position; and this is usually found to be the case.

This general connection between a symmetrical form or position and a maximum or minimum value is not exact enough to constitute a *proof* in any particular problem; but by means of it a situation is suggested, which on further examination may be shewn to give the maximum or minimum value sought for.

For example, suppose it is required
to determine the greatest straight line that may be drawn perpendicular to the chord of a segment of a circle and intercepted between the chord and the arc:

we immediately anticipate that the greatest perpendicular is that which occupies a *symmetrical* position in the figure, namely the perpendicular which passes through the middle point of the chord; and on further examination this may be proved to be the case by means of I. 19, and I. 34.

Again we are able to find at what point a geometrical magnitude, varying under certain conditions, assumes its Maximum or Minimum value, if we can discover a construction for drawing the magnitude so that it may have an *assigned* value: for we may then examine between what limits the assigned value must lie in order that the construction may be possible; and the higher or lower limit will give the Maximum or Minimum sought for.


It was pointed out in the chapter on the Intersection of Loci, [see page 119] that if under certain conditions existing among the data, *two* solutions of a problem are possible, and under other conditions, *no* solution exists, there will always be some intermediate condition under which *one and only one* distinct solution is possible.

Under these circumstances this single or limiting solution will always be found to correspond to the maximum or minimum value of the magnitude to be constructed.

1. For example, suppose it is required
to divide a given straight line so that the rectangle contained by the two segments may be a maximum.

We may first attempt to divide the given straight line so that the rectangle contained by its segments may have a *given* area—that is, be equal to the square on a given straight line.

Let AB be the given straight line, and K the side of the given square:



it is required to divide the st. line AB at a point M , so that the rect. AM , MB may be equal to the sq. on K .

Adopting a construction suggested by II. 11,

describe a semicircle on AB ; and at any point X in AB , or AB produced, draw XY perp. to AB , and equal to K .

Through Y draw YZ par^l to AB , to meet the arc of the semicircle at P .

Then if the perp. PM is drawn to AB , it may be shewn after the manner of II. 11, or by III. 35 that

$$\begin{aligned} \text{the rect. } AM, MB &= \text{the sq. on } PM. \\ &= \text{the sq. on } K. \end{aligned}$$

So that the rectangle AM , MB increases as K increases.

Now if K is less than the radius CD , then YZ will meet the arc of the semicircle in two points P , P' ; and it follows that AB may be divided at *two* points, so that the rectangle contained by its segments may be equal to the square on K . If K increases, the st. line YZ will recede from AB , and the points of intersection P , P' will continually approach one another; until, when K is equal to the radius CD , the st. line YZ (now in the position $Y'Z'$) will meet the arc in *two coincident points*, that is, will touch the semicircle at D ; and there will be only *one* solution of the problem.

If K is greater than CD , the straight line YZ will not meet the semicircle, and the problem is impossible.

Hence the greatest length that K may have, in order that the construction may be possible, is the radius CD .

\therefore the rect. AM , MB is a maximum, when it is equal to the square on CD ;

that is, when PM coincides with DC , and consequently when M is the middle point of AB .

Obs. The special feature to be noticed in this problem is that the maximum is found at the transitional point between *two* solutions and *no* solution; that is, when the two solutions coincide and become identical.

The following example illustrates the same point.

2. *To find at what point in a given straight line the angle subtended by the line joining two given points, which are on the same side of the given straight line, is a maximum.*

Let CD be the given st. line, and A, B the given points on the same side of CD :

it is required to find at what point in CD the angle subtended by the st. line AB is a maximum.

First determine at what point in CD , the st. line AB subtends a given angle.

This is done as follows:—

On AB describe a segment of a circle containing an angle equal to the given angle. III. 33.

If the arc of this segment intersects CD , two points in CD are found at which AB subtends the given angle: but if the arc does not meet CD , no solution is given.

In accordance with the principles explained above, we expect that a maximum angle is determined at the limiting position, that is, when the arc touches CD ; or meets it at two coincident points.

[See page 213.]

This we may prove to be the case.

Describe a circle to pass through A and B , and to touch the st. line CD .

[Ex. 21, p. 235.]

Let P be the point of contact.

Then shall the $\angle APB$ be greater than any other angle subtended by AB at a point in CD on the same side of AB as P .

For take Q , any other point in CD , on the same side of AB as P ;

and join AQ, QB .

Since Q is a point in the tangent other than the point of contact, it must be without the circle,

\therefore either BQ or AQ must meet the arc of the segment APB .

Let BQ meet the arc at K : join AK .


Then the $\angle APB =$ the $\angle AKB$, in the same segment: but the ext. $\angle AKB$ is greater than the int. opp. $\angle AQB$.

\therefore the $\angle APB$ is greater than AQB .

Similarly the $\angle APB$ may be shewn to be greater than any other angle subtended by AB at a point in CD on the same side of AB :

that is, the $\angle APB$ is the greatest of all such angles. Q. E. D.

NOTE. Two circles may be described to pass through A and B , and to touch CD , the points of contact being on opposite sides of AB :



hence two points in CD may be found such that the angle subtended by AB at each of them is greater than the angle subtended at any other point in CD on the same side of AB .

We add two more examples of considerable importance.

3. In a straight line of indefinite length find a point such that the sum of its distances from two given points, on the same side of the given line, shall be a minimum.

Let CD be the given st. line of indefinite length, and A, B the given points on the same side of CD : it is required to find a point P in CD such that the sum of AP, PB is a minimum.

Draw AF perp. to CD ; and produce AF to E , making FE equal to AF .

$C \quad F$

Join EB , cutting CD at P .

Join AP, PB .

Then of all lines drawn from A and B to a point in CD ,

the sum of AP, PB shall be the least

For, let Q be any other point in CD .

Join AQ, BQ, EQ .

Now in the $\Delta AFP, EFP$,

Because $\left\{ \begin{array}{l} AF = EF, \\ \text{and } FP \text{ is common,} \\ \text{and the } \angle AFP = \text{the } \angle EFP, \text{ being rt. angles.} \end{array} \right.$ Constn
i. 1.

$\therefore AP = EP$.

Similarly it may be shewn that

$AQ = EQ$.

Now in the ΔEQB , the two sides EQ, QB are together greater than EB ;

hence, AQ, QB are together greater than EB ,
that is, greater than AP, PB .

Similarly the sum of the st. lines drawn from A and B to any other point in CD may be shewn to be greater than AP, PB .

\therefore the sum of AP, PB is a minimum.

Q. E. D.

NOTE. It follows from the above proof that

$\left\{ \begin{array}{l} \text{the } \angle APF = \text{the } \angle EPF \\ \qquad \qquad \qquad = \text{the } \angle BPD. \end{array} \right.$ i. 4.
i. 15.


Thus the sum of AP, PB is a minimum, when those lines are equally inclined to CD .

4. Given two intersecting straight lines AB , AC , and a point P between them; shew that of all straight lines which pass through P and are terminated by AB , AC , that which is bisected at P cuts off the triangle of minimum area.

Let EF be the st. line, terminated by AB , AC , which is bisected at P :
then the $\triangle FAE$ shall be of minimum area.

For let HK be any other st. line passing through P :
through E draw EM par^l to AC .

Then in the $\triangle HPF$, MPE ,



Because {	the $\angle HPF$ = the $\angle MPE$,	1. 15.
	and the $\angle HFP$ = the $\angle MEP$,	1. 29.
	and FP = EP ;	<i>Hyp.</i>
	the $\triangle HPF$ = the $\triangle MPE$	1. 26, Cor.

But the $\triangle MPE$ is less than the $\triangle KPE$,
the $\triangle HPF$ is less than the $\triangle KPE$:
to each add the fig. $AHPE$;
then the $\triangle FAE$ is less than the $\triangle HAK$.

Similarly it may be shewn that the $\triangle FAE$ is less than any other triangle formed by drawing a st. line through P :
that is, the $\triangle FAE$ is a minimum.

EXAMPLES.

1. Two sides of a triangle are given in length; how must they be placed in order that the area of the triangle may be a maximum?

2. Of all triangles of given base and area, the isosceles is that which has the least perimeter.

3. Given the base and vertical angle of a triangle; construct it so that its area may be a maximum.

4. Find a point in a given straight line such that the tangents drawn from it to a given circle contain the greatest angle possible.

5. A straight rod slips between two straight rulers placed at right angles to one another; in what position is the triangle intercepted between the rulers and rod a maximum?

6. Divide a given straight line into two parts, so that the sum of the squares on the segments may

- (i) be equal to a given square,
- (ii) may be a minimum.

7. Through a point of intersection of two circles draw a straight line terminated by the circumferences,

- (i) so that it may be of given length,
- (ii) so that it may be a maximum.

8. Two tangents to a circle cut one another at right angles; find the point on the intercepted arc such that the sum of the perpendiculars drawn from it to the tangents may be a minimum.

9. Straight lines are drawn from two given points to meet one another on the convex circumference of a given circle: prove that their sum is a minimum when they make equal angles with the tangent at the point of intersection.

10. Of all triangles of given vertical angle and altitude, the isosceles is that which has the least area.

11. Two straight lines CA , CB of indefinite length are drawn from the centre of a circle to meet the circumference at A and B ; then of all tangents that may be drawn to the circle at points on the arc AB , that whose intercept is bisected at the point of contact cuts off the triangle of minimum area.

12. Given two intersecting tangents to a circle, draw a tangent to the convex arc so that the triangle formed by it and the given tangents may be of maximum area.

13. Of all triangles of given base and area, the isosceles is that which has the greatest vertical angle.

14. Find a point on the circumference of a circle at which the straight line joining two given points (of which both are within, or both without the circle) subtends the greatest angle.

15. A bridge consists of three arches, whose spans are 49 ft., 32 ft. and 49 ft. respectively: shew that the point on either bank of the river at which the middle arch subtends the greatest angle is 63 feet distant from the bridge.

16. From a given point P without a circle whose centre is C , draw a straight line to cut the circumference at A and B , so that the triangle ACB may be of maximum area.

17. Shew that the greatest rectangle which can be inscribed in a circle is a square.

18. A and B are two fixed points without a circle: find a point P on the circumference such that the sum of the squares on AP , PB may be a minimum. [See p. 147, Ex. 24.]

5. Given an angle and the radii of the inscribed and circumscribed circles; construct the triangle.

6. Given the base, an angle at the base, and the distance between the centre of the inscribed circle and the centre of the escribed circle which touches the base; construct the triangle.

7. In a given circle inscribe a triangle such that two of its sides may pass through two given points, and the third side be of given length.

8. In any triangle ABC , I, I_1, I_2, I_3 are the centres of the inscribed and escribed circles, and S_1, S_2, S_3 are the centres of the circles circumscribed about the triangles BIC, CIA, AIB : shew that the triangle $S_1S_2S_3$ has its sides parallel to those of the triangle $I_1I_2I_3$, and is one-fourth of it in area: also that the triangles ABC and $S_1S_2S_3$ have the same circumscribed circle.

9. O is the orthocentre of a triangle ABC : shew that

$$AO^2 + BC^2 = BO^2 + CA^2 = CO^2 + AB^2 = d^2,$$

where d is the diameter of the circumscribed circle.

10. If from any point within a regular polygon of n sides perpendiculars are drawn to the sides the sum of the perpendiculars is equal to n times the radius of the inscribed circle.

11. The sum of the perpendiculars drawn from the vertices of a regular polygon of n sides on any straight line is equal to n times the perpendicular drawn from the centre of the inscribed circle.

12. The area of a cyclic quadrilateral is independent of the order in which the sides are placed in the circle.

13. Given the orthocentre, the centre of the nine points circle, and the middle point of the base: construct the triangle.

14. Of all polygons of a given number of sides, which may be inscribed in a given circle, that which is regular has the maximum area and the maximum perimeter.

15. Of all polygons of a given number of sides circumscribed about a given circle, that which is regular has the minimum area and the minimum perimeter.

16. Given the vertical angle of a triangle in position and magnitude, and the sum of the sides containing it: find the locus of the centre of the circumscribed circle.

17. P is any point on the circumference of a circle circumscribed about an equilateral triangle ABC : shew that $PA^2 + PB^2 + PC^2$ is constant.

7. PQ is a fixed chord in a circle, and PX , QY any two parallel chords through P and Q : shew that XY touches a fixed concentric circle.

8. Two equal circles intersect at A and B ; and from C any point on the circumference of one of them a perpendicular is drawn to AB , meeting the other circle at O and O' : shew that either O or O' is the orthocentre of the triangle ABC . Distinguish between the two cases.

9. Three equal circles pass through the same point A , and their other points of intersection are B , C , D . shew that of the four points A , B , C , D , each is the orthocentre of the triangle formed by joining the other three.

10. From a given point without a circle draw a straight line to the concave circumference so as to be bisected by the convex circumference. When is this problem impossible?

11. Draw a straight line cutting two concentric circles so that the chord intercepted by the circumference of the greater circle may be double of the chord intercepted by the less.

12. ABC is a triangle inscribed in a circle, and A' , B' , C' are the middle points of the arcs subtended by the sides (remote from the opposite vertices): find the relation between the angles of the two triangles ABC , $A'B'C'$; and prove that the pedal triangle of $A'B'C'$ is equilateral to the triangle ABC .

13. The opposite sides of a quadrilateral inscribed in a circle are produced to meet: shew that the bisectors of the two angles so formed are perpendicular to one another.

14. If a quadrilateral can have one circle inscribed in it, and another circumscribed about it; shew that the straight lines joining the opposite points of contact of the inscribed circle are perpendicular to one another.

15. Given the base of a triangle and the sum of the remaining sides; find the locus of the foot of the perpendicular from one extremity of the base on the bisector of the exterior vertical angle.

16. Two circles touch each other at C , and straight lines are drawn through C at right angles to one another, meeting the circles at P , P' and Q , Q' respectively: if the straight line which joins the centres is terminated by the circumferences at A and A' , shew that

$$P'P^2 + Q'Q^2 = A'A^2.$$

17. Two circles cut one another orthogonally at A and B ; P is any point on the arc of one circle intercepted by the other, and PA , PB are produced to meet the circumference of the second circle at C and D : shew that CD is a diameter.

18. $\triangle ABC$ is a triangle, and from any point P perpendiculars PD , PE , PF are drawn to the sides: if S_1 , S_2 , S_3 are the centres of the circles circumscribed about the triangles EPF , FPD , DPE , shew that the triangle $S_1S_2S_3$ is equiangular to the triangle ABC , and that the sides of the one are respectively half of the sides of the other.

19. Two tangents PA , PB are drawn from an external point P to a given circle, and C is the middle point of the chord of contact AB : if XY is any chord through P , shew that AB bisects the angle XPY .

20. Given the sum of two straight lines and the rectangle contained by them (equal to a given square) find the lines.

21. Given the sum of the squares on two straight lines and the rectangle contained by them: find the lines.

22. Given the sum of two straight lines and the sum of the squares on them: find the lines.

23. Given the difference between two straight lines, and the rectangle contained by them: find the lines.

24. Given the sum or difference of two straight lines and the difference of their squares: find the lines.

25. ABC is a triangle, and the internal and external bisectors of the angle A meet BC , and BC produced, at P and P' : if O is the middle point of PP' , shew that OA is a tangent to the circle circumscribed about the triangle ABC .

26. ABC is a triangle, and from P , any point on the circumference of the circle circumscribed about it, perpendiculars are drawn to the sides BC , CA , AB meeting the circle again in A' , B' , C' ; prove that

- (i) the triangle $A'B'C'$ is identically equal to the triangle ABC .
- (ii) AA' , BB' , CC' are parallel.

27. Two equal circles intersect at fixed points A and B , and from any point in AB a perpendicular is drawn to meet the circumferences on the same side of AB at P and Q : shew that PQ is of constant length.

28. The straight lines which join the vertices of a triangle to the centre of its circumscribed circle, are perpendicular respectively to the sides of the pedal triangle.

29. P is any point on the circumference of a circle circumscribed about a triangle ABC ; and perpendiculars PD , PE are drawn from P to the sides BC , CA . Find the locus of the centre of the circle circumscribed about the triangle PDE .

30. P is any point on the circumference of a circle circumscribed about a triangle ABC : shew that the angle between Simson's line for the point P and the side BC , is equal to the angle between AP and the diameter of the circumscribed circle through A .

31. Shew that the circles circumscribed about the four triangles formed by two pairs of intersecting straight lines meet in a point.

32. Shew that the orthocentres of the four triangles formed by two pairs of intersecting straight lines are collinear.

33. Given the vertical angle, one of the sides containing it, and the length of the perpendicular from the vertex on the base: construct the triangle.

34. Given the feet of the perpendiculars drawn from the vertices on the opposite sides: construct the triangle.

35. Given the base, the altitude, and the radius of the circumscribed circle: construct the triangle.

36. Given the base, the vertical angle, and the sum of the squares on the sides containing the vertical angle: construct the triangle.

37. Given the base, the altitude and the sum of the squares on the sides containing the vertical angle: construct the triangle.

38. Given the base, the vertical angle, and the difference of the squares on the sides containing the vertical angle: construct the triangle.

39. Given the vertical angle, and the lengths of the two medians drawn from the extremities of the base: construct the triangle.

40. Given the base, the vertical angle, and the difference of the angles at the base: construct the triangle.

41. Given the base, and the position of the bisector of the vertical angle: construct the triangle.

42. Given the base, the vertical angle, and the length of the bisector of the vertical angle: construct the triangle.

43. Given the perpendicular from the vertex on the base, the bisector of the vertical angle, and the median which bisects the base: construct the triangle.

44. Given the bisector of the vertical angle, the median bisecting the base, and the difference of the angles at the base: construct the triangle.

BOOK IV.

Book IV. ~~is~~ ^{consists} entirely of problems, dealing with various rectilinear ~~figures~~ ^{figures} in relation to the circles which pass through their ~~or~~ ^{or} points, or are touched by their sides.

DEFINITIONS.

1. A **Polygon** is a rectilinear figure bounded by more than four sides.


A Polygon of	<i>five</i> sides	is called a	Pentagon,
"	<i>six</i> sides	"	Hexagon,
"	<i>seven</i> sides	"	Heptagon,
"	<i>eight</i> sides	"	Octagon,
"	<i>ten</i> sides	"	Decagon,
"	<i>twelve</i> sides	"	Dodecagon,
"	<i>fifteen</i> sides	"	Quindecagon.

2. A Polygon is **Regular** when all its sides are equal, and all its angles are equal.

3. A rectilinear figure is said to be **inscribed** in a circle, when all its angular points are on the circumference of the circle: and a circle is said to be **circumscribed about** a rectilinear figure, when the circumference of the circle passes through all the angular points of the figure.




4. A rectilinear figure is said to be **circumscribed about** a circle, when each side of the figure is a tangent to the circle: and a circle is said to be **inscribed in** a rectilinear figure, when the circumference of the circle is touched by each side of the figure.



5. A straight line is said to be **placed in** a circle, when its extremities are on the circumference of the circle.

PROPOSITION I. PROBLEM.

In a given circle to place a chord equal to a given straight line, which is not greater than the diameter of the circle.



Let ABC be the given circle, and D the given straight line not greater than the diameter of the circle :

it is required to place in the $\odot ABC$ a chord equal to D .

Draw CB , a diameter of the $\odot ABC$.

Then if $CB = D$, the thing required is done.

But if not, CB must be greater than D . *Hyp.*

From CB cut off CE equal to D : 1. 3.

and from centre C , with radius CE , describe the $\odot AEF$,
cutting the given circle at A .

Join CA .

Then CA shall be the chord required.

For $CA = CE$, being radii of the $\odot AEF$.

and $CE = D$.

$\therefore CA = D$.

Constr.

$\odot E, F.$

EXERCISES.


1. In a given circle place a chord of given length so as to pass through a given point (i) without, (ii) within the circle.

When is this problem impossible?

2. In a given circle place a chord of given length so that it may be parallel to a given straight line.

PROPOSITION 2. PROBLEM.

In a given circle to inscribe a triangle equiangular to a given triangle.



Let ABG be the given circle, and DEF the given triangle. it is required to inscribe in the $\odot ABC$ a triangle equiangular to the $\triangle DEF$.

At any point A , on the \odot of the $\odot ABC$, draw the tangent GAH . III. 17.

At A make the $\angle GAB$ equal to the $\angle DFE$; I. 23.
and make the $\angle HAC$ equal to the $\angle DEF$. I. 23.

Join BC .

Then ABC shall be the triangle required.

Because GH is a tangent to the $\odot ABC$, and from A its point of contact the chord AB is drawn,

\therefore the $\angle GAB =$ the $\angle ACB$ in the alt. segment: III. 32.

\therefore the $\angle ACB =$ the $\angle DFE$. Constr.

Similarly the $\angle HAC =$ the $\angle ABC$, in the alt. segment:

\therefore the $\angle ABC =$ the $\angle DEF$. Constr.



Hence the third $\angle BAC =$ the third $\angle EDF$,
for the three angles in each triangle are together equal to two rt. angles. I. 32.

\therefore the $\triangle ABC$ is equiangular to the $\triangle DEF$, and it is inscribed in the $\odot ABC$.

Q. E. F.

PROPOSITION 3. PROBLEM.

About a given circle to circumscribe a triangle equiangular to a given triangle.



Let ABC be the given circle, and DEF the given \triangle : it is required to circumscribe about the $\odot ABC$ a triangle equiangular to the $\triangle DEF$.

Produce EF both ways to G and H .

Find K the centre of the $\odot ABC$, and draw any radius KB . III. 1.

At K make the $\angle BKA$ equal to the $\angle DEG$; and make the $\angle BKC$ equal to the $\angle DFH$. I. 23.

Through A, B, C draw LM, MN, NL perp. to KA, KB, KC .

Then LMN shall be the triangle required.

Because LM, MN, NL are drawn perp. to radii at their extremities,

$\therefore LM, MN, NL$ are tangents to the circle. III. 16.

And because the four angles of the quadrilateral $AKBM$ together, four rt. angles; I. 32. Cor.

and of these, the $\angle KAM, KBM$, are rt. angles; Constr.

\therefore the $\angle AKB, AMB$, together = two rt. angles.

But the $\angle DEG, DEF$ together = two rt. angles; I. 13.

\therefore the $\angle AKB, AMB$ = the $\angle DEG, DEF$;

and of these, the $\angle AKB$ = the $\angle DEG$; Constr.

\therefore the $\angle AMB$ = the $\angle DEF$.


Similarly it may be shewn that the $\angle LNM$ = the $\angle DFE$.

\therefore the third $\angle MLN$ = the third $\angle EDF$. I. 32.

\therefore the $\triangle LMN$ is equiangular to the $\triangle DEF$, and it is circumscribed about the $\odot ABC$. Q.E.F.

PROPOSITION 4. PROBLEM.

To inscribe a circle in a given triangle.



Let $\triangle ABC$ be the given triangle :
it is required to inscribe a circle in the $\triangle ABC$.

Bisect the \angle^s ABC , ACB by the st. lines BI , CI , which intersect at I . I. 9.

From I draw IE , IF , IG perp. to AB , BC , CA : I. 12.

Then in the \triangle^s EIB , FIB ,
Because $\left\{ \begin{array}{l} \text{the } \angle EBI = \text{the } \angle FBI ; \\ \text{and the } \angle BEI = \text{the } \angle BFI, \text{ being rt. angles ;} \\ \text{and } BI \text{ is common ;} \end{array} \right. \quad \begin{array}{l} \text{Constr.} \\ \\ \end{array}$
 $\therefore IE = IF$. I. 26.

Similarly it may be shewn that $IF = IG$.

$\therefore IE$, IF , IG are all equal.

From centre I , with radius IE , describe a circle :
this circle must pass through the points E , F , G ;
and it will be inscribed in the $\triangle ABC$.

For since IE , IF , IG are radii of the $\odot EFG$;
and since the \angle^s at E , F , G are rt. angles ;
the $\odot EFG$ is touched at these points by AB , BC , CA :
III. 16.

\therefore the $\odot EFG$ is inscribed in the $\triangle ABC$.

Q. E. F.

Now. From page 103 it is seen that if AI be joined, then AI bisects the angle BAC .

Hence it follows that the bisectors of the angles of a triangle are concurrent, the point of intersection being the centre of the inscribed circle.

The centre of the circle inscribed in a triangle is sometimes called its **in-centre**.

DEFINITION.

A circle which touches one side of a triangle and the other two sides produced is said to be an **escribed circle** of the triangle.

To draw an escribed circle of a given triangle.

Let ABC be the given triangle, of which the two sides AB , AC are produced to E and F :

it is required to describe a circle touching BC , and AB , AC produced.

Bisect the $\angle CBE$, BCF by the st. lines BI_1 , CI_1 , which intersect at I_1 . I. 9.

From I_1 draw I_1G , I_1H , I_1K perp. to AE , BC , AF . I. 12.

Then in the ΔI_1BG , I_1BH ,
Because $\left\{ \begin{array}{l} \text{the } \angle I_1BG = \text{the } \angle I_1BH, \text{ Constr.} \\ \text{and the } \angle I_1GB = \text{the } \angle I_1HB, \\ \text{being rt. angles;} \\ \text{also } I_1B \text{ is common;} \end{array} \right.$
 $\therefore I_1G = I_1H$.

Similarly it may be shewn that $I_1H = I_1K$;
 $\therefore I_1G$, I_1H , I_1K are all equal.

From centre I_1 with radius I_1G , describe a circle:

this circle must pass through the points G , H , K ;
and it will be an escribed circle of the ΔABC .


For since I_1H , I_1G , I_1K are radii of the $\odot HGK$,
and since the angles at H , G , K are rt. angles,
the $\odot GHK$ is touched at these points by BC , and by AB , AC
produced:

\therefore the $\odot GHK$ is an escribed circle of the ΔABC . Q.E.F.

It is clear that every triangle has *three* escribed circles.


NOTE. From page 104 it is seen that if AI_1 be joined, then AI_1 bisects the angle BAC : hence it follows that

The bisectors of two exterior angles of a triangle and the bisector of the third angle are concurrent, the point of intersection being the centre of an escribed circle.



PROPOSITION 5. PROBLEM.

To circumscribe a circle about a given triangle.



Let ABC be the given triangle ;
it is required to circumscribe a circle about the $\triangle ABC$.

Draw DS bisecting AB at rt. angles ; i. 11.
and draw ES bisecting AC at rt. angles,
then since AB, AC are neither par^l, nor in the same st. line,
 $\therefore DS$ and ES must meet at some point S .

Join SA ;

and, if S be not in BC , join SB, SC ;

Then in the $\triangle ADS, BDS$,

Because $\left\{ \begin{array}{l} AD = BD \\ \text{and } DS \text{ is common to both ;} \\ \text{and the } \angle ADS = \text{the } \angle BDS, \text{ being rt. angles ;} \end{array} \right.$
 $\therefore SA = SB$.

Similarly it may be shewn that $SC = SA$.

$\therefore SA, SB, SC$ are all equal.

From centre S , with radius SA , describe a circle :
this circle must pass through the points A, B, C , and is
therefore circumscribed about the $\triangle ABC$. Q. E. F.

It follows that

(i) when the centre of the circumscribed circle falls *within* the triangle, each of its angles must be acute, for each angle is then in a segment greater than a semicircle :

(ii) when the centre falls *on one of the sides* of the triangle, the angle opposite to this side must be a right angle, for it is the angle in a semicircle :

(iii) when the centre falls *without* the triangle, the angle opposite to the side beyond which the centre falls, must be obtuse, for it is the angle in a segment less than a semicircle.

Therefore, conversely, if the given triangle be acute angled, the centre of the circumscribed circle falls within it: if it be a right-angled triangle, the centre falls on the hypotenuse: if it be an obtuse-angled triangle, the centre falls without the triangle.

NOTE. From page 103 it is seen that if S be joined to the middle point of BC , then the joining line is perpendicular to BC .

Hence the perpendiculars drawn to the sides of a triangle from their middle points are concurrent, the point of intersection being the centre of the circle circumscribed about the triangle.

The centre of the circle circumscribed about a triangle is sometimes called its **circum-centre**.

EXERCISES.

ON THE INSCRIBED, CIRCUMSCRIBED, AND EScribed CIRCLES OF A TRIANGLE.

1. An equilateral triangle is inscribed in a circle, and tangents are drawn at its vertices, prove that

- (i) the resulting figure is an equilateral triangle;
- (ii) its area is four times that of the given triangle.

2. Describe a circle to touch two parallel straight lines and a third straight line which meets them. Shew that two such circles can be drawn, and that they are equal.

3. Triangles which have equal bases and equal vertical angles have equal circumscribed circles.

4. I is the centre of the circle inscribed in the triangle ABC , and I_1 is the centre of the circle which touches BC and AB, AC produced: shew that A, I, I_1 are collinear.

5. If the inscribed and circumscribed circles of a triangle are concentric, shew that the triangle is equilateral; and that the diameter of the circumscribed circle is double that of the inscribed circle.

6. ABC is a triangle; and I, S are the centres of the inscribed and circumscribed circles; if A, I, S are collinear, shew that $AB = AC$.

7. The sum of the diameters of the inscribed and circumscribed circles of a right-angled triangle is equal to the sum of the sides containing the right angle.

8. If the circle inscribed in a triangle ABC touches the sides at D, E, F , shew that the triangle DEF is acute-angled; and express its angles in terms of the angles at A, B, C .

9. If I is the centre of the circle inscribed in the triangle ABC , and I_1 the centre of the escribed circle which touches BC ; shew that I, B, I_1, C are concyclic.

10. In any triangle the difference of two sides is equal to the difference of the segments into which the third side is divided at the point of contact of the inscribed circle.

11. In the triangle ABC the bisector of the angle BAC meets the base at D , and from I the centre of the inscribed circle a perpendicular IE is drawn to BC ; shew that the angle BID is equal to the angle CIE .

12. In the triangle ABC , I and S are the centres of the inscribed and circumscribed circles: shew that IS subtends at A an angle equal to half the difference of the angles at the base of the triangle.

13. In a triangle ABC , I and S are the centres of the inscribed and circumscribed circles, and AD is drawn perpendicular to BC : shew that AI is the bisector of the angle DAS .

14. Shew that the area of a triangle is equal to the rectangle contained by its semi-perimeter and the radius of the inscribed circle.

15. The diagonals of a quadrilateral $ABCD$ intersect at O : shew that the centres of the circles circumscribed about the four triangles AOB, BOC, COD, DOA are at the angular points of a parallelogram.

16. In any triangle ABC , if I is the centre of the inscribed circle, and if AI is produced to meet the circumscribed circle at O ; shew that O is the centre of the circle circumscribed about the triangle BIC .

17. Given the base, altitude, and the radius of the circumscribed circle; construct the triangle.


18. Describe a circle to intercept equal chords of given length on three given straight lines.

19. In an equilateral triangle the radii of the circumscribed and escribed circles are respectively double and treble of the radius of the inscribed circle.

20. Three circles whose centres are A, B, C touch one another externally two by two at D, E, F : shew that the inscribed circle of the triangle ABC is the circumscribed circle of the triangle DEF .

PROPOSITION 6. PROBLEM.

To inscribe a square in a given circle.



Let ABCD be the given circle.
it is required to inscribe a square in the \odot ABCD.

Find E the centre of the circle : m. 1.
and draw two diameters AC, BD perp. to one another. r. 11.
Join AB, BC, CD, DA.

Then the fig. ABCD shall be the square required.

For in the Δ 's BEA, DEA,
Because $\left\{ \begin{array}{l} BE = DE, \\ \text{and } EA \text{ is common;} \\ \text{and the } \angle BEA = \text{the } \angle DEA, \text{ being rt. angles;} \end{array} \right.$
 $\therefore BA = DA.$ l. 4.

Similarly it may be shewn that $CD = DA$, and that $BC = CD$.
 \therefore the fig. ABCD is equilateral.


And since BD is a diameter of the \odot ABCD,
 \therefore BAD is a semicircle;
 \therefore the \angle BAD is a rt. angle. m. 31.

Similarly the other angles of the fig. ABCD are rt. angles.
 \therefore the fig. ABCD is a square,
and it is inscribed in the given circle.

Q. E. D.

PROPOSITION 7. PROBLEM.

To circumscribe a square about a given circle.



Let $\odot ABCD$ be the given circle.
it is required to circumscribe a square about it.

Find E the centre of the $\odot ABCD$: iii. 1.
and draw two diameters AC, BD perp. to one another. i. 11.
Through A, B, C, D draw FG, GH, HK, KF perp. to EA, EB, EC, ED .

Then the fig. GK shall be the square required.

Because FG, GH, HK, KF are drawn perp. to radii at their extremities,

$\therefore FG, GH, HK, KF$ are tangents to the circle. iii. 16

And because the \angle 's AEB, EBG are both rt. angles, Constr.

$\therefore GH$ is par^l to AC . i. 28.

Similarly FK is par^l to AC :

and in like manner GF, BD, HK are par^l.

Hence the figs. GK, GC, AK, GD, BK, GE are par^{ms}.

$\therefore GF$ and HK each = BD ;

also GH and FK each = AC :

but $AC = BD$;

$\therefore GF, FK, KH, HG$ are all equal
that is, the fig. GK is equilateral.

And since the fig. GE is a par^m,

\therefore the \angle $EGA =$ the \angle BEA ;

i. 31.

but the \angle BEA is a rt. angle .

Constr.

\therefore the \angle at G is a rt. angle


Similarly the \angle 's at F, K, H are rt. angles.

\therefore the fig. GK is a square, and it has been circumscribed about the $\odot ABCD$.

Q. E. F.

PROPOSITION 8. PROBLEM.

To inscribe a circle in a given square.



Let $ABCD$ be the given square.
it is required to inscribe a circle in the sq. $ABCD$.

Bisect the sides AB , AD at F and E . I. 10.

Through E draw EH par^l to AB or DC : I. 31.
and through F draw FK par^l to AD or BC , meeting EH at G .

Now $AB = AD$, being the sides of a square;
and their halves are equal; Constr.

$\therefore AF = AE$. I. 7.

But the fig. AG is a par^m; Constr.

$\therefore AF = GE$, and $AE = GF$;

$\therefore GE = GF$.

Similarly it may be shown that $GE = GK$, and $GK = GH$:

$\therefore GF, GE, GK, GH$ are all equal.

From centre G , with radius GE , describe a circle;

this circle must pass through the points F, E, K, H :

and it will be touched by BA, AD, DC, CB ; III. 16.

for GF, GE, GK, GH are radii;


and the angles at F, E, K, H are rt. angles. I. 29.

Hence the $\odot FEKH$ is inscribed in the sq. $ABCD$.

Q. E. F.

PROPOSITION 9. PROBLEM.

To circumscribe a circle about a given square.



Let $ABCD$ be the given square :
it is required to circumscribe a circle about the sq. $ABCD$.

Join AC , BD , intersecting at E .

Then in the Δ^s BAC , DAC ,

Because $\begin{cases} BA = DA, & \text{I. Def. 28.} \\ \text{and } AC \text{ is common;} \\ \text{and } BC = DC; & \text{I. Def. 28.} \end{cases}$

\therefore the $\angle BAC =$ the $\angle DAC$: I. 8.

that is, the diagonal AC bisects the $\angle BAD$.

Similarly the remaining angles of the square are bisected by the diagonals AC or BD .

Hence each of the \angle^s EAD , EDA is half a rt. angle,

\therefore the $\angle EAD =$ the $\angle EDA$:

$\therefore EA = ED$. I. 6.

Similarly it may be shewn that $ED = EC$, and $EC = EB$.

$\therefore EA, EB, EC, ED$ are all equal.

From centre E , with radius EA , describe a circle :
this circle must pass through the points A, B, C, D , and is
therefore circumscribed about the sq. $ABCD$. Q. E. F.

DEFINITION. A rectilineal figure about which a circle may be described is said to be **Cyclic**.

EXERCISES ON PROPOSITIONS 6 - 9.

1. If a circle can be inscribed in a quadrilateral, shew that the sum of one pair of opposite sides is equal to the sum of the other pair.

2. If the sum of one pair of opposite sides of a quadrilateral is equal to the sum of the other pair, shew that a circle may be inscribed in the figure.

[Bisect two adjacent angles of the figure, and so describe a circle to touch three of its sides. Then prove indirectly by means of the last exercise that this circle must also touch the fourth side.]

3. Prove that a rhombus and a square are the only parallelograms in which a circle can be inscribed.

4. All cyclic parallelograms are rectangular.

5. The greatest rectangle which can be inscribed in a given circle is a square.

6. Circumscribe a rhombus about a given circle.

7. All squares circumscribed about a given circle are equal.

8. The area of a square circumscribed about a circle is double of the area of the inscribed square.

9. $ABCD$ is a square inscribed in a circle, and P is any point on the arc AD . shew that the side AD subtends at P an angle three times as great as that subtended at P by any one of the other sides.

10. Inscribe a square in a given square $ABCD$ so that one of its angular points should be at a given point X in AB .

11. In a given square inscribe the square of minimum area.


12. Describe (i) a circle, (ii) a square about a given rectangle.

13. Inscribe (i) a circle, (ii) a square in a given quadrant.

14. $ABCD$ is a square inscribed in a circle, and P is any point on the circumference; shew that the sum of the squares on PA , PB , PC , PD is double the square on the diameter. [See Ex. 24, p. 147.]

PROPOSITION 10. PROBLEM.

To describe an isosceles triangle having each of the angles at the base double of the third angle.



Take any straight line AB.

Divide AB at C, so that the rect. BA, BC = the sq. on AC.

II. 11.

From centre A, with radius AB, describe the \odot BDE;

and in it place the chord BD equal to AC.

IV. 1.

Join DA.

Then ABD shall be the triangle required.

Join CD;

and about the Δ ACD circumscribe a circle. IV. 5.

Then the rect. BA, BC = the sq. on AC Constr.

= the sq. on BD. Constr.

Hence BD is a tangent to the \odot ACD: III. 37.

and from the point of contact D a chord DC is drawn;

\therefore the \angle BDC = the \angle CAD in the alt. segment. III. 32.

To each of these equals add the \angle CDA:

then the whole \angle BDA = the sum of the \angle s CAD, CDA.

But the ext. \angle BCD = the sum of the \angle s CAD, CDA; I. 32.

\therefore the \angle BCD = the \angle BDA.

And since AB = AC, being radii of the \odot BDE,

\therefore the \angle DBA = the \angle BDA: I. 5.

\therefore the \angle DBC = the \angle DCB;

$\therefore DC = DB$; I. 6.
 that is, $DC = CA$; *Constr.*
 \therefore the $\angle CAD =$ the $\angle CDA$, I. 5.
 \therefore the sum of the $\angle^s CAD, CDA =$ twice the angle at A.
 But the $\angle ADB =$ the sum of the $\angle^s CAD, CDA$; *Proved.*
 \therefore each of the $\angle^s ABD, ADB =$ twice the angle at A.
Q. E. F.

EXERCISES ON PROPOSITION 10

1. In an isosceles triangle in which each of the angles at the base is double of the vertical angle, shew that the vertical angle is one-fifth of two right angles.

2. Divide a right angle into five equal parts.

3. Describe an isosceles triangle whose vertical angle shall be three times either angle at the base. Point out a triangle of this kind in the figure of Proposition 10.

4. In the figure of Proposition 10, if the two circles intersect at F, shew that $BD = DF$.

5. In the figure of Proposition 10, shew that the circle ACD is equal to the circle circumscribed about the triangle ABD.

6. In the figure of Proposition 10, if the two circles intersect at F, shew that

(i) BD, DF are sides of a regular decagon inscribed in the circle EBD.


(ii) AC, CD, DF are sides of a regular pentagon inscribed in the circle ACD.

7. In the figure of Proposition 10, shew that the centre of the circle circumscribed about the triangle DBC is the middle point of the arc CD.

8. In the figure of Proposition 10, if I is the centre of the circle inscribed in the triangle ABD, and I', S' the centres of the inscribed and circumscribed circles of the triangle DBC, shew that $S'I = S'I'$.

PROPOSITION 11. PROBLEM.

To inscribe a regular pentagon in a given circle.



Let $\odot ABC$ be a given circle :
it is required to inscribe a regular pentagon in the $\odot ABC$.

Describe an isosceles $\triangle FGH$, having each of the angles at G and H double of the angle at F . IV. 10.

In the $\odot ABC$ inscribe the $\triangle ACD$ equiangular to the $\triangle FGH$, IV. 2.
so that each of the $\angle^s ACD, ADC$ is double of the $\angle CAD$.

Bisect the $\angle^s ACD, ADC$ by CE and DB , which meet the \odot at E and B . I. 9.

Join AB, BC, AE, ED .

Then $ABCDE$ shall be the required regular pentagon.

Because each of the $\angle^s ACD, ADC$ twice the $\angle CAD$;
and because the $\angle^s ACD, ADC$ are bisected by CE, DB ,
 \therefore the five $\angle^s ADB, BDC, CAD, DCE, ECA$ are all equal.
 \therefore the five arcs $\widehat{AB}, \widehat{BC}, \widehat{CD}, \widehat{DE}, \widehat{EA}$ are all equal. III. 26.
 \therefore the five chords AB, BC, CD, DE, EA are all equal. III. 29.
 \therefore the pentagon $ABCDE$ is equilateral.

Again the arc \widehat{AB} = the arc \widehat{DE} ; Proved.
to each of these equals add the arc \widehat{BCD} ;
 \therefore the whole arc \widehat{ABCD} = the whole arc \widehat{BCDE} :

\therefore hence the angles at the \odot which stand upon these equal arcs are equal ; III. 27.
that is, the $\angle AED$ = the $\angle BAE$.


In like manner the remaining angles of the pentagon may be shewn to be equal ;

\therefore the pentagon is equiangular.

Hence the pentagon, being both equilateral and equiangular, is regular ; and it is inscribed in the $\odot ABC$. Q. E. F.

PROPOSITION 12 PROBLEM.

To circumscribe a regular pentagon about a given circle.



Let $ABCD$ be the given circle :
it is required to circumscribe a regular pentagon about it.

Inscribe a regular pentagon in the $\odot ABCD$, iv. 11.
and let A, B, C, D, E be its angular points.

At the points A, B, C, D, E draw GH, HK, KL, LM, MG ,
tangents to the circle. iii. 17.

Then shall $GHKLM$ be the required regular pentagon.

Find F the centre of the $\odot ABCD$, iii. 1

and join FB, FK, FC, FL, FD .

Then in the two $\triangle BFK, CFK$,

Because $\left\{ \begin{array}{l} BF = CF, \text{ being radii of the circle,} \\ \text{and } FK \text{ is common :} \\ \text{and } KB = KC, \text{ being tangents to the circle from} \\ \text{the same point } K. \end{array} \right.$ iii. 17. Cor.

\therefore the $\angle BFK =$ the $\angle CFK$, i. 8.

also the $\angle BKF =$ the $\angle CKF$. i. 8. Cor.

Hence the $\angle BFC =$ twice the $\angle CFK$,

and the $\angle BKC =$ twice the $\angle CKF$.

Similarly it may be shewn

that the $\angle CFD =$ twice the $\angle CFL$,


and that the $\angle CLD =$ twice the $\angle CLF$.

But since the arc $BC =$ the arc CD , iv. 11.

\therefore the $\angle BFC =$ the $\angle CFD$; iii. 27

and the halves of these angles are equal,

that is, the $\angle CFK =$ the $\angle CFL$.



Then in the \triangle s CFK, CFL,
 the \angle CFK = the \angle CFL, *Proved.*
 Because { and the \angle FCK = the \angle FCL, being rt. angles, III. 18.
 and FC is common :
 \therefore CK = CL, I. 26.
 and the \angle FKC = the \angle FLC.

Hence KL is double of KC ; similarly HK is double of KB.

And since KC = KB, III. 17. Cor.

\therefore KL = HK.

In the same way it may be shewn that every two consecutive sides are equal ;

\therefore the pentagon GHKLM is equilateral.

Again it has been proved that the \angle FKC = the \angle FLC,
 and that the \angle s HKL, KLM are respectively double of these angles :

\therefore the \angle HKL = the \angle KLM.

In the same way it may be shewn that every two consecutive angles of the figure are equal ;


\therefore the pentagon GHKLM is equiangular.

\therefore the pentagon is regular, and it is circumscribed about the \odot ABCD. Q. E. F.

* COROLLARY. Similarly it may be proved that if tangents are drawn at the vertices of any regular polygon inscribed in a circle, they will form another regular polygon of the same species circumscribed about the circle.

PROPOSITION 13. PROBLEM.

To inscribe a circle in a given regular pentagon.



Let $ABCDE$ be the given regular pentagon :
it is required to inscribe a circle within it.

Bisect two consecutive \angle^s BCD , CDE by CF and DF
which intersect at F . I. 9.

Join FB ;

and draw FH , FK perp. to BC , CD . I. 12.

Then in the \triangle^s BCF , DCF ,


Because $\left\{ \begin{array}{l} BC = DC, \\ \text{and } CF \text{ is common to both,} \\ \text{and the } \angle BCF = \text{the } \angle DCF, \end{array} \right. \quad \begin{array}{l} \text{Hyp.} \\ \text{Constr.} \end{array}$
 \therefore the $\angle CBF =$ the $\angle CDF$. I. 4.

But the $\angle CDF$ is half an angle of the regular pentagon :
 \therefore also the $\angle CBF$ is half an angle of the regular pentagon :
that is, FB bisects the $\angle ABC$.

So it may be shewn that if FA , FE were joined, these
lines would bisect the \angle^s at A and E .

Again, in the \triangle^s FCH , FCK ,
Because $\left\{ \begin{array}{l} \text{the } \angle FCH = \text{the } \angle FCK, \\ \text{and the } \angle FHC = \text{the } \angle FKC \text{ being rt. angles;} \\ \text{also } FC \text{ is common;} \end{array} \right. \quad \text{Constr.}$
 $\therefore FH = FK$. I. 26.

Similarly if FG , FM , FL be drawn perp. to BA , AE , ED ,
it may be shewn that the five perpendiculars drawn from F
to the sides of the pentagon are all equal.



From centre F , with radius FH , describe a circle, this circle must pass through the points H, K, L, M, G ; and it will be touched at these points by the sides of the pentagon, for the \angle 's at H, K, L, M, G are rt \angle 's. *Constr.* \therefore the $\odot HKLMG$ is inscribed in the given pentagon. Q.E.D.

COROLLARY. *The bisectors of the angles of a regular pentagon meet at a point.*

In the same way it may be shewn that the bisectors of the angles of any regular polygon meet at a point. [See Ex. I, p. 271.]

[For Exercises on Regular Polygons see p. 276.]

MISCELLANEOUS EXERCISES.

1. Two tangents AB, AC are drawn from an external point A to a given circle: describe a circle to touch AB, AC and the convex arc intercepted by them on the given circle.


2. $\triangle ABC$ is an isosceles triangle, and from the vertex A a straight line is drawn to meet the base at D and the circumference of the circumscribed circle at E : shew that AB is a tangent to the circle circumscribed about the triangle BDE .

3. An equilateral triangle is inscribed in a given circle: shew that twice the square on one of its sides is equal to three times the area of the square inscribed in the same circle.

4. $\triangle ABC$ is an isosceles triangle in which each of the angles at B and C is double of the angle at A : shew that the square on AB is equal to the rectangle AB, BC with the square on BC .

PROPOSITION 14. PROBLEM.

To circumscribe a circle about a given regular pentagon.



Let $ABCDE$ be the given regular pentagon :
It is required to circumscribe a circle about it.

Bisect the \angle s BCD , CDE by CF , DF intersecting at F . I. 9.
Join FB , FA , FE .

Then in the \triangle s BCF , DCF ,

Because $\left\{ \begin{array}{l} BC = DC, \\ \text{and } CF \text{ is common to both;} \\ \text{and the } \angle BCF = \text{the } \angle DCF; \end{array} \right. \quad \begin{array}{l} \text{Hyp.} \\ \text{Constr.} \end{array}$
 $\therefore \text{the } \angle CBF = \text{the } \angle CDF.$ I. 4.

But the $\angle CDF$ is half an angle of the regular pentagon :
 \therefore also the $\angle CBF$ is half an angle of the regular pentagon :
that is, FB bisects the $\angle ABC$.

So it may be shewn that FA , FE bisect the \angle s at A and E .

Now the \angle s FCD , FDC are each half an angle of the given regular pentagon ;

$\therefore \text{the } \angle FCD = \text{the } \angle FDC,$ Iv. Def.
 $\therefore FC = FD.$ I. 6.

Similarly it may be shewn that FA , FB , FC , FD , FE are all equal.


From centre F , with radius FA describe a circle :
this circle must pass through the points A , B , C , D , E ,
and therefore is circumscribed about the given pentagon.

Q. E. D.

In the same way a circle may be circumscribed about any regular polygon.

PROPOSITION 15. PROBLEM.

To inscribe a regular hexagon in a given circle.



Let $ABDF$ be the given circle ;
it is required to inscribe a regular hexagon in it.

Find G the centre of the $\odot ABDF$; III. 1.
and draw a diameter AGD .

From centre D , with radius DG , describe the $\odot EGCH$.
Join CG, EG , and produce them to cut the \odot^{co} of the
given circle at F and B .

Join AB, BC, CD, DE, EF, FA .

Then $ABCDEF$ shall be the required regular hexagon.

Now $GE = GD$, being radii of the $\odot ACE$;
and $DG = DE$, being radii of the $\odot EHC$;

$\therefore GE, ED, DG$ are all equal, and the $\triangle EGD$ is equilateral.

Hence the $\angle EGD =$ one-third of two rt. angles. I. 32.

Similarly the $\angle DGC =$ one-third of two rt. angles.

But the $\angle^s EGD, DGC, CGB$ together = two rt. angles ; I. 13.

\therefore the remaining $\angle CGB =$ one third of two rt. angles.

\therefore the three $\angle^s EGD, DGC, CGB$ are equal to one another.

And to these angles the vert. opp. $\angle^s BGA, AGF, FGE$
are respectively equal :

\therefore the $\angle^s EGD, DGC, CGB, BGA, AGF, FGE$ are all equal ;

\therefore the arcs ED, DC, CB, BA, AF, FE are all equal ; III. 26.

\therefore the chords ED, DC, CB, BA, AF, FE are all equal : III. 29.

\therefore the hexagon is equilateral.

Again the arc $FA =$ the arc DE : *Proved.*

to each of these equals add the a.c. $ABCD$;

then the whole arc $FABCD =$ the whole arc $ABCDE$:

hence the angles at the \odot^{co} which stand on these equal arcs
are equal.

that is, the $\angle FED$ = the $\angle AFE$.

III. 27.

In like manner the remaining angles of the hexagon may be shewn to be equal.

\therefore the hexagon is equiangular :


\therefore the hexagon is regular, and it is inscribed in the \odot $ABDF$.

Q. E. F.

COROLLARY. *The side of a regular hexagon inscribed in a circle is equal to the radius of the circle.*

PROPOSITION 16. PROBLEM.

To inscribe a regular quindecagon in a given circle.



Let $ABCD$ be the given circle.

It is required to inscribe a regular quindecagon in it.

In the $\odot ABCD$ inscribe an equilateral triangle, IV. 2.
and let AC be one of its sides.

In the same circle inscribe a regular pentagon, IV. 11.
and let AB be one of its sides.

Then of such equal parts as the whole \odot contains fifteen,
the arc AC , which is one-third of the \odot , contains five;
and the arc AB , which is one-fifth of the \odot , contains three;
 \therefore their difference; the arc BC , contains two.

Bisect the arc BC at E : III. 30.

then each of the arcs BE , EC is one-fifteenth of the \odot .

\therefore if BE , EC be joined, and st. lines equal to them be placed successively round the circle, a regular quindecagon will be inscribed in it.

Q. E. F.

NOTE ON REGULAR POLYGONS.


The following propositions, proved by Euclid for a regular pentagon, hold good for all regular polygons.

1. *The bisectors of the angles of any regular polygon are concurrent.*

Let D, E, A, B, C be consecutive angular points of a regular polygon of any number of sides.

Bisect the \angle 's EAB, ABC by AO, BO, which intersect at O.

Join EO.



It is required to prove that EO bisects the \angle DEA.

For in the Δ 's EAO, BAO,
 Because $\left\{ \begin{array}{l} \text{EA} = \text{BA, being sides of a regular polygon,} \\ \text{and AO is common,} \\ \text{and the } \angle \text{EAO} = \text{the } \angle \text{BAO;} \end{array} \right. \quad \begin{array}{l} \text{Constr.} \\ 1. 4. \end{array}$
 $\therefore \text{the } \angle \text{OEA} = \text{the } \angle \text{OBA}.$

But the $\angle \text{OBA}$ is half the $\angle \text{ABC}$, Constr.
 also the $\angle \text{ABC} = \text{the } \angle \text{DEA}$, since the polygon is regular;
 $\therefore \text{the } \angle \text{OEA is half the } \angle \text{DEA};$
 that is, EO bisects the $\angle \text{DEA}.$

Similarly if O be joined to the remaining angular points of the polygon, it may be proved that each joining line bisects the angle to whose vertex it is drawn.

That is to say, the bisectors of the angles of the polygon meet at the point O. Q. E. D.

COROLLARIES. Since the $\angle \text{EAB} = \text{the } \angle \text{ABC};$ Hyp.
 and since the \angle 's OAB, OBA are respectively half of the \angle 's EAB, ABC;
 $\therefore \text{the } \angle \text{OAB} = \text{the } \angle \text{OBA}.$

$\therefore \text{OA} = \text{OB}.$ 1. 6.
 Similarly $\text{OE} = \text{OA}.$

Hence *The bisectors of the angles of a regular polygon are all equal; and a circle described from the centre O, with radius OA, will be circumscribed about the polygon.*

Also it may be shewn as in Proposition 13, that perpendiculars drawn from O to the sides of the polygon are all equal; therefore a circle described from centre O with any one of these perpendiculars as radius will be inscribed in the polygon.

2. If a polygon inscribed in a circle is equilateral, it is also equiangular.

Let AB, BC, CD be consecutive sides of an equilateral polygon inscribed in the $\odot ADK$;
then shall this polygon be equiangular.

Because the chord $AB =$ the chord DC , *Hyp.*
 \therefore the minor arc $AB =$ the minor arc DC , III. 28.

To each of these equals add the arc AKD ;

then the arc $BAKD =$ the arc $AKDC$;


\therefore the angles at the \odot , which stand on these equal arcs, are equal;

that is, the $\angle BCD =$ the $\angle ABC$. III. 27.

Similarly the remaining angles of the polygon may be shown to be equal:

the polygon is equiangular.



Q. E. D.



3. If a polygon inscribed in a circle is equiangular, it is also equilateral, provided that the number of its sides is odd.

[Observe that Theorems 2 and 3 are only true of polygons inscribed in a circle.]

The accompanying figures are sufficient to shew that otherwise a polygon may be equilateral without being equiangular, Fig. 1; or equiangular without being equilateral, Fig. 2.]



NOTE. The following extensions of Euclid's constructions for Regular Polygons should be noticed.

By continual bisection of arcs, it is enabled to divide the circumference of a circle,

by means of Proposition 6, into 4, 8, 16, \dots , $2 \cdot 2^n$, equal parts;

by means of Proposition 15, into 3, 6, 12, \dots , $3 \cdot 2^n$, equal parts;

by means of Proposition 11, into 5, 10, 20, \dots , $5 \cdot 2^n$, equal parts;

by means of Proposition 16, into 15, 30, 60, \dots , $15 \cdot 2^n$, equal parts.

Hence we can inscribe in a circle a regular polygon the number of whose sides is included in any one of the formulæ $2 \cdot 2^n$, $3 \cdot 2^n$, $5 \cdot 2^n$, $15 \cdot 2^n$, n being any positive integer. In addition to these, it has been shown that a regular polygon of $2^n + 1$ sides, provided $2^n + 1$ is a prime number, may be inscribed in a circle.

EXERCISES ON PROPOSITIONS 11--16.

1. Express in terms of a right angle the magnitude of an angle of the following *regular* polygons:

- (i) a pentagon, (ii) a hexagon, (iii) an octagon,
(iv) a decagon, (v) a quindecagon.

2. The angle of a regular pentagon is trisected by the straight line which join it to the opposite vertices.

3. In a polygon of n sides the straight lines which join any angular point to the vertices not adjacent to it, divide the angle into $n - 2$ equal parts.

4. Shew how to construct on a given straight line

- (i) a regular pentagon, (ii) a regular hexagon, (iii) a regular octagon.

5. An equilateral triangle and a regular hexagon are inscribed in a given circle; shew that

- (i) the area of the triangle is half that of the hexagon;
(ii) the square on the side of the triangle is three times the square on the side of the hexagon.

6. $ABCDE$ is a regular pentagon, and AC , BE intersect at H : shew that

- (i) $AB = CH = EH$.
(ii) AB is a tangent to the circle circumscribed about the triangle BHC .
(iii) AC and BE cut one another in medial section.

7. The straight lines which join alternate vertices of a regular pentagon intersect so as to form another regular pentagon.

8. The straight lines which join alternate vertices of a regular polygon of n sides, intersect so as to form another regular polygon of n sides.

If $n = 6$, shew that the area of the resulting hexagon is one-third of the given hexagon.

9. By means of iv. 16, inscribe in a circle a triangle whose angles are at the numbers 2, 5, 8.

10. Shew that the area of a regular hexagon inscribed in a circle is three-fourths of that of the corresponding circumscribed hexagon.

THEOREMS AND EXAMPLES ON BOOK IV.

I. ON THE TRIANGLE AND ITS CIRCLES.

1. D, E, F are the points of contact of the inscribed circle of the triangle ABC , and D_1, E_1, F_1 the points of contact of the escribed circle, which touches BC and the other sides produced: a, b, c denote the lengths of the sides BC, CA, AB ; s the semi-perimeter of the triangle, and r, r_1 the radii of the inscribed and escribed circles.

Prove the following equalities.

$$\begin{aligned} \text{(i)} \quad & AE = AF = s - a, \\ & BD = BF = s - b, \\ & CD = CE = s - c. \end{aligned}$$


$$\text{(ii)} \quad AE_1 = AF_1 = s.$$

$$\begin{aligned} \text{(iii)} \quad & CD_1 = CE_1 = s - b, \\ & BD_1 = BF_1 = s - c. \end{aligned}$$


$$\text{(iv)} \quad CD = BD_1 \text{ and } BD = CD_1$$

$$\text{(v)} \quad EE_1 = FF_1 = r.$$

$$\begin{aligned} \text{(vi)} \quad & \text{The area of the } \triangle ABC \\ & = rs = r_1(s - a). \end{aligned}$$



2. In the triangle ABC , I is the centre of the inscribed circle, and I_1, I_2, I_3 the centres of the escribed circles touching respectively the sides BC, CA, AB and the other sides produced.



Prove the following properties :-

- (i) The points A, I, I_1 are collinear; so are B, I, I_2 ; and C, I, I_3 .
- (ii) The points I_2, A, I_3 are collinear; so are I_3, B, I_1 ; and I_1, C, I_2 .
- (iii) The triangles BI_1C, CI_2A, AI_3B are equiangular to one another.
- (iv) The triangle $I_1I_2I_3$ is equiangular to the triangle formed by joining the points of contact of the inscribed circle.
- (v) Of the four points I, I_1, I_2, I_3 each is the orthocentre of the triangle whose vertices are the other three.
- (vi) The four circles, each of which passes through three of the points I, I_1, I_2, I_3 , are all equal.

3. With the notation of page 277, shew that in a triangle ABC , if the angle at C is a right angle,

$$r = s - c; \quad r_1 = s - b; \quad r_2 = s - a; \quad r_3 = s.$$

4. With the figure given on page 278, shew that if the circles whose centres are I, I_1, I_2, I_3 touch BC at D, D_1, D_2, D_3 , then

$$(i) \quad DD_2 - D_1D_3 = b.$$

$$(ii) \quad DD_1 - D_1D_2 = c.$$

$$(iii) \quad D_2D_3 = b + c.$$

$$(iv) \quad DD_1 = b + c.$$

5. Shew that the orthocentre and vertices of a triangle are the centres of the inscribed and escribed circles of the pedal triangle.

[See Ex. 20, p. 225.]

6. Given the base and vertical angle of a triangle, find the locus of the centre of the inscribed circle.

[See Ex. 36, p. 228.]

7. Given the base and vertical angle of a triangle, find the locus of the centre of the escribed circle which touches the base.

8. Given the base and vertical angle of a triangle, shew that the centre of the circumscribed circle is fixed.

9. Given the base BC , and the vertical angle A of a triangle, find the locus of the centre of the escribed circle which touches AC .

10. Given the base, the vertical angle, and the radius of the inscribed circle; construct the triangle.

11. Given the base, the vertical angle, and the radius of the escribed circle, (i) which touches the base, (ii) which touches one of the sides containing the given angle; construct the triangle.

12. Given the base, the vertical angle, and the point of contact with the base of the inscribed circle; construct the triangle.

13. Given the base, the vertical angle, and the point of contact with the base, or base produced, of an escribed circle; construct the triangle.

14. From an external point A two tangents AB, AC are drawn to a given circle; and the angle BAC is bisected by a straight line which meets the circumference in I and I_1 ; shew that I is the centre of the circle inscribed in the triangle ABC , and I_1 the centre of one of the escribed circles.

15. I is the centre of the circle inscribed in a triangle, and I_1, I_2, I_3 the centres of the escribed circles; shew that II_1, II_2, II_3 are bisected by the circumference of the circumscribed circle.

16. ABC is a triangle, and I_2, I_3 the centres of the escribed circles which touch AC , and AB respectively; shew that the points B, C, I_2, I_3 lie upon a circle whose centre is on the circumference of the circle circumscribed about ABC .

17. With three given points as centres describe three circles touching one another two by two. How many solutions will there be?

18. Two tangents AB , AC are drawn to a given circle from an external point A ; and in AB , AC two points D and E are taken so that DE is equal to the sum of DB and EC : shew that DE touches the circle.

19. Given the perimeter of a triangle, and one angle in magnitude and position: shew that the opposite side always touches a fixed circle.

20. Given the centres of the three escribed circles; construct the triangle.

21. Given the centre of the inscribed circle, and the centres of two escribed circles; construct the triangle.

22. Given the vertical angle, perimeter, and the length of the bisector of the vertical angle; construct the triangle.

23. Given the vertical angle, perimeter, and altitude; construct the triangle.

24. Given the vertical angle, perimeter, and radius of the inscribed circle; construct the triangle.

25. Given the vertical angle, the radius of the inscribed circle, and the length of the perpendicular from the vertex to the base; construct the triangle.

26. Given the base, the difference of the sides containing the vertical angle, and the radius of the inscribed circle; construct the triangle. [See Ex. 16, p. 258.]

27. Given a vertex, the centre of the circumscribed circle, and the centre of the inscribed circle, construct the triangle.

28. In a triangle ABC , I is the centre of the inscribed circle; shew that the centres of the circles circumscribed about the triangles BIC , CIA , AIB lie on the circumference of the circle circumscribed about the given triangle.

29. In a triangle ABC , the inscribed circle touches the base BC at D ; and r , r_1 are the radii of the inscribed circle and of the escribed circle which touches BC : shew that $r \cdot r_1 = BD \cdot DC$.

30. ABC is a triangle, D , E , F the points of contact of its inscribed circle; and $D'E'F'$ is the pedal triangle of the triangle DEF : shew that the sides of the triangle $D'E'F'$ are parallel to those of ABC .

31. In a triangle ABC , the inscribed circle touches BC at D . Shew that the circles inscribed in the triangles ABD , ACD touch one another.

ON THE NINE-POINTS CIRCLE.

32. In any triangle the middle points of the sides, the feet of the perpendiculars drawn from the vertices to the opposite sides, and the middle points of the lines joining the orthocentre to the vertices are concyclic.

In the $\triangle ABC$, let X, Y, Z be the middle points of the sides BC, CA, AB ; let D, E, F be the feet of the perps drawn to these sides from A, B, C ; let O be the orthocentre, and α, β, γ the middle points of OA, OB, OC :

then shall the nine points $X, Y, Z, D, E, F, \alpha, \beta, \gamma$ be concyclic.

Join $XY, XZ, X\alpha, Ya, Za$.

Now from the $\triangle ABO$, since $AZ = ZB$,
and $A\alpha = \alpha C$, *Hyp.*

$\therefore Z\alpha$ is par^l to BO . Ex. 2, p. 96.
And from the $\triangle ABC$, since $BZ = ZA$,
and $BX = XC$, *Hyp.*

$\therefore ZX$ is par^l to AC .

But BO makes a rt. angle with AC ;

\therefore the $\angle XZ\alpha$ is a rt. angle. *Hyp.*

Similarly, the $\angle XY\alpha$ is a rt. angle. I. 29.

\therefore the points X, Z, α, Y are concyclic:

that is, α lies on the C^{ce} of the circle, which passes through X, Y, Z ; and $X\alpha$ is a diameter of this circle.


Similarly it may be shewn that β and γ lie on the C^{ce} of the circle which passes through X, Y, Z .

Again, since αDX is a rt. angle, *Hyp.*

\therefore the circle on $X\alpha$ as diameter passes through D .

Similarly it may be shewn that E and F lie on the circumference of the same circle.

\therefore the points $X, Y, Z, D, E, F, \alpha, \beta, \gamma$ are concyclic. Q.E.D.



From this property the circle which passes through the middle points of the sides of a triangle is called the **Nine-Points Circle**; many of its properties may be derived from the fact of its being the circle circumscribed about the pedal triangle.

33. To prove that

- (i) the centre of the nine-points circle is the middle point of the straight line which joins the orthocentre to the circumscribed centre:
- (ii) the radius of the nine-points circle is half the radius of the circumscribed circle:
- (iii) the centroid is collinear with the circumscribed centre, the nine-points centre, and the orthocentre.

In the $\triangle ABC$, let X, Y, Z be the middle points of the sides; D, E, F the feet of the perp.; O the orthocentre; S and N the centres of the circumscribed and nine-points circles respectively.

(i) To prove that N is the middle point of SO .

It may be shewn that the perp. to XD from its middle point bisects SO ; Ex. 14, p. 98.

Similarly the perp. to EY at its middle point bisects SO :

that is, these perp. intersect at the middle point of SO :

And since XD and EY are chords of the nine-points circle, \therefore the intersection of the lines which bisect XD and EY at rt. angles is its centre: III. 1.

\therefore the centre N is the middle point of SO .

(ii) To prove that the radius of the nine-points circle is half the radius of the circumscribed circle.

By the last Proposition, Xa is a diameter of the nine-points circle. \therefore the middle point of Xa is its centre:

but the middle point of SO is also the centre of the nine-points circle. (Proved.)

Hence Xa and SO bisect one another at N .

Then from the $\triangle SNX, ONa$

Because $\begin{cases} SN = ON, \\ \text{and } NX = Na, \\ \text{and the } \angle SNX = \text{the } \angle ONa; \end{cases}$ I. 15.

$\therefore SX = Oa$ I. 4.

$= Aa.$


And SX is also par^l to Aa ,

$\therefore SA = Xa.$ I. 33.

But SA is a radius of the circumscribed circle;

and Xa is a diameter of the nine-points circle;

\therefore the radius of the nine-points circle is half the radius of the circumscribed circle.



(in) To prove that the centroid is collinear with points S, N, O .

Join AX and draw ag par^l to SO .

Let AX meet SO at G .

Then from the $\triangle AGO$, since $Aa = aO$ and ag is par^l to OG ,

$$\therefore Ag = gG. \quad \text{Ex. 13, p. 98.}$$

And from the $\triangle Xag$, since $aN = NX$, and NG is par^l to ag ,

$$\therefore gG = GX. \quad \text{Ex. 13, p. 98.}$$

$$AG = \frac{2}{3} \text{ of } AX;$$

$\therefore G$ is the centroid of the triangle ABC .

That is, the centroid is collinear with the points S, N, O . \square E.D.

34. Given the base and vertical angle of a triangle, find the locus of the centre of the nine-points circle.

35. The nine-points circle of any triangle ABC , whose orthocentre is O , is also the nine-points circle of each of the triangles AOB, BOC, COA .

36. If I, I_1, I_2, I_3 are the centres of the inscribed and escribed circles of a triangle ABC , then the circle circumscribed about ABC is the nine-points circle of each of the four triangles formed by joining three of the points I, I_1, I_2, I_3 .

37. All triangles which have the same orthocentre and the same circumscribed circle, have also the same nine points circle.

38. Given the base and vertical angle of a triangle, shew that one angle and one side of the pedal triangle are constant.

39. Given the base and vertical angle of a triangle, find the locus of the centre of the circle which passes through the three escribed centres.

NOTE For another important property of the Nine-points Circle see Ex. 60, p. 382.

II. MISCELLANEOUS EXAMPLES.

1. If four circles are described to touch every three sides of a quadrilateral, shew that their centres are concyclic.

2. If the straight lines which bisect the angles of a rectilineal figure are concurrent, a circle may be inscribed in the figure.

3. Within a given circle describe three equal circles touching one another and the given circle.

4. The perpendiculars drawn from the centres of the three escribed circles of a triangle to the sides which they touch are concurrent.

APPENDIX.

ON POLE AND POLAR

DEFINITIONS.


(1) If in any straight line drawn from the centre of a circle two points are taken such that the rectangle contained by their distances from the centre is equal to the square on the radius, each point is said to be the **inverse** of the other.

Thus in the figure given below, if O is the centre of the circle, and if $OP \cdot OQ = (\text{radius})^2$, then each of the points P and Q is the inverse of the other.

It is clear that if one of these points is within the circle the other must be without it.

(ii) The **polar** of a given point with respect to a given circle is the straight line drawn through the inverse of the given point at right angles to the line which joins the given point to the centre: and with reference to the polar the given point is called the **pole**.

Thus in the adjoining figure, if $OP \cdot OQ = (\text{radius})^2$, and if through



P and Q , LM and HK are drawn perp. to OP ; then HK is the polar of the point P , and P is the pole of the st. line HK : also LM is the polar of the point Q , and Q the pole of LM .

It is clear that the polar of an *external* point must intersect the circle, and that the polar of an *internal* point must fall without it: also that the polar of a point *on the circumference* is the tangent at that point.


1. Now it has been proved [see Ex. 1, page 233] that if from an external point P two tangents PH , PK are drawn to a circle, of which O is the centre, then OP cuts the chord of contact HK at right angles at Q , so that

$$OP \cdot OQ = (\text{radius})^2,$$

$\therefore HK$ is the polar of P with respect to the circle. Def. (ii).

Hence we conclude that

The Polar of an external point with reference to a circle is the chord of contact of tangents drawn from the given point to the circle.



The following Theorem is known as the **Reciprocal Property of Pole and Polar**.

2. *If A and P are any two points, and if the polar of A , with respect to any circle passes through P , then the polar of P must pass through A .*

Let BC be the polar of the point A with respect to a circle whose centre is O , and let BC pass through P : then shall the polar of P pass through A .

Join OP ; and from A draw AQ perp. to OP . We shall shew that AQ is the polar of P .

Now since BC is the polar of A ,

\therefore the $\angle ABP$ is a rt. angle;

Def. (ii), p. i.

and the $\angle AQP$ is a rt. angle: Constr.

\therefore the four points A, B, P, Q are concyclic;

$\therefore OQ \cdot OP = OA \cdot OB$ III. 36.

(radius)², for CB is the polar of A :


$\therefore P$ and Q are inverse points with respect to the given circle.

And since AQ is perp. to OP ,

$\therefore AQ$ is the polar of P .

That is, the polar of P passes through A .

Q. E. D.



A similar proof applies to the case when the given point A is without the circle, and the polar BC cuts it.

3. To prove that the locus of the intersection of tangents drawn to a circle at the extremities of all chords which pass through a given point is the polar of that point.

Let A be the given point within the circle, of which O is the centre.

Let HK be any chord passing through A ; and let the tangents at H and K intersect at P :

it is required to prove that the locus of P is the polar of the point A .

I. To shew that P lies on the polar of A .

Join OP cutting HK in Q .
Join OA : and in OA produced take the point B ,

so that $OA \cdot OB = (\text{radius})^2$. II. 11.
Then since A is fixed, B is also fixed.

Join PB .

Then since HK is the chord of contact of tangents from P ,

$\therefore OP \cdot OQ = (\text{radius})^2$. Ex. 1, p. 233.

But $OA \cdot OB = (\text{radius})^2$;

$\therefore OP \cdot OQ = OA \cdot OB$:

the four points A, B, P, Q are concyclic.

the \angle at Q and B together = two rt. angles. III. 22.

But the \angle at Q is a rt. angle;

\therefore the \angle at B is a rt. angle.

And since the point B is the inverse of A ;

$\therefore PB$ is the polar of A ;

that is, the point P lies on the polar of A .

II. To shew that any point on the polar of A satisfies the given conditions.

Let BC be the polar of A , and let P be any point on it. Draw tangents PH, PK , and let HK be the chord of contact.

Now from Ex. 1, p. ii, we know that the chord of contact HK is the polar of P ,

and we also know that the polar of P must pass through A ; for P is on BC , the polar of A :

that is, HK passes through A .

$\therefore P$ is the point of intersection of tangents drawn at the extremities of a chord passing through A .

From I. and II. we conclude that the required locus is the polar of A .

NOTE. If A is *without* the circle, the theorem demonstrated in Part I. of the above proof still holds good; but the converse theorem in Part II. is not true for *all* points in BC . For if A is without the

circle, the polar BC will intersect it: and no point on that part of the polar which is within the circle can be the point of intersection of tangents.

We now see that

(i) *The Polar of an external point with respect to a circle is the chord of contact of tangents drawn from it.*

(ii) *The Polar of an internal point is the locus of the intersections of tangents drawn at the extremities of all chords which pass through it.*

(iii) *The polar of a point on the circumference is the tangent at that point.*

EXAMPLES ON POLE AND POLAR.

1. *The straight line which joins any two points is the polar with respect to a given circle of the point of intersection of their polars.*

2. *The point of intersection of any two straight lines is the pole of the straight line which joins their poles.*

3. *Find the locus of the poles of all straight lines which pass through a given point.*

4. *Find the locus of the poles, with respect to a given circle, of tangents drawn to a concentric circle.*

5. *If two circles cut one another orthogonally and PQ be any diameter of one of them; shew that the polar of P with regard to the other circle passes through Q .*

6. *If two circles cut one another orthogonally, the centre of each circle is the pole of their common chord with respect to the other circle.*

7. *Any two points subtend at the centre of a circle an angle equal to one of the angles formed by the polars of the given points.*

8. *O is the centre of a given circle, and AB a fixed straight line. P is any point in AB ; find the locus of the point inverse to P with respect to the circle.*

9. *Given a circle, and a fixed point O on its circumference: P is any point on the circle: find the locus of the point inverse to P with respect to any circle whose centre is O .*

10. *Given two points A and B , and a circle whose centre is O ; shew that the rectangle contained by OA and the perpendicular from B on the polar of A is equal to the rectangle contained by OB and the perpendicular from A on the polar of B .*

II. ON THE RADICAL AXIS.

1. To find the locus of points from which the tangents drawn to two given circles are equal.

Fig. 1.





Fig. 2.



Let A and B be the centres of the given circles, whose radii are a and b ; and let P be any point such that the tangent PQ drawn to circle (A) is equal to the tangent PR drawn to the circle (B):

it is required to find the locus of P.

Join PA, PB, AQ, BR, AB; and from P draw PS perp. to AB.

Then because $PQ = PR$, $\therefore PQ^2 = PR^2$.

But $PQ^2 = PA^2 - AQ^2$; and $PR^2 = PB^2 - BR^2$; I. 47.

$\therefore PA^2 - AQ^2 = PB^2 - BR^2$;

that is, $PS^2 + AS^2 - a^2 = PS^2 + SB^2 - b^2$; I. 17.

or, $AS^2 - a^2 = SB^2 - b^2$.

Hence AB is divided at S, so that $AS^2 - SB^2 = a^2 - b^2$:

\therefore S is a fixed point.

Hence all points from which equal tangents can be drawn to the two circles lie on the straight line which cuts AB at rt. angles, so that the difference of the squares on the segments of AB is equal to the difference of the squares on the radii.


Again, by simply retracing these steps, it may be shewn that in Fig. 1 every point in SP, and in Fig. 2 every point in SP exterior to the circles, is such that tangents drawn from it to the two circles are equal.

Hence we conclude that in Fig. 1 the whole line SP is the required locus, and in Fig. 2 that part of SP which is without the circles.

In either case SP is said to be the **Radical Axis** of the two circles.

COROLLARY. *If the circles cut one another as in Fig. 2, it is clear that the Radical Axis is identical with the straight line which passes through the points of intersection of the circles; for it follows readily from III. 36 that tangents drawn to two intersecting circles from any point in the common chord produced are equal.*

2. *The Radical Axes of three circles taken in pairs are concurrent.*



Let there be three circles whose centres are A, B, C.

Let CZ be the radical axis of the \odot^s (A) and (B);
and OY the Radical Axis of the \odot^s (A) and (C), O being the point of intersection:

then shall the radical axis of the \odot^s (B) and (C) pass through O.

It will be found that the point O is either *without* or *within* all the circles.

I. When O is without the circles.

From O draw OP, OQ, OR tangents to the \odot^s (A), (B), (C).

Then because O is a point on the radical axis of (A) and (B); *Hyp.*

$$\therefore OP = OQ.$$

And because O is a point on the radical axis of (A) and (C), *Hyp.*

$$\therefore OP = OR,$$

$$\therefore OQ = OR;$$


\therefore O is a point on the radical axis of (B) and (C),
i.e. the radical axis of (B) and (C) passes through O.

II. If the circles intersect in such a way that O is within them all;

the radical axes are then the common chords of the three circles taken two and two; and it is required to prove that these common chords are concurrent. This may be shewn indirectly by III. 35.

DEFINITION. The point of intersection of the radical axes of three circles taken in pairs is called the **radical centre**.

3. To draw the radical axis of two given circles.



Let A and B be the centres of the given circles:
it is required to draw their radical axis.

If the given circles intersect, then the st. line drawn through their points of intersection will be the radical axis. [Ex. 1, Cor. p. vi.]

But if the given circles do not intersect,

describe any circle so as to cut them in E, F and G, H:

Join EF and HG, and produce them to meet in P.

Join AB; and from P draw PS perp. to AB.

Then PS shall be the radical axis of the \odot s (A), (B).

DEFINITION. If each pair of circles in a given system have the same radical axis, the circles are said to be **co-axal**.

EXAMPLES.

1. Show that the radical axis of two circles bisects any one of their common tangents.

2. If tangents are drawn to two circles from any point on their radical axis; shew that a circle described with this point as centre and any one of the tangents as radius, cuts both the given circles orthogonally.

3. O is the radical centre of three circles, and from O a tangent OT is drawn to any one of them: shew that a circle whose centre is O and radius OT cuts all the given circles orthogonally.

4. If three circles touch one another, taken two and two, shew that their common tangents at the points of contact are concurrent.

5. If circles are described on the three sides of a triangle as diameters, their radical centre is the orthocentre of the triangle.

6. All circles which pass through a fixed point and cut a given circle orthogonally, pass through a second fixed point.

7. Find the locus of the centres of all circles which pass through a given point and cut a given circle orthogonally.

8. Describe a circle to pass through two given points and cut a given circle orthogonally.

9. Find the locus of the centres of all circles which cut two given circles orthogonally.

10. Describe a circle to pass through a given point and cut two given circles orthogonally.

11. The difference of the squares on the tangents drawn from a point to two circles is equal to twice the rectangle contained by the straight line joining their centres and the perpendicular from the given point on their radical axis.

12. In a system of co-axial circles which do not intersect, any point is taken on the radical axis; shew that a circle described from the point as centre with radius equal to the tangent drawn from it to any one of the circles, will meet the line of centres in two fixed points.

[These fixed points are called the **Limiting Points** of the system.]

